

Comparative Evaluation of Nutritional Qualities of *Nymphaea lotus* and *Nymphaea pubescens* Seeds

ABSTRACT

Aims: The study is aimed at evaluating the nutritional qualities of *Nymphaea lotus* and *Nymphaea pubescens* seeds.

Study design: It was designed to estimate the proximate and amino acid profiles of *Nymphaea lotus* and *Nymphaea pubescens* seeds to indicate their nutritional potentials.

Place and Duration of Study: Department of Biochemistry, Faculty of Basic Medical Sciences, Bayero University, Kano State-Nigeria, between August 2016 and January 2017.

Methodology: Fresh seeds were collected, dried and ground to smaller particle size. Standard official methods were employed in the proximate analysis while amino acid analysis was carried out using Technicon Sequential Multisample Amino Acid Analyser.

Results: The result of the proximate composition revealed the richness of both seeds in carbohydrates, crude lipids and proteins. The amino acid analysis revealed that amino acids were more concentrated in *N. lotus* seed with the total value of 73.82g/100g than 70.70g/100g of *N. pubescens* seed. The former also exceeded the latter in total essential amino acids by difference of 1.12g/100g. The highest scoring amino acid was arginine in *N. lotus* seed and leucine in *N. pubescens* seed while methionine was the most limiting amino acid in both the samples. All the essential amino acids in both samples satisfied WHO/FAO reference protein except histidine, valine and methionine. The P-PER of *Nymphaea lotus* seed and *Nymphaea pubescens* seed were found to be 1.69 and 1.54 respectively, meaning *Nymphaea pubescens* seed may be slightly more bioavailable than the *Nymphaea lotus* seed. The Lys/Arg ratio of *Nymphaea lotus* seed (0.84) and *Nymphaea pubescens* seed (1.09) indicate their low atherogenic potential.

Conclusion: These results re-enforce the growing awareness that wild and semi-wild plant seeds can contribute useful amounts of essential nutrients to human diets.

Keywords: *Nymphaea lotus*, *Nymphaea pubescens*, Amino acid, Essential amino acid, Protein

1. INTRODUCTION

Malnutrition is a condition that results from eating a diet in which nutrients are either not enough or are too much such that the diet causes health problem. It may involve calories, protein, carbohydrates, vitamins and minerals [1]. Not enough nutrients is called undernutrition or undernourishment while too much nutrients is called overnutrition. Malnutrition is often specifically to refer to undernutrition where there is not enough calories, protein and micronutrients [2]. There are number of causes of malnutrition. It results from: inadequate or unbalanced diet, problems with digestion or absorption and certain medical conditions.

Nigeria has been ranked second after India in the list of countries with highest cases of malnourished children in the world as it accounts for 10% of the 160 million stunted children globally. The malnutrition in

Nigeria is not very encouraging as about 37% of the children are stunted, children under the age of five have chronic malnutrition and 29% are underweight, while 18% have acute malnutrition [3]. However, protein energy malnutrition affects children the most because they have less protein intake. The few rare case found in the world are almost entirely found in children as a result of diets or ignorance in the nutritional needs of children particularly in the cases of milk [4].

Nymphaea lotus (white waterlily) and *Nymphaea pubescens* (red waterlily) also known as *Bado gero* and *Bado dawa* in the native Hausa language, are perennial plant that grows up to 45cm in height. They are herbaceous aquatic plants, whose leaves float or are submerged in water [5]. These plants prefer clear, warm and slightly acidic water and is localized to central and Southern Europe, Asia, the Middle East, Northern Africa, and Tropical Mountain of West Africa especially Nigeria [6]. *Nymphaea lotus* and *Nymphaea pubescens* fruits are either eaten raw or cooked for food when dried by local people of Northern Nigeria. This study evaluated and compared the nutritional qualities of *Nymphaea lotus* and *Nymphaea pubescens* seeds

2. METHODOLOGY

2.1 COLLECTION AND IDENTIFICATION OF PLANT MATERIALS

Nymphaea lotus and *Nymphaea pubescens* seeds were obtained from Guzu- Guzu Dam in Kabo Local Government Area, Kano state of Nigeria. The samples were authenticated by the Department of Plant Biology, Bayero University Kano with accession number BUKHAN 0356 and BUKHAN 0357 for *Nymphaea lotus* and *Nymphaea pubescens* seeds respectively.

2.2 PREPARATION OF SAMPLES

The samples were thoroughly washed to remove sand and the drained parts were air dried later. The samples were grounded using wooden mortar and pestle until powder was obtained to ensure homogeneity. The fine powdered samples were then stored at 25°C into labeled plastic containers prior to use.

2.3 PROXIMATE ANALYSIS

2.3.1 Determination of Moisture Content [7]

Three clean dried petri dishes were weighed (W_1) and 5g of the pulverized sample was placed in each of them and weighed (W_2). They were placed in an oven at 100°C for 17 hours. The dishes were removed and cooled in a dessicator for 30minutes and finally weighed (W_3).

Calculation

The loss in weight due to drying is equal to the moisture content of the sample,

$$\% \text{moisture} = \frac{\text{loss of weight}}{\text{weight of sample taken}} \times 100$$

$$= \frac{W_2 - W_3}{W_2 - W_1} \times 100$$

Where

W_1 = weight of empty petri dish

W_2 = weight of empty petri dish + sample

W_3 = weight of empty petri dish + sample after drying

2.3.2 Determination of Ash Content [8].

Porcelain crucible or silica disc was ignited in a hot Bunsen burner flame or in a muffle furnace for about one minute; it was then transferred into a dessicator for cooling and weighing (W_1). The sample (2g) was placed into the crucible and weighed again (W_2). A crucible containing the sample was heated gently on a muffle furnace at 550-570°C to burn off all the organic matter. The carbon char burn off as CO_2 leaving a white ash. The crucible taken out was immediately covered and placed in a dessicator for cooling and then weighed (W_3).

Calculation

$$\text{Ash \%} = \frac{W_3 - W_1}{W_2 - W_1} \times 100$$

Where

W_1 = weight of crucible

W_2 = weight of crucible + sample

W_3 = weight of crucible after all organic matter was burnt off

2.3.3 Determination of Crude Fat using soxhlet extraction method [7].

Exactly 3g of the sample was carefully weighed (W_1) into a folded fat-free filter paper. This was properly tied with a thread at both ends and weighed (W_2). This was carefully placed in the extraction thimble and small cotton wool placed on top. The whole apparatus was then connected after the addition of 300ml of hexane into the weighed extraction flask. The extraction was then carried out for 3 hours using the heating mantle at 60-80°C making sure there was continuous flow of water in the condenser (which served as the cooling system). The sample was then removed, air-dried and then placed in an oven at 80°C until a constant weight was obtained (W_3).

78 Calculation

$$79 \text{ Crude fat (\%)} = \frac{(W_2 - W_3)}{W_1} \times 100$$

80 Where:

81 W_1 = Weight of sample

82 W_2 = Weight of sample and filter paper

83 W_3 = Weight of sample after lipid extraction and filter paper

84 **2.3.4 Determination of Crude protein [7].**

85 Exactly 0.15g of dried (moisture free) sample was weighed and transferred into the Kjeldahl digestion
86 flask (30-35ml). 0.8g of catalyst (0.7g sodium sulphate, 0.06g copper sulphate and 0.04g mercury (II)
87 oxide red) was added into the digestion flask. To this mixture was added 2ml of concentrated sulphuric
88 acid of which color turned black. The mixture in the digestion flask was heated on the heating mantle for 1
89 hour until liquid became clear green. The digest was cooled and made alkaline with 15ml 40% NaOH.
90 The digest was then transferred to the steamed out apparatus using minimum volume of water. The
91 ammonia steamed is distilled into 10ml 2% boric acid solution with 5 drops of methyl red indicator for 15
92 minutes. The distilled ammonia was titrated with 0.02M hydrochloric acid. The same procedure was
93 carried out for the blank.

94 Calculation

$$95 \% \text{Nitrogen} = \frac{0.00056 \times V \times 100}{W}$$

96 Where:

97 V = volume of acid used (titre value)

98 W = weight of sample = 0.15g

99 %Crude protein = %Nitrogen $\times$ 6.25 (multiplication factor)

100

101 **2.3.5 Determination of crude fiber content [7].**

102 Exactly 3g of the sample was weighed "W" into the extraction apparatus and extracted three times with
103 light petroleum ether by stirring, settling and decanting. The air-dried extracted sample was transferred to
104 a dry 100ml conical flask. 200ml of 0.127M H_2SO_4 measured at ordinary temperature and brought to its
105 boiling point. The sample was then dispersed by some of this solution. This was boiled for 30mins, while
106 a constant volume was maintained. The flask was rotated every few minutes in order to mix the contents
107 and remove particles from the side.

108 The Buchner funnel was prepared while waiting for the mixture to boil by fixing the Buchner funnel to a
109 perforated plate and to the funnel and ash free filter paper was also fixed to cover the holes in the plate.
110 Then the boiled water was poured into the funnel and allowed to remain until the funnel was hot and then
111 drained by Applying suction.

At the end of 30minutes of boiling, the mixture was allowed to stand for 1minute and then, the mixture was poured immediately into a shallow layer of hot water under gentle suction in the prepared funnel. The suction was adjusted so that filtration of the bulk of 200ml was completed within 10minutes.

The insoluble matter was washed with boiled water for several times until the washings were free of acid. It was washed back into the original flask by means of a wash bottle containing 200ml of 0.313 NaOH solution measured at ordinary temperature and brought to boiling point. The mixture was boiled for 30minutes with the same precautions as those used in the earlier boiling acid treatment. After 30minutes of boiling, it was allowed to stand for 1minute and then filtered immediately. The insoluble material was transferred to the filter paper by means of boiling water, washed with 1%HCl and finally, it was washed again with boiling water until it was free from acid. This was then washed twice with alcohol (ethanol) and thrice with petroleum ether. The insoluble matter was then transferred to a dried weighed crucible and dried at 100⁰C to a constant weight. The crucible and its content were placed on a heating mantle in a fume cupboard to burn off the organic matter. It was transferred to a muffle furnace at 550⁰C for 3hours. The ash content after cooling was then determined by weighing.

Calculation

$$\text{Crude fiber (\%)} = \frac{W_1 - W_2}{\text{Weight of sample}} \times 100$$

Where:

W_1 = Weight of sample extracted + filter paper

W_2 = Weight of "W" after ashing.

2.3.6 Estimation of Total Carbohydrates

Total carbohydrate was estimated by calculation/ difference [8].

Calculation

% Carbohydrates= 100- (% Crude Fat + % Crude Protein + % Ash + % Crude fibre).

2.3 ANALYSIS OF AMINO ACID PROFILE OF *Nymphaea lotus* AND *Nymphaea pubescens* SEEDS

The amino acid profile of the *Nymphaea lotus* and *Nymphaea pubescens* seeds were determined using methods described by [9]. The samples were dried to constant weight, defatted, hydrolysed, evaporated in a rotary evaporator and then loaded into the Technicon Sequential Multisample Amino Acid Analyser (TSM). (Technicon Instruments Corporation, New York).

2.3.1 Defatting of sample

A known weight of the dried sample was weighed into extraction thimble and the fat was extracted with chloroform/methanol mixture in ratio of 2: 1 using soxhlet extraction apparatus as described by [7]. The extraction lasted for 15 hours.

2.3.2 Percentage Nitrogen Determination

The sample (2g) was weighed, wrapped in whatman filter paper (No.1) and put in the Kjeldhal digestion flask. Concentrated sulphuric acid (10ml) was added. Catalyst mixture (0.5) containing sodium sulphate (Na_2SO_4), copper sulphate (CuSO_4) and selenium oxide (SeO_2) in the ration of 10:5:1 was added into the flask to facilitate digestion. Four pieces of anti-bumping granules were added. The flask was then placed in Kjeldhal digestion apparatus for 3 hours until the liquid turned light green. The digested sample was cooled and diluted with distilled water to 100ml in standard volumetric flask. Aliquot (10ml) of the diluted solution with 10ml of 45% sodium hydroxide was placed into the Markham distillation apparatus and distilled with 10ml of 2% boric acid containing 4 drops of methyl red indicator until 70ml of distillate was collected. The distillate was then titrated with standardize 0.01N hydrochloric acid to grey coloured end point, the percentage nitrogen in the original sample was calculated using the formula:

$$\% \text{ N} = \frac{(a-b) \times 0.01 \times 14 \times v}{w \times c} \times 100$$

Where:

- a = Titre value of the digested sample
- b = Titre value of blank sample
- v = Volume after dilution (100ml)
- w = Weight of dried sample (g)
- c = Aliquot of the sample used (10ml)
- 14 = Nitrogen constant in g

2.3.3 Hydrolysis of the samples.

The defatted sample (1.204g) was weighed into glass ampoule. 7 ml of 6N HCl was added and oxygen was expelled by passing nitrogen into the ampoule (This is to avoid possible oxidation of some amino acids during hydrolysis). The glass ampoule was then sealed with Bunsen burner flame and placed in an oven preset at $105^\circ\text{C} \pm 5^\circ\text{C}$ for 22 hours to effect hydrolysis. The ampoule was allowed to cool before it was broken open at the tip and the content was filtered to remove the humins. The filtrate was then evaporated to dryness at 40°C under vacuum in a rotary evaporator. The residue was dissolved with 5 ml of acetate buffer (pH 2.0) and stored in plastic specimen bottles, which were kept in deep freezer.

2.3.4 Loading of the hydrolysate into the TSM analyzer.

The amount loaded was 60microlitres. This was injected into the cartridge of the analyzer. The TSM analyzer is designed to analyse free acidic, neutral and basic amino acids of the hydrolysate. The period of an analysis lasted for 75 min. The amino acids present in the samples were identified by matching their peak retention time in the chromatogram with those peaks of standard mixture of amino acids with norleucine as internal standard.

2.3.5 Calculating the amino acid values from the chromatogram peaks

An integrator attached to the analyzer calculates the peak area proportional to the concentration of each of the amino acids. The net heights of each peak produced by the chart recorder of the TSM (each representing an amino acid) were measured. The half-height of the peak on the chart was found and the width of the peak at the half- height was accurately measured and result recorded. Approximate area of each peak was then obtained by multiplying the height with the width at half height. The norleucine equivalent (NE) for each amino acid in the standard mixture was calculated using the formular:

$$NE = \frac{(\text{area of norleucine peak})}{(\text{area of each amino acid})}$$

Constant S was calculated for each amino acid in the standard mixture:

$$S_{std.} = NE_{std.} \times \text{Mol. weight} \times \mu\text{Mol. AA}_{std.}$$

Finally the amount of each amino acid present in the samples was calculated in grams per 100 g protein using the following formula:

$$\text{Concentration (g/100 of protein)} = NH \times NH/2 \times S_{std.} \times C$$

$$\text{Where: } C = \frac{[\text{Dilution} \times 16]}{\text{Sample Wt(g)} \times \text{Vol.} \frac{\text{Loaded}}{NH} \times W(nleu)}$$

Where Wt= weight, NH = net height, W = width, nleu = norleucine

2.3.6 Determination of Tryptophan

Tryptophan is a difficult amino acid to determine in proteins and peptides because it chemically decomposes during acid hydrolysis. It should be noted that tryptophan is destroyed by 6N HCL during hydrolysis. Alkaline hydrolysis was improved by using sodium hydroxide (NaOH) instead of barium hydroxide to prevent problems with both precipitation and adsorption of tryptophan.

The tryptophan in the known sample was hydrolyzed with 4.2 M Sodium hydroxide. The known sample was dried to constant weight, defatted, hydrolyzed, evaporated in a rotary evaporator and loaded into the Applied Biosystems PTH Amino Acid Analyzer as described above.

2.3.7 Estimation of Seed Samples Protein Quality: To estimate the quality of dietary protein in the seed samples, the Total Essential Amino Acid (TEAA), Total Non-Essential Amino Acid (TNEAA), Total Sulphur Amino Acid (TSAA), percentage of cysteine in TSAA (%Cys/TSAA), Total Aromatic Amino Acid (TArAA), Leu/Ile ratio, Lys/Arg ratio etc, were calculated as described by FAO, (1995). The Predicted Protein Efficiency Ratio (P-PER) was also determined using the equation of Alsmeyer's (1973). I.e. P-PER = - 0.468 + 0.454 (Leu) – 0.105 (Tyr). The percentage amino acid scores of the seed samples were also calculated using the formula:

$$\% \text{ Amino acid score} = \frac{\text{Amount of amino acid per test protein (g/100g)}}{\text{Amount of amino acid per protein in FAO/WHO (1995) reference pattern (g/100g)}} \times 100$$

3. RESULTS AND DISCUSSION

3.1 PROXIMATE COMPOSITION

Table 1 presents the proximate composition of the *Nymphaea lotus* and *Nymphaea pubescens* seeds. The moisture, crude fibre contents of *Nymphaea lotus* and *Nymphaea pubescens* seeds are not significantly different (p>0.05). However, ash, lipid, crude proteins and carbohydrates contents of the *Nymphaea lotus* seed and *Nymphaea pubescens* seeds recorded significant difference (p<0.05).

Table 1: Proximate compositions of *Nymphaea lotus* seed and *Nymphaea pubescens* seeds

Parameters (%)	<i>Nymphaea lotus</i> seed	<i>Nymphaea pubescens</i> seed
Moisture	5.31±0.32	6.20±0.87
Ash	1.33±0.11 ^a	3.00±0.30 ^a
Proteins	4.92±0.34 ^a	4.14±0.06 ^a
Lipids	13.23±1.01 ^a	9.28±0.15 ^a
Fibre	5.17±0.51	5.00±0.26
Carbohydrates	75.35±1.38 ^a	78.58±0.89 ^a

All values are expressed as mean ± standard deviation for three determinations. Rows with similar superscript are statistically significant (p<0.05).

The percentage of ash, moisture, crude protein, crude fat (lipids), crude fibre and carbohydrate contents of *Nymphaea lotus* and *Nymphaea pubescens* seeds are presented in table 1. The *Nymphaea pubescens* seed had higher percentage value of ash content compared to *Nymphaealotus* seed. This is higher than

the reported value of *Nymphaea lotus* seed by [12][5]. Thus, the result indicated that *Nymphaea lotus* and *Nymphaea pubescens* seed could serve as good source of mineral element since ash content measured the minerals content of food substances and were found to be within the range reported for some leguminous seed [13]. However, the percentage for the moisture content in *Nymphaea lotus* and *Nymphaea pubescens* seeds were 5.31% and 6.20% respectively which is less than 8.5% in *Annona muricata* and 9.7 % in pride of Barbados seeds reported by [13][14] respectively. These results are comparable to the literature value of 6.00% reported for *Nymphaea lotus* seed [12]. However, the results were lower than 8.20% reported by [5]. The low moisture content of both *Nymphaea* seeds may signify longer shelf life since seeds with moisture content greater than 15% are subjected to deterioration from mould growth, heat, insect damage and sprouting [13]. Moreover, the percentage of crude fat (lipids) in *Nymphaea lotus* seed was found to be higher than in *Nymphaea pubescens* seeds. In addition [12] had earlier reported that *Nymphaea lotus* seeds have a lipid or crude fat content of 9.33% which was lower than value reported in the present study. Therefore, both *Nymphaea* seeds could serve as the secondary source of energy.

Furthermore, *Nymphaea lotus* seed was found to have higher crude protein content than the reported value of (3.09%) by [5]. It was earlier reported that *Nymphaea lotus* seed could be good source of protein [15]. The percentage value of crude protein obtained for *Nymphaea pubescens* seed was slightly lower than in *Nymphaea lotus* seed. Thus, this study suggests that *Nymphaea lotus* seeds possess high protein content compared to *Nymphaea pubescens* seeds. Proteins ensure proper growth and also serve as source of some essential amino acids for body use. Analysis revealed slightly higher percentage of crude fibre in *Nymphaea lotus* seed than obtained in the *Nymphaea pubescens* seeds. However these values are within the range reported by [15][5][12] for *Nymphaea lotus* seeds. Thus, ingestion of foodstuff which has reasonable amount of dietary fiber reduce postprandial glucose response after carbohydrate rich meals as well as lowering total LDL cholesterol levels [16].

Similarly, the carbohydrate content of both *Nymphaea lotus* and *Nymphaea pubescens* seeds were lower compared to the reported value in *Nymphaea lotus* seed (80.96%) by [12]. The difference in carbohydrates contents could be attributed to difference in climate and soil condition. The carbohydrate composition present in both *Nymphaea* seeds is enough to prevent ketosis as breakdown of 1gram of (glucose) carbohydrate yield 4 kcal of energy [17]. Similar carbohydrate content was reported by Mohammed *et al.* (2013) which reported that, seeds of *Nymphaea lotus* serves as good source of carbohydrate and is enough to prevent ketosis. The role of Carbohydrate is not limited to energy intake as they also play a vital role in both the taste of the food and the pleasure of eating [18].

Therefore, the proximate composition of the *Nymphaea lotus* and *Nymphaea pubescens* seeds indicates they are highly nutritious as they contains high protein content hence could supplement other protein sources such as beans, peas and groundnuts especially in dry seasons and in arid regions [19] when or where most common protein sources might be scarce and expensive, hence could help to reduce malnutrition and food insecurity

259 The amino acid composition and percentage Amino acid score of *Nymphaea lotus* and *Nymphaea*
 260 *pubescens* seeds are presented in Table 2. The result indicates the presence of all the essential amino
 261 acids while two non-essential amino acids were totally absent in the two samples screened.

262 3.2 AMINO ACIDS COMPOSITIONS

263 **Table 2: Amino acid composition and percentage Amino acid Score of *Nymphaea lotus* and**
 264 ***Nymphaea pubescens* seeds**

Amino Acid	Concentration(g/100g)		FAO/WHO Ref.	%Amino Acid Score	
	<i>N. lotus</i>	<i>N. pubescens</i>		<i>N. lotus</i>	<i>N. pubescens</i>
Leucine	5.49	5.14	4.2	130.71	122.38
Lysine	4.48	4.88	4.2	106.67	116.19
Isoleucine	4.52	4.19	4.2	107.62	99.76
Phenylalanine	2.48	2.39	2.3	107.83	103.91
Tryptophan	1.21	0.92	1.1	110.00	83.63
Valine	4.01	3.71	4.2	95.48	88.33
Methionine	1.23	1.20	2.2	55.91	54.55
Arginine	5.33	4.47	4.0	133.25	111.75
Histidine	2.43	2.20	3.4	71.47	64.71
Threonine	3.16	3.25	2.8	112.86	116.71
Cysteine	1.15	1.27	2.0	57.50	63.50
Proline	3.55	3.96	-	-	-
Tyrosine	3.10	3.10	2.8	110.71	110.71
Alanine	4.02	3.41	-	-	-
Glutamic acid	12.11	11.58	6.3	192.22	183.81

Glycine	3.61	3.94	-	-	-
Serine	3.94	3.59	-	-	-
Aspartic acid	8.00	7.50	-	-	-

265 - Means Not reported/Not determined

266 From the various parameters presented in Table 2, total amino acids (TAA), total non-essential amino
 267 acids (TNEAA), total essential amino acids (TEAA), their respective percentages, the predicted protein
 268 efficiency ratio (P-PER), Leu/Ile ratio and Essential amino acid index (EAAI) values of *Nymphaea lotus*
 269 and *Nymphaea pubescens* seed were calculated and presented in Table 3.

270 **Table 3: Nutritional quality indices of protein**

Amino Acid Grouping	<i>Nymphaea lotus</i> Seed	<i>Nymphaea pubescens</i> Seed
Total amino acid (TAA)	73.82	70.70
Total non-essential amino acid (TNEAA)	39.48	38.38
Total essential amino acid (TEAA)		
-with His	34.34	32.32
-without His	31.91	30.12
% TNEAA	53.48	54.29
% TEAA		
-with His	46.52	45.71
-without His	43.23	42.60
Total Sulphur amino acid (TSAA)	2.38	2.47
% TSAA	3.22	3.49
% Cys in TSAA	48.32	51.42
Total Aromatic amino acid (TArAA)	6.79	6.41
% TArAA	9.19	9.07
Predicted Protein Efficiency Ratio (P-PER)	1.69	1.54
Leu/Ile ratio	1.21	1.23
Lys/Arg	0.84	1.09
Essential amino acid index(EAAI)	87.6	84.2

271

272

273

274

275

The amino acids contents of *Nymphaea lotus* seed and *Nymphaea pubescens* seed were presented in Table 2 showing only eighteen amino acids out of the common twenty amino acids found in proteins, including ten essential amino acids and eight non-essential amino acids. This may be due to the conversion of the amide glutamine and asparagine to their corresponding amino acids, glutamate and aspartate respectively [20].

All essential amino acids in both *Nymphaea lotus* and *Nymphaea pubescens* seeds were found to satisfied the FAO/WHO ideal protein value in children except for valine, histidine and methionine which were closer to the value established by FOA/WHO ideal protein in children. Hence, with proper processing, the seeds could meet up with the WHO ideal protein value for both children and adults.

Among the essential amino acids, leucine had the highest concentration in both the samples, followed by isoleucine. The least essential amino acid is methionine with *Nymphaea lotus* seed having the higher content. *Nymphaea lotus* seed had higher concentration of all amino acids compared to *Nymphaea pubescens* seed except for lysine, cysteine, proline and glycine. This could be attributed to genetic difference between the samples.

The arginine content of *Nymphaea lotus* and *Nymphaea pubescens* seeds was higher than the FAO/WHO (1991) recommendations for infants (Arginine 4.0g/100g). Arginine is good for children and it is considerably high in both the *Nymphaea lotus* and *Nymphaea pubescens* seeds with respectively. Isoleucine is an essential amino acid both for young and old. Methionine is needed for the synthesis of choline which in turn forms lecithin and other phospholipids in the body. When the diet is low in protein, for instance in alcoholism and kwashiorkor, insufficient choline may be formed; this may cause accumulation of fat in the liver [22].

The amino acid scores(a rating of the quality of a test protein by comparing its amino acid pattern with that of FAO/WHO reference ideal protein for pre-school children) of *Nymphaea lotus* and *Nymphaea pubescens* seeds are shown on Table 2. The highest scoring essential amino acid was Arginine (133.25%) in *Nymphaea lotus* seed and leucine in *Nymphaea pubescens* seed while methionine was the lowest scoring amino acid (55.91 and 54.55%) in both the samples making it the first limiting amino acid. This is in conformity with the report that the essential amino acids most often acting in a limiting capacity are Methionine and Cysteine, Lysine and Tryptophan [23]. Leucine, Isoleucine and Tryptophan are also present in significant quantities. Methionine is needed for the synthesis of other important substances including choline. The intake of methionine may prevent the fatty liver and causes development and growth [21].

Furthermore, Table 3 shows the nutritional quality determinations of *Nymphaea lotus* and *Nymphaea pubescens* seeds. The nutritive value of a protein depends primarily on the capacity to satisfy the needs for nitrogen and essential amino acids [24]. The total amino acids, TAA (73.82g/100g) of *Nymphaea lotus* seed is higher than *Nymphaea pubescens* seed (70.70g/100g). This difference could be attributed to genetic differences among the *Nymphaea* species and the values obtained in the present study are

comparable with that of some plant foods which range between 39.3 – 76.5g/100g protein as reported by earlier workers [25][26].

The predicted protein efficiency ratio (P-PER) value is one of the quality parameters used in the evaluation of proteins [29]. The P-PER of *Nymphaea lotus* seed and *Nymphaea pubescens* seed were found to be 1.69 and 1.54 respectively, meaning *Nymphaea pubescens* seed may be slightly bioavailable than the *Nymphaea lotus* seed. This is slightly lower than the reported values for true digestible protein of whole dried honey bees (*Apis mellifera* L.), 2.47 and 2.50 for casein [27]. Due to the fact that the experimentally determined PER usually ranged from 0.0 for a very poor protein to a maximum possible value of just over 4 [28], the result of this work shows that both *Nymphaea lotus* and *Nymphaea pubescens* seeds could be efficiently utilized in human body.

Furthermore, the Leu/Ile ratio value of *Nymphaea lotus* and *Nymphaea pubescens* seeds were found to be 1.21 and 1.23 respectively, of which Ile was less than half that of Leu. It has been suggested that an amino acid imbalance from excess leucine might be a factor in the development of pellagra due to sorghum consumption [29]. High Leucine in the diet impairs tryptophan and niacin metabolism and is responsible for niacin deficiency in sorghum eaters [30].

The Lysine/Arginine (Lys/Arg) ratio has been reported to play a role in the artherogenocity of a protein. [31] Reported that a Lys/Arg ratio as high as 2.0, increases the artherogenic potential of a diet. The Lys/Arg ratio of the *Nymphaea lotus* (0.84) and *Nymphaea pubescens* seed (1.09) are much lower than that of casein with Lys/Arg ratio 2.0. A similar observation has been made for *Kerstingiella geocarpa* [32]. Thus, since *Nymphaea lotus* and *Nymphaea pubescens* seeds showed a low Lys/Arg ratio, the samples will possess a low artherogenic potential.

The essential amino acid index (EAAI) and nutritional index of *Nymphaea lotus* seed were higher than the *Nymphaea pubescens* seed. However, the nutritional indexes of the both *Nymphaea* seeds were much higher compared to the *A. bisporus* (13.69%) and *P. florida* (12.59%) as reported by [33]. The essential amino acid index can be useful as a rapid tool to evaluate food formulations for protein quality [34]. However, it does not account for differences in protein quality due to various processing methods or certain chemical reactions [34]. Protein material is said to be of good nutritional quality when its essential amino acid index (EAAI) is above 90% and to be useful as food when the values are around 80% and are inadequate for food material when values are below 70% [35]. From the present study it is observed that the EAAI values were generally appreciable and could be attributed to the fact that *Nymphaea* seeds, a wild haebaceous plant, possessed most essential amino acids in appropriate quantities.

4. CONCLUSION

From the results obtained from this work, it can be concluded that *Nymphaea lotus* and *Nymphaea pubescens* seeds are of high nutritional qualities. Both the seeds are rich in essential amino acids that if consumed in sufficient amount, it could contribute to meeting human nutritional needs and helps to combat diseases associated with malnutrition. The level of arginine and histidine in the seeds warrants them to be recommended for children food since children need arginine and histidine in their foods. This

nutritional information could be of great use to nutritionists, industrialists, researchers, policy makers development agencies and encourage the consumption of *Nymphaea lotus* and *Nymphaea pubescens* seeds so that they become part of normal diet rather than being considered as 'famine' or 'poor peoples' food especially in arid regions.

REFERENCES

- [1]. United Nations Children's Fund, UNICEF. *Facts for life* (4th edition). 2010. New York. Pp 61 and 79
- [2]. Young, E.M. *Food and Development*. Abingdon, oxon. 2012; pp. 36-38.
- [3]. Federal Ministry of Health, Nutrition Department (FMH). Prevalence of Malnutrition in Nigeria. *This Day News Paper* 22 may, 2014;10.
- [4]. Liu, T., Howard, R. M., Mancini, A. J. I., Weston, W. L., Paller, A. S., Drolet, B. A., Esterly, N. B. and Levy, M. L. Kwashiorkor in the United States: fad diets, perceived and true milk allergy and nutritional ignorance. *Archives of Dermatology*. 2012; 1375:630-636
- [5]. Musa, A., Birnin-Yauri, A., Muhammad, C. and A. Umar. Proximate composition and mineral analysis of *Nymphaea lotus* seeds. *African Journal of Food Science and Technology*. 2012; 3(7) PP4
- [6]. Akinjogunla O.J, Yah C.S, Eghafona N.O and Ogbemudia F.O. Antibacterial Activity of Leave Extracts of *Nymphaea lotus* on Methicilin Resistant *Staphylococcus aureus*(MRSA) and Vancomycin-Resistant *Staphylococcus aureus* (VRSA), Isolated from Clinical Samples. *Annals of Biological Research*. 2010; (2):174-184.
- [7]. AOAC. Official methods of analysis, Association of Official Analytical Chemists, Washington, D.C., USA. 17th Edition. 1997; Pp. 807-928.
- [8]. AOAC International. Official method of Analysis of AOAC .17th edition current through 1st revision Gaithersburg, MD, USA, Association of Analytical communities. 2002
- [9] Benitez, L. V. Amino Acid and fatty acid profiles in aquaculture nutrition studies, p. 23- 35. in S.S. De Silva (ed.) *Fish Nutrition Research in Asia*. Proceedings of the Third Asian Fish Nutrition Network Meeting. *Asian fish. Society Special Publication*. 1989; 4:166
- [10]. AOAC, (2010). Official Method of Analysis of the Association of Official Analytical Chemists, Washington, D. C., AOAC. Asian Fisheries Society, Manila Philippines. 18th Edition. 2010; 810-931.
- [11]. Alsmeyer, R. H., Cunningham, A.E. and Happich, M.L. Equations to predict PER for amino acid analysis. *Food Technol*. 1974; 28, 34-38.
- [12]. Wasagu, R. S. U., Lawal, M., Galadima, L. G. and Aliero, A. A. Nutritional composition, Anti-nutritional factors, Elemental analysis of *Nymphaea lotus* (Water lily). *Bayero Journal of Pure and Applied Sciences*, 2015 8(1): 1 – 5.
- [13]. Onimawo, I. A., Oleno, F., Orakpo, G. and Akubur, P. I. Physiochochemical and Nutrient Evaluation of African Bush Mango (*Irvingia gabonensis*) seeds and pulp. *Plant foods for human nutrition*. 2003; 58:1-6

- [14]. Olefe, O., Ogunlade, I. and Jibunoh, N. Proximate, minerals and functional properties of pride of Barbados (*Caesulphina pulcherima*) seeds: proceeding of the international conference of chemical society of Nigeria. Pp 2004;130-134.
- [15]. Mohammed, H. N., Uka, U. N. and Yauri, Y. A. B. Evaluation of Nutritional Composition of Water Lily (*Nymphaea lotus* Linn.) from Tatabu Flood plain North. Central, Nigeria. *Journal of fisheries and Aquatic Science*, 2013;8:261-264.
- [16]. Jenkins, D., Alexandra L., Jenkins, T., Woleve, M. S., Lilian, H. T. and Venkat R. Simple and Complex Carbohydrates. *Nutrition*.1986;3:7:43
- [17]. Genton, L., Melzer, K. and Pichard, C. Energy and Micronutrients Requirements for Physical Fitness in Exercising Subject. *Journal of Agriculture and Food Chemistry*, 2010; 31, 833-836.
- [18]. Sagna, M. B., Aly, D., Papa, S. S., Aliou, G., Ousmane, N. and Rebecca, G. Biochemical composition and nutritional value of *Balanites aegyptiaca* (L.) Del. Fruit pulps from Northern Ferlo in Senegal. *Academic journals*. 2014;13(2) 336-342.
- [19]. Okia, C.A., Agea, J.G., Kimondo, J.M., Abohassan, R. A. A., Okiror, P., Obua, J. and Teklehaimanot, Z. Use and management of *Balanites aegyptiaca* in the drylands of Uganda. *Research Journal of Biological Sciences* 6(1): 15-24, 2011. [20]. Alhassan, A.J. and Sule, M.S. (2011). Characterization and nutritional quality of degutted whitegrubs' protein. *Biological and Environmental Sciences Journal for the Tropics*. 2011;8(3): 26-30.
- [21]. FAO/WHO,. Protein quality evaluation. Report of a Joint FAO/WHO Expert Consultation Food and Agriculture Organization of the United Nations. FAO, Rome, Food and nutrition paper No 51,1991;1-66.
- [22]. Adeyeye, E.I. Amino acid composition of some aquaculture fauna resources in Nigeria. *Chemical and Process Engineering Research* www.iiste.org ISSN.2009;(3)2224-7467
- [23]. FAO/WHO/UNU. Energy and protein requirements. WHO Technical Report Series No 724. WHO Geneva,1985;13-205.
- [24]. Pellet, P.L. and Young, V.R. *Nutritional evaluation of protein foods*, Report of a working group sponsored by the International Union of Nutritional Sciences and the United Nations University World Hunger Programme. The United Nations University,1980;167.
- [25]. Olaofe, O., Adeyemi, F.O. and Adeniran, G.O. Amino acid, mineral composition and functional properties of some oil seeds. *Journal of Agric and Food Chemistry*, 1994;42(4) pp.879 – 881
- [26]. Ogungbenle, H.N. Chemical composition, functional properties and amino acid composition of some edible seeds. *La Rivista Italiana Delle Sostanze Marzo*,2006;83, pp.81 – 86.
- [28]. Muller, H.E. and Tobin, G. *Nutrition and Food Processing*. London: Crown Helm. 1980; 34-36.
- [29]. FAO. Sorghum and millets in human nutrition. Rome, Italy: FAO Food Nutrition Series. 1995
- [30]. Belavady, B. and Udayasekhara-Rao, P. Leucine and Isoleucine content of jower and its pellagragenicity. *Indian Journal of Experimental Biology*. 1996;17: 659-661.
- [31]. Kritchevsky, D. Vegetable protein and atherosclerosis. *Journal of American Oil Chemical Society*, 1999;56;135 – 140

- [32]. Ajayi, O.B. and Oyetayo, F.L. Potentials of Kerstingiella geocarpa as a health food.*Journal of Medicinal Food*, 2009;12(1), pp.184 - 187
- [33]. Oluwole S. I. and Oluremi, O. K. Comparison between the amino acid, fatty acid,mineral and nutritional quality of raw, germinated and fermented African locust bean(*Parkia biglobosa*) flour.*Acta Sci. Pol., Technol. Aliment.* 2012;11(2) 151-165
- [34]. Nielsen S.S.. *Introduction to the chemical analysis of foods*. CBS Publ. New Delhi. 2002; 235-247.
- [35]. Oser, B.L. *An integrated essential amino acid index for predicting the biological value of proteins*. In: Protein and amino acid nutrition. Ed. A.A. Albanese. Academic Press New York, 1998;295-311.