

Proximate Analysis, Hydrogen Cyanide and some Essential Mineral Contents of Sweet  
*Manihot utilisima* Cassava variety (Sweet Cassava variety *Manihot Utilisima*) and Bitter  
Cassava variety *Manihot palmate* (Bitter Cassava variety *Manihot Palmate*) cultivated in  
Kachia Local Government Area of Kaduna State, Nigeria.

Formatted: Font: Italic

ABSTRACT

This study investigated the proximate composition and the hydrogen cyanide content of two different cassava species, bitter cassava and sweet cassava species (*Manihot esculenta* Crantz) variety *utilisima* (sweet cassava) and variety *palmate* (bitter cassava) harvested from Kachia Local Government Area of Kaduna State, Nigeria. The proximate and essential minerals composition analyses were determined using standard methods and titration methods. The mean results of proximate analysis of both species varieties showed that the fresh samples of sweet cassava variety had chloride content value of  $0.0024 \pm 0.0002$  mg/L, ash content:  $0.87 \pm 0.44\%$ , moisture content:  $41.05 \pm 2.26\%$ , dry matter:  $58.9 \pm 0.14\%$  and fat content:  $0.57 \pm 0.42\%$ . The bitter cassava variety had chloride content value of  $0.0028 \pm 0.0000$  mg/L, ash content:  $0.94 \pm 0.06\%$ , moisture content:  $38.2 \pm 2.69\%$ , dry matter:  $61.7 \pm 2.69\%$  and fat content:  $0.49 \pm 0.42\%$ . These values were within the Food and Agricultural Organization and World Health Organization (FAO/WHO) standards for proximate composition in cassava except for the high moisture content of both species which was above the permissible limit. The mean total HCN content in bitter cassava was 89.60 mg/Kg, while the sweet cassava had an HCN content value of 80.60 mg/Kg. The yeast fermentation process was employed and the mean extracted hydrogen cyanide content obtained was as follows:  $69.33 \pm 2.80$  HCN mg/Kg for the sweet cassava and  $85.33 \pm 4.61$  HCN mg/Kg content in the bitter cassava species; the remaining free HCN in both cassava varieties were within recommended consumption limit. The mineral concentration in both cassava varieties were however lower than the recommended mineral values as stated by USDA, 2016.

Formatted: Space Before: 12

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Italic

**Keywords:** Cassava, mineral concentration, hydrogen cyanide, proximate analysis.

1.0. Introduction

Cassava (*Manihot esculenta* Crantz) is cultivated mainly in the tropic and sub-tropic regions of the world, over a wide range of environmental and soil conditions (Richardson, 2011). Latest statistics show that Nigeria is the world largest producer of cassava roots with an annual production of about 45 million metric tonnes (Daily Trust, 2016). This quick spread of cassava development was because of its simplicity of cultivation and minimal effort of production in cost. It develops on poor and ineffectively prepared soils, it can stand extreme drought and insects attack, and the roots can remain underground for a considerable length of time without deterioration, even after its leaves have fallen (Teles, 2002). It is rich in carbohydrates, calcium, vitamin B and C and essential minerals. However, nutrient composition differs according to variety, age of the harvested crop, soil conditions, climate and other environmental conditions during cultivation (IITA, 2009). Besides satisfying the dietary needs of the greater part of the population of Nigeria, there is a huge demand for the roots as raw material for the manufacture of livestock feed, bio fuel, pharmaceutical and textile industries (NRCRI, 2016).

Formatted: Font: Not Italic

Cassava cultivars can be classified into two major types, which are the bitter and sweet variety (Ubwa *et al.*, 2015). Sweet cassava roots contain less than 50 mg per kg HCN on fresh weight basis, whereas that of bitter variety may contain up to 400 mg per kg HCN (CFS, 2008). However, the rate of cyanide in the cassava roots can be reduced with different processing and fermentation methods (Enidioket *et al.*, 2008; Emorotuet *et al.*, 2012). Hydrocyanic or prussic acid (HCN) is produced in cassava, when two major cyanogenic glucosides (linamarin and lotaustralin) are hydrolysed and the acid released when the cassava roots are ruptured (Marcus & Adesina, 2011). Therefore making it very toxic for consumers to eat the cassava root and leaves raw (Emmanuel *et al.*, 2012). Several cases of cassava poisoning have been recorded in Nigeria, all resulting to improper fermentation and processing of cassava (Premium Times, 2017).

57

Chandra *et al.*, (1980) reported that workers exposed to HCN for more than 5 years showed an increase in symptoms such as headache, weakness, changes in taste and smell, irritation of throat, vomiting, lacrimation, abdominal colic, pericardial pain and nervous instability. Acute intoxication, with symptoms of dizziness, nausea, diarrhoea and sometimes death were observed in humans as a result of poor processing cassava products (Mlingiet *et al.*, 1992). Priya *et al.*, (2011) reported that an increase in the triglyceride and Atherogenic index of plasma (AIP) in the blood of cassava workers. While it was reported that tropical ataxic neuropathy (TAN) or similar degenerative neuropathies that causes poor vision occur in Nigeria mainly amongst older people who have consumed cyanide mainly from garri over a long period of years (Oshuntokun, 1994; Nhassico *et al.*, 2008).

Kaduna State in the North West region of Nigeria produces 2 million tonnes of cassava a year, which is comparable in output to many of the states in the southern regions which produce less than 6 million tonnes of cassava (FAO, 2007). With increasing farming activities in the cultivation of cassava in Kaduna State, more interest should be employed on the quality of the cassava products grown in the soils. Kachia Local Government in Kaduna state is home to large cassava, ginger and maize cultivation (Ishaya, 2014). This study was to evaluate the proximate analysis and the use of yeasts fermentation process to reduce the amount of HCN content in the sweet and bitter cassava varieties grown on Kachia soil.

## 2.0. Material and Methods

### 2.1. Sample Collection and Preparation

*Manihot esculenta* Crantz (Cassava tubers (Bitter and Sweet) (*Manihot esculenta* Crantz) were collected from farmlands located in Kachia Local Government Area of Kaduna State Nigeria. The cassava samples were mixed together and representative samples were picked at random. The two species sweet and bitter cassava were cleaned with water and rinsed with distilled water to avoid surface contamination, peeled and stored in a plastic container for fermentation, mineral content analysis and proximate content respectively. Analyses on the cassava samples were prepared in duplicate for proximate content, mineral concentrations and triplicate for fermentation and hydrogen cyanide content determination (Madubuike *et al.*, 2014).




Figure 2.0 Map of Kachia Local government area Showing sampling Location (Source: Nigerian Defence Academy (NDA) Geography Department, Kaduna State).

## 2.2.SAMPLE METHODS OF ANALYSIS

### 2.2.1. PROXIMATE ANALYSIS

**1. Moisture/Dry Matter:** The sliced cassava samples of 2g were accurately weighed into pre-labelled, pre-weighed beaker and transferred to vacuum dry oven to dry at a temperature of 130°C. The samples were heated within a time range of 1hr, 1hr -30 min, 2hrs, 2 hrs -30 min, 3hrs respectively and weighed till a constant Weight was achieved. All sampling and analysis were done in duplicate (Kenneth, 2011).The formula used for the calculation of moisture content and dry matter can be seen below:

$$\text{sampleweight} - \text{moisturecontent} = \text{drymatter}$$

$$\% = \frac{\text{drymatter}}{\text{weightofsample}} \times 100$$

**2. Ash:** Prepared samples were weighed into pre-weighed, porcelain crucibles. The samples were transferred to a muffle furnace and ash at 550°C for 8 hours. The crucibles were allowed to cool in desiccators and then weighed(Kenneth, 2011). The formula used for the calculation of ash content can be seen below:

$$\% = \frac{W\text{tofAsh}}{\text{weightofsample}} \times 100$$

**3. Fat:**The fat content sample was determine by dissolving 8 g of the cassava sample in a 200cm<sup>3</sup> beaker containing 8.4 cm<sup>3</sup> of hydrochloride acid and heated in a water bath for 1h

after heating the sample solution were allowed to cool and extracted with petroleum ether in a separating funnel. After extraction, the sample solution was then heated to dryness and the weight collected after cooling (Kenneth, 2011). The formula used for the calculation of fat content can be seen below:

$$\% \text{ fat} = \frac{\text{weight loss of sample (Extract)}}{\text{weight of sample}} \times 100$$

**4. Chloride:** The ash of the cassava samples weighing 0.064 g was dissolved in a 250 cm<sup>3</sup> beakers with 50 cm<sup>3</sup> of 0.1M silver nitrate (AgNO<sub>3</sub>) and 20 cm<sup>3</sup> HNO<sub>3</sub> was added into the wet ash. The mixture was transferred to conical flask and titrated with 0.1M KSCN, and 0.1M FeNH<sub>4</sub> (SO<sub>4</sub>)<sub>2</sub> were used as indicator. The appearance of brown reddish precipitate marked the end point which was noted. All sampling and analysis were done in duplicate. The chloride concentration (mg/L) was thus calculated (Kenneth, 2011).

#### 2.2.2. DETERMINATION OF MINERAL CONTENT (Ca, Fe, Mg, k, Na.)

Cassava samples of 2 g was washed and transferred into a 250 cm<sup>3</sup> beaker containing 5 cm<sup>3</sup> of aqua and heated for 10 minutes. After digestion, the sample solution was allowed to cool, then filtered and transferred into a 100 cm<sup>3</sup> volumetric flasks. The solution was made up to the mark with deionized water and transferred to a 100 cm<sup>3</sup> sterile plastic container which was then carried to the Atomic Absorption Spectrophotometer and Flame Emission Spectrophotometer for determination of mineral concentration (Kenneth, 2011).

128  
129  
130  
131  
132  
133  
134

Formatted: Indent: First line:

#### 2.2.2. METHOD OF FERMENTATION

##### Procedure: Fermentation of samples using baker's Yeast.

The sample was washed with tap water and after with distilled water respectively. The technique employed involved the addition of 0.5g of hydrated baker's yeast was cultured into a volumetric flask containing 200 cm<sup>3</sup> of distilled water. Thereafter the fermented yeast was transferred to already weighed 20 g of the sample and the retting time was 60 hours. The mixture was then filtered using filter paper and the filtrate was analyzed using UV-Spectroscopy analysis. All sampling and analysis was done in duplicate (Piracicaba, *et al.*, 2009).

#### 2.3.3. PREPARATION OF CALIBRATION STOCK SOLUTION


This was prepared by dissolving 69 g K<sub>2</sub>CO<sub>3</sub> 0.5 M in a 100 cm<sup>3</sup> beaker and transferred to 500 cm<sup>3</sup> volumetric flask and made up to the mark with distilled water. standard solution KCN 0.1M of 0.25 g of potassium cyanide was added to 10 cm<sup>3</sup> of 0.5M K<sub>2</sub>CO<sub>3</sub> solution in a beaker, transferred to a 100 cm<sup>3</sup> volumetric flask and made up to the mark with distilled water. Working standards of concentration 2,4,6,8 and 10 part per million (ppm) were

prepared by serial dilution and taken to the UV-spectrophotometer. The calibration curve for the cyanide determination was obtained by plotting absorbance against concentration of the standard cyanide solutions. Then the graph factor obtained from the plot was then used for the calculation of the cyanide content in the cassava samples (Younoussaet al., 2014).

#### 2.3.4. DETERMINATION OF TOTAL CYANIDE CONTENT IN CASSAVA SAMPLES USING TITRIMETRIC DISTILLATION METHOD

Twenty (20) grams of the cassava sample (bitter and sweet cassava) was weighted and transferred into a 1000cm<sup>3</sup> beaker containing 100 cm<sup>3</sup> of distilled water, 200 cm<sup>3</sup> of phosphate buffer (pH 6.0) and 10 cm<sup>3</sup> of 2 % mercuric chloride solution and left overnight. The sample was then transferred to 500 cm<sup>3</sup> round bottom flask and 5.0 g of hydrated stannous chloride was added to the sample and steam distillation method was used to separate the hydrocyanic acid from the prepared sample. After the mixture was heated by steam distillation, the released hydrogen cyanide was collected in a conical flask containing 50 cm<sup>3</sup> of 1% Alcoholic sodium hydroxyl (NaOH) solution until the volume of the distillate was 200 cm<sup>3</sup>. The distillate was then titrated against 0.02M AgNO<sub>3</sub> using 1cm<sup>3</sup> of freshly prepared 0.5% w/v dithizone in ethanol as indicator and the amount of cyanide in the sample was determined. The end point of the titration was accommodated with a colour change of the indicator from red to purple (FIIRO, 2000-2004).

Chemical equation of the reaction:



1 ml of 0.02m AgNO<sub>3</sub>=0.5mgCN<sup>-</sup>

$$Concentration\ of\ CN^{-} = \frac{Titre\ value \times 0.52 \times 1000}{Weight\ of\ the\ sample\ taken}$$

170

171

### 3.0. Results and Discussion

The following results displayed in the Table 3.1 – 3.4 below were obtained as a result of the analysis done on the determination of the proximate analysis on the cassava samples, their hydrogen cyanide content and mineral concentrations.

**TABLE3.1:** The proximate analysis results of the cassava samples

| Cassava sample | Chloride content (mg/L) | Ash content (%) | Moisture content (%) | Dry matter (%) | Fat content (%) |
|----------------|-------------------------|-----------------|----------------------|----------------|-----------------|
|----------------|-------------------------|-----------------|----------------------|----------------|-----------------|

| varieties                      |               |           |            |           |           |
|--------------------------------|---------------|-----------|------------|-----------|-----------|
| Kachia bitter cassava sample 1 | 0.0028        | 1.02      | 40.10      | 59.80     | 0.78      |
| Kachia bitter cassava sample 2 | 0.0028        | 0.93      | 36.30      | 63.60     | 0.20      |
| Mean Result                    | 0.0028±0.0000 | 0.94±0.06 | 38.20±2.69 | 61.7±2.69 | 0.49±0.42 |
| Kachia sweet cassava sample 1  | 0.0022        | 1.18      | 41.20      | 58.80     | 0.86      |
| Kachia sweet cassava sample 2  | 0.0025        | 0.55      | 42.80      | 59.00     | 0.27      |
| Mean Result                    | 0.0024±0.0002 | 0.87±0.44 | 41.05±2.26 | 58.9±0.14 | 0.57±0.42 |
| Permissible limit<br>FAO/WHO   | 250mg/L       | 3.0%      | 13%        |           |           |

177

178 **TABLE3.2:** Extracted HCN (mg/Kg) content in the yeast fermented cassava samples  
179 (Triplicate analysis).

| TREATMENT                    | Kachia sweet cassava<br>(mg/Kg) | Kachia bitter cassava (mg/Kg) |
|------------------------------|---------------------------------|-------------------------------|
| Cassava sample 1             | 66.00                           | 88.00                         |
| Cassava sample 2             | 71.00                           | 80.00                         |
| Cassava sample 3             | 71.00                           | 88.00                         |
| Mean Results                 | 69.33                           | 85.33                         |
| WHO/FAO<br>Permissible limit | 10 mg/Kg HCN                    |                               |

180

181 Note: KCS: Kachia Sweet Cassava Variety; KCB: Kachia Bitter Cassava Variety.

182

183 **TABLE3.3:** Result of HCN content, free cyanide content, total cyanide content and Permissible limits

| Cassava<br>sample<br>varieties | Total<br>Cyanide<br>content<br>(mg/Kg) | Extracted<br>HCN content<br>(mg/Kg) | Free Cyanide<br>(mg/Kg) | Permissible<br>Limit<br>(mg/Kg) | Above /<br>Below<br>Limits |
|--------------------------------|----------------------------------------|-------------------------------------|-------------------------|---------------------------------|----------------------------|
| Kachia<br>sweet<br>cassava     | 80.60                                  | 66.00                               | 14.60                   | 10.00                           | BPL                        |

|                                |       |       |      |       |     |
|--------------------------------|-------|-------|------|-------|-----|
| sample 1                       |       |       |      |       |     |
| Kachia sweet cassava sample 2  | 80.60 | 71.00 | 9.60 | 10.00 | BPL |
| Kachia sweet cassava sample 3  | 80.60 | 71.00 | 9.60 | 10.00 | BPL |
| Kachia bitter cassava sample 1 | 89.60 | 88.00 | 1.60 | 10.00 | BPL |
| Kachia bitter cassava sample 2 | 89.60 | 80.00 | 9.60 | 10.00 | BPL |
| Kachia bitter cassava sample 3 | 89.60 | 88.00 | 1.60 | 10.00 | BPL |

184 **Note:** APL: Above Permissible Limit, BPL: Below Permissible Limit.

185 **TABLE3.4:** The mineral concentration results of the cassava samples

| Cassava sample varieties         | Calcium(mg/L) | Potassium(mg/L) | Iron(mg/L) | Sodium(mg/L) | Magnesium(mg/L) |
|----------------------------------|---------------|-----------------|------------|--------------|-----------------|
| Kachia bitter cassava sample     | 42.5          | 39.04           | 2.25       | 31.59        | 27.47           |
| Kachia sweet cassava sample      | 32.5          | 34.57           | 4.00       | 32.52        | 34.37           |
| USDA Recommended Value (mg/100g) | 16            | 271             | 0.27       | 14           | 21              |

186

## 187 3.2. Discussion

### 188 3.2.1. Proximate analysis

189 **1.Chloride content:** The bitter cassava variety had higher chloride content than the sweet  
190 cassava variety as harvested in Kachia LGA, Kaduna State. The values as shown in Table 3.1  
191 were within the WHO/FAO permissible limits. Several researchers have reviewed chloride  
192 content to be higher than the reported values. As such as seen in Danso *et al.*, (2001) reported  
193 a chloride content values ranging from  $46.8 \pm 14.8$  -  $48.8 \pm 27.3$  mg/L with some cassava  
194 varieties having no chloride content in them. Edogu, (2007) reported a zero presence of  
195 chloride content in cassava samples analysed in Ebonyi State, Nigeria. Berardi, (2003) stated

196 that chloride is acidic and when taken in high concentration can alter the acid-base load of the  
197 body and obviously manifest as net acid load, if the diet source does not provide enough  
198 basic cations (K, Ca and Mg) to balance it. Soetanet *al.*, (2010) stated that chloride plays a  
199 major role in plants as they are required for water-splitting step of photosynthesis, functions  
200 in water balance, role in root, shoot growth, plays a role in photolysis and deficiency of them  
201 leads to wilting, chlorosis, death of some plants leaves

202 **2. Ash Content:** It was observed that the bitter cassava variety had a higher ash content  
203 percentage value of  $0.94 \pm 0.06\%$  than the sweet cassava variety which had a value of  
204  $0.87 \pm 0.06\%$ . These values were however within the recommended limits as shown in Table  
205 3.1. However, the percentage ash content values were lower than some reviewed literature  
206 studies in Nigeria, as Emmanuelet *al.*, (2012) reported an ash content in six cassava cultivars  
207 in Ghana ranging from  $1.71 \pm 0.32\%$  –  $2.34 \pm 0.15\%$ . Eleazu and Eleazu, (2012) reported an  
208 ash content value ranging from  $1.44 \pm 0.52\%$  –  $2.35 \pm 0.35\%$  from 6 yellow and white cassava  
209 variety cultivated in Umudike, Abia State, Nigeria. Ash content represents the total mineral  
210 content in food after it has been burned at a very high temperature. The variance in the ash  
211 content values of the cassava in study and the reviewed cassava could be as a result of  
212 environmental factors and location of harvesting.

213 **3. Moisture Content:** The sweet cassava variety had a moisture content value of  $41.05$   
214  $\pm 2.26\%$ , which was higher than the bitter cassava variety which had a value of  $38.20 \pm 2.69\%$ .  
215 These values were however higher than the recommended moisture content limits as shown  
216 in Table 3.1. The consequence of having higher moisture content is that the cassava samples  
217 can't be stored for a long period of time. Akpabio *et al.*, 2012 reported a higher moisture  
218 content value of 71.53% and 72.07% for sweet and bitter cassava cultivated in Abia State,  
219 Nigeria. Somendrikaet *al.*, (2017) reported an ash content value of  $52.70 \pm 0.52\%$  for cassava  
220 flesh and  $58.88 \pm 0.25\%$  for cassava peel in a CARI-555 cassava variety in Gannoruwa, Sri-  
221 Lanka. These variations in moisture content values as shown above could be as a result of  
222 cassava variety, locations of planting and time of harvesting.

223 **4. Dry Matter:** The bitter cassava variety had high dry matter content value of  $61.7 \pm 2.69\%$  as  
224 compared to the sweet cassava variety, which had a value of  $58.9 \pm 0.14\%$ . These values are  
225 very good as Nigerian cassava have been reported to have higher dry matter percentage in  
226 cassava varieties. Ngikiet *al.*, (2014) reported a dry matter value of 92.50% for cassava roots  
227 and 90.90% for cassava leaves which is been used for livestock feeds. Dry matter  
228 determination in different genotypes of cassava is important since nutrition and energy  
229 calculation is based on magnitude and nature of dry matter content. Higher dry matter content  
230 naturally would provide greater yield (Fakir *et al.*, 2012).

231 **5. Fat:** The bitter cassava variety had high fat content with a value of  $0.31 \pm 0.02\%$ , while the  
232 sweet cassava variety had a value of  $0.25 \pm 0.05\%$ . These values were within the  
233 recommended limit for fat as shown in Table 3.1. The low fat content however corresponds  
234 with Adegbolaet *al.*, (1992) value of 0.42% for fresh cassava peels. Devendra, (1977)  
235 reported percentage fat content of 1.2% and Ukpabioet *al.*, (2012) also reported a similar low  
236 fat content value of 0.33% for cassava cultivars cultivated from Umudike, Abia State

237 **3.2.2. Total Extracted Cyanide Content:** The total extracted hydrogen cyanide results as  
238 shown-presented in Table 3.2 shows that the bitter cassava variety had the highest extracted  
239 HCN content with a value of  $85.33 \pm 4.61 \text{ mg/Kg}$ , than the sweet cassava variety with a value  
240 of  $69.33 \pm 2.80 \text{ mg/Kg}$ . It can also be seen that the fermentation method using yeast agent was  
241 able to detoxify a large amount of hydrogen cyanide from both cassava variety as shown in

Table 3.3 as reported by (Okaforet *al.*, 1998; Oboh *et al.*, 2003). The fermentation process made the cassava edible for consumption, as it was able to remove the higher HCN level to the recommended permissible level. This is possible because it is one of the natural floras involved in cassava fermentation during processing (Tweyongtete *et al.*, 2002). Babalola, (2014) reported HCN value of  $11.78 \pm 2.02$  mg/Kg and  $15.30 \pm 3.10$  mg/Kg for Garri processed in Ado-Ekiti and Egbira in Ekiti State, Nigeria. Adebayo-Oyetoro *et al.*, (2013) reported a HCN value ranging from  $10.51 \pm 0.13$  mg/Kg –  $16.02 \pm 0.12$  mg/Kg for Garri processed in Ogun and Oyo State, Nigeria and attributed it to the short period of fermentation sometimes employed by the processors. Symptoms of acute cyanide intoxication appear four or more hours after ingesting raw or poorly processed cassava such as vertigo, vomiting, and collapse. In some cases, death may result within one or two hours. It can be treated easily with an injection of thiosulfate, which makes sulphur available for the patient's body to detoxify by converting the poisonous cyanide into thiocyanate (FAO, 1990).

255

**3.2.3. Total Hydrogen Cyanide:** The total hydrogen cyanide present in the cassava samples as shown in Table 3.3 shows that the bitter cassava variety had more hydrogen cyanide with a mean value of 89.60 mg/Kg, while the sweet cassava variety had a value of 80.60 mg/Kg. This supports the fact that the bitter cassava variety has a higher HCN content than the sweet cassava variety. The result also shows why it is harmful for people to eat cassava raw without any fermentation and processing method, in order to detoxify the cassava product (Adindu *et al.*, 2003). Reports have shown that age, variety and environmental conditions influence the occurrence and concentration of hydrogen cyanide in various parts of the cassava plant and at different stages of development respectively (Albert *et al.*, 2005). This means that some varieties generally considered sweet (low cyanide content) can have a high cyanogenic potential under certain conditions (Westby, 2002). Cardoso *et al.*, (2005) reported that in Amazonia (the original source of cassava) and in Africa different varieties have a range of total cyanide contents in the parenchyma from very low to very high (1–1550 ppm). Classical examples of variations in HCN content in cassava varieties can be seen as Seri *et al.*, (2013) reported an HCN value of different varieties with Nigerian origin harvested in Senegal, Soya cassava variety had a value of  $231.2 \pm 10.2$  mg/Kg, Kombo cassava variety had  $104.2 \pm 3.9$  mg/Kg, Gniargui cassava variety had  $270.8 \pm 120$  mg/Kg and TMS 30572 cassava variety had  $171.6 \pm 5.4$  mg/Kg. Agiriga and Iwe (2015) reported an improved cassava variety HCN value with 10, 13 and 16 months of harvesting with TMS 98/0505 having a value of  $43.68 \pm 33$  mg/Kg,  $44.34 \pm 4.92$  mg/Kg and  $47.85 \pm 4.92$  mg/Kg; TMS 30752 had  $42.80 \pm 0.00$  mg/Kg,  $42.68 \pm 45$  mg/Kg and  $46.44 \pm 3.25$  mg/Kg; and TME 419 had value of  $41.75 \pm 28$  mg/Kg,  $40.92 \pm 1.91$  mg/Kg and  $44.85 \pm 6.15$  mg/Kg.

#### 278 **3.2.4. Mineral Concentration**

The mineral (Ca, K, Fe, Na and Mg) were present in the bitter and sweet cassava variety but however lower than the recommended amount of minerals in raw cassava as stated by USDA, (2016) in Table 3.4. Calcium supplies the strength to bones that support locomotion, but it also serves as a reservoir to maintain serum calcium levels (Weaver and Peacock, 2011). Although the requirement for Calcium varies according to age, the range is between 1000 and 1500 mg/day (Bailey *et al.*, 2010). Since Nigeria is the highest producer of cassava and it is one of the main staple foods for the large population of Nigerian, the low calcium content observed shows an urgent input of calcium in the agricultural soils, if deficiency is to be avoided. Calcium insufficiency manifests as decreased bone mass and osteoporotic fracture. In the rapidly growing child, calcium deficiency causes rickets. Low levels of intestinal

calcium resulting from low dietary intakes have also been associated with increased risk of kidney stones and colon cancer (Weaver and Heaney, 2006). The same trend of extremely low amount mineral can be found in iron where the bitter cassava had a value of 2.25 mg/L and 4.00 mg/L for the sweet cassava. Iron exists in the blood mainly as haemoglobin in the erythrocytes and as transferrin in the plasma. It is transported as transferrin; stored as ferritin or hemosiderin and it is lost in sloughed cells and by bleeding (Murray *et al.*, 2000). The average adult stores about 1-3 g of iron in his or her body. A fine balance between dietary uptake and loss maintains this balance (Abbaspouret *al.*, 2014). The most significant and common cause of anaemia is iron deficiency. If iron intake is limited or inadequate due to poor dietary intake, anaemia may occur as a result. This is called iron deficiency anaemia (DeBenoist *et al.*, 2008).

Magnesium is the fourth most abundant mineral and the second most abundant intracellular divalent cation in the body. It is a required mineral that is involved in more than 300 metabolic reactions in the body. Magnesium helps maintain normal nerve and muscle function, heart rhythm (cardiac excitability), vasomotor tone, blood pressure, immune system, bone integrity, and blood glucose levels and promotes calcium absorption (Volpe, 2015). Chen *et al.*, (2014) reported that magnesium enhances exercise performance via increasing glucose availability in the blood, muscle, and brain during exercise. The total body magnesium content of an average adult is 25 g, or 1000 mmol. Approximately 60% of the body's magnesium is present in bone, 20% is in muscle, and another 20% is in soft tissue and the liver. Approximately 99% of total body magnesium is intracellular or bone-deposited, with only 1% present in the extracellular space (Drueke and Lacour, 2007).

Magnesium was however low in both cassava varieties when compared to USDA, (2016) and as such could not reach up the necessary dietary needs for human maximum utilization. Dierck-Hartmut and Dierck-Ekkehard, (2004) reported that patients with diagnoses of depression, epilepsy, diabetes mellitus, tremor, parkinsonism, arrhythmias, circulatory disturbances (stroke, cardiac infarction, arteriosclerosis), hypertension, migraine, cluster head-ache, cramps, neuro-vegetative disorders, abdominal pain, osteoporosis, asthma, stress dependent disorders, tinnitus, ataxia, confusion, preeclampsia, weakness, might also be consequences of the magnesium deficiency syndrome. Potassium and sodium content in the cassava were not enough to meet the necessary recommended value as needed by the human system. Potassium is the main intracellular cation in the body and is principally involved in membrane potential and electrical excitation of both nerve and muscle cells and acid-base regulation (Lanham and Lambert, 2012). Lennon-Edwards (2014) reported that increased potassium intake has been shown to lower blood pressure (BP) even in the presence of high sodium consumption. Low levels of magnesium have been associated with a number of chronic diseases, such as Alzheimer's disease, insulin resistance and type-2 diabetes mellitus, hypertension, cardiovascular disease (e.g., stroke), migraine headaches, and attention deficit hyperactivity disorder (ADHD) (Grober *et al.*, 2015). Sodium is the principal cation in extracellular fluids (Soetan *et al.*, 2010). Sodium is an essential nutrient involved in the maintenance of normal cellular homeostasis and in the regulation of fluid and electrolyte balance and blood pressure (BP). Its role is crucial for maintaining extracellular fluids volume because of its important osmotic action and is equally important for the excitability of muscle and nerve cells and for the transport of nutrients and substrates through plasma membranes (Seldin and Giebisch, 1990). However deficiency of sodium is unlikely despite its low concentration in cassava and other foods. Instead its overconsumption of sodium and its salts that has the major health effects (Strazzullo and Leclerc, 2014).

336

### 3.3, Conclusion and Recommendation

In view on the status of cassava is to the economy and how it is a major tool in alleviating worldwide hunger. Special emphasis must be employed on the varieties of cassava harvested in the country and their HCN and mineral concentration. The cassava variety cultivated in Kachia LGA of Kaduna State show that they can't be stored for a long period of times, as a result of their high moisture content. Also observed in the study, is the low mineral values of Ca, Na, K, Mg and Fe in both cassava varieties, which encourages mineral supplements during diets to meet up with required daily dietary mineral use for humans. Geological locations, environmental factors, soil types, fertilizer applications and mode of cultivation also play a major role in the concentration of minerals and HCN in cassava. The study also shows the high amount of HCN in both cassava variety, this discourage the consumption of raw cassava without processing as toxicity is paramount.

With increasing awareness on the toxicity of HCN in cassava and the emancipation of different techniques in HCN detoxification, the yeast fermentation method could be employed, as it effectively detoxified large HCN content to a permissible acceptable limit. It is recommended that the Government of Kaduna State increase efforts on HCN toxicity in cassava awareness to farmers and consumers and the need for improved processing sites, where farmers can processes their cassava before it is been sold to the general public. A more comprehensive data on HCN content on various cassava varieties should be created for effective monitoring of cassava HCN values, soil location and its nature and mode of plantation and how more genetically improved cassava varieties with low HCN content could be grown in the State.

### References

- Abbaspour, N., Hurrell, R. and Keloshadi, R. (2014). Review on iron and its importance for human health. *Journal of Research on Medical Sciences* 19(2):164 – 174.
- Adebayo-Oyetoro, A. O., Oyewole, O. B., Obadina, A. O and Omemu, M. A. (2013). Cyanide and heavy metal concentration of fermented cassava flour (lafun) available in the markets of Ogun and Oyo States of Nigeria. *International Journal of Biological, Biomolecular, Agricultural, Food and Biotechnological Engineering* 7(7):645 – 648.
- Adegbola, A. A., and Asaolu, O. (1986). Preparation of cassava peels for use in small ruminant production in Western Nigeria. In: *Proceeding of a Workshop Held at the University of Alexandria, Egypt*, Preston T. R. and Nuwanyakpa, M. Y. (eds). pp. 105-115.
- Adegbola, O. B., Smith, J. B., and Okeudo, M. J. (1992). Responses of West African Dwarf Sheep Fed Cassava Peel and Poultry Manure Baked Diets. Department of Animal Science, Obafemi Awolowo.

375 Agriga,A.N and Iwe,M.O. (2015). Optimization of chemical properties of cassava varieties  
 376 harvested at different times using response surface methodology. *American Journal of*  
 377 *Advanced Food Science and Technology* 4(1):10-21.

378 Akpabio,U.D., Akpakpani, A. and Nwo5kocha, G.C. (2012). Comparative Study on the  
 379 physicochemical properties of two varieties of Cassava peels. *International Journal of*  
 380 *Environment and Bioenergy*; 2(1):19-32.

381 Albert, L.C., Klanarong,S&Tzou-chi, H. (2005). Proximate composition, mineral contents,  
 382 hydrogen cyanide and phytic acid of five cassava genotypes.*Food Chemistry* 92(4),  
 383 615-620.

384 Ampe, F., Brauman , A. and Trache, S. Agossou, A. (1994).Cassava retting Optimization  
 385 of a traditional fermentation by an experimental research method. *J. the Sci. of Food*  
 386 *and Agric* 65: 355 -361.

387 Babalola,O.O.(2014). Cyanide content of commercial garri from different areas of Ekiti State,  
 388 Nigeria.*World Journal of Nutrition and Health* 2(4):58-60.

389 Bailey, R.L., Dodd, KW., Goldman, J.A., Gahche, J.J., Dwyer, J.T., Moshfegh, A.J., Sempos,  
 390 C.T and Picciano, M.F. (2010). Estimation of total usual calcium and vitamin D intakes  
 391 in the United States.*JNutr*140(4):817-22.

392 Berardi,J.M. (2003). Covering nutritional bases: The importance of acid-base balance.Culled  
 393 from [www.e-mag.com](http://www.e-mag.com).

394 Cardoso A.P., Mirione, E., Ernesto, M., Massaza, F., Cliff, J andHaque, M.R.  
 395 (2005).Processing of cassava roots to remove cyanogens.*J Food Comp Anal* 18:451–  
 396 460.

397 Chen, H-Y., Cheng, F-C., Pan, H-C., Hsu, J-C., Wang, M-F. (2014). Magnesium Enhances  
 398 Exercise Performance via Increasing Glucose Availability in the Blood, Muscle, and  
 399 Brain during Exercise. *PLoS ONE* 9(1): e85486.

400

401 Committee on World Food Security (2008) 34th Session, Cyanide poisoning and cassava  
 402 food: Safety focus (19<sup>th</sup> issue, February 2008).Incident in focus  
 403 [www.cfs.gov.hk/englisin\\_multimedia/multimedia-pus/multidia-pubfsf-19-](http://www.cfs.gov.hk/englisin_multimedia/multimedia-pus/multidia-pubfsf-19-01.html)  
 404 [01.html](http://www.cfs.gov.hk/englisin_multimedia/multimedia-pus/multidia-pubfsf-19-01.html).Accessed on the 13 April, 2016.

405 Daily Trust,(2014). Nigeria: Country Now World’s largest producer of cassava. Culled from  
 406 [www.allafrica.com/stores/201603290264.html](http://www.allafrica.com/stores/201603290264.html)

407 Danso,K.E., Serfor-Armah, Y., Nyarko, B.J.B., Osae, S and Osae, E.K. (2001).  
 408 Determination of some mineral components of cassava using instrumental neutron  
 409 activation analysis.*Journal of radioanalytical and nuclear Chemistry*250(1) :139-142  
 410 De Benoist B, McLean E, Egli I, Cogswell M, editors. Geneva: WHO Press, World Health  
 411 Organization; 2008. WHO/CDC.Library Cataloguing-in-Publication Data. Worldwide  
 412 prevalence of anaemia 1993-2005: WHO global database on anaemia; p. 40.  
 413 Devendra, C. (1977). Cassava as a feed source for ruminants In Cassava as Animal Feed,  
 414 Mestle, B., and Graham, M. (eds). IDRC, Canada, pp. 107-119.  
 415 Dierck-Hartmut,L and Dierck-Ekkehard,L.(2004). About the misdiagnosis of magnesium  
 416 deficiency.*Journal of the American College of Nutrition* 23(6):730S – 731S.  
 417 Drueke,T.N., Lacour,B.(2007). Magnesium homeostasis and disorder of magnesium  
 418 metabolism.Feebally,J., Floege,J and Johnson,R.J.eds. Comprehensive clinical  
 419 nephrology 3<sup>rd</sup> eds. Philadelphia PA: Mosby. 136-138.  
 420 Eindio, S.E., Attah, .L.E and Otuechere, C.A. (2008). Evaluation of Moisture, Total  
 421 Cyanide and Fiber Contents of Garri produced from cassava (*Manihot utilisima*)  
 422 Varieties Obtained from Awassa in Southern Ethiopia. *Pakistan Journal of Nutrition*; 7  
 423 (5): 625-629.  
 424 Eleazu C.O and Eleazu,K.C.(2012). Determination of the Proximate Composition, Total  
 425 Carotenoid, Reducing Sugars and Residual Cyanide Levels of Flours of 6 New Yellow  
 426 and White Cassava (*Manihot esculenta Crantz*) Varieties. *American Journal of Food*  
 427 *Technology*, 7: 642-649.  
 428 Emmanuel, O.A, Clement. A. Agnes, S.B. Chiwona-Karlton, L and Drinah, B.N. (2012).  
 429 Chemical Composition and Cyanogenic potential of Traditional and High Yielding  
 430 CMD resistant Cassava (*manihotesculantacrantz*) varieties.*International Food Research*  
 431 *Journal* 19(1): 175-181.  
 432 Emurotu., J.E., Balehdeen, U.M. and Ayeni, O.M. (2012). Assessment of heavy metals level  
 433 in cassava flour sold in ayigba market kogi state, Nigeria. *Advances in Applied science*  
 434 *Research*; 3(5): 2544-2548.  
 435 Etonihu, A.C. Olajubu, E.O. Ekanem and S.S. Bako (2011). Titrimetric Evaluation of  
 436 Cyanogens in Paart of Nigerian Cassava Species. *Pakistan Journal of Nutrition* 10(3):  
 437 260-263.

438 Fakir, M.S.A., Jannat, m., Mostafa, M.G. and Seal, H. (2012). Starch and flour Extraction and  
 439 Nutrient Composition of tuber in seven cassava  
 440 accessions. *J. Bangladesh Agril. Univ.* 10(2): 217-222.

441 FAO (Food and Agriculture Organization of the United Nations) Yearbook, (2007). Website:  
 442 [www.fao.org](http://www.fao.org). Accessed

443 FAO/WHO. 1991. Joint FAO/WHO food standards programme. In: Codex Alimentarius  
 444 Commission XII (suppl. 4). Rome, Italy: FAO.

445

446 Food and Agriculture Organization of the United Nations (1990). "Roots, tubers, plantains  
 447 and bananas in human nutrition", Rome, 1990, Ch. 7 "Toxic substances and  
 448 antinutritional factors", under sub-heading "Acute cyanide intoxication." Document  
 449 available online at <http://www.fao.org/docrep/t0207e/T0207E00.htm#Contents>.

450 Food Science. (2012). Ash content in food. Culled from [www.foodscience-avenue.com/2012](http://www.foodscience-avenue.com/2012/11/ash-content-in-food.html?m=1)  
 451 [/11/ash-content-in-food.html?m=1](http://www.foodscience-avenue.com/2012/11/ash-content-in-food.html?m=1).

452 Grober, U., Schmidt, J. and Kister, K. (2015). Magnesium in prevention and therapy. *Nutrient*  
 453 *7*(9):8199-8226.

454 International Institute of Agriculture (IITA). (2009). Culled from  
 455 [www.IITA.org/Cassava.html](http://www.IITA.org/Cassava.html)

456 Ishaya, p. (2014). Analysis of the Ginger, Maize and Soyabeans marketing in kachia local  
 457 Government area, Kaduna state, Nigeria (unpublished master thesis) Ahmadu Bello  
 458 University, Zaria, Nigeria.

459 International Scientific Meeting of Cassava Biotechnology Network; CIAT, Cali Columbia,  
 460 (1987). Culled from [http://onlinelibrary.wiley.com/doi/10.1111/j.1467-](http://onlinelibrary.wiley.com/doi/10.1111/j.1467-7652.2006.00195.x/pdf/379-383)  
 461 [7652.2006.00195.x/pdf/379-383](http://onlinelibrary.wiley.com/doi/10.1111/j.1467-7652.2006.00195.x/pdf/379-383). Accessed on the 6<sup>th</sup> April 2015

462 Jiang, X., Lu, X.W., Zhang, Q.H., Yang, Z.P. (2014). Potential Ecological Risk Assessment  
 463 and prediction of soil Heavy Metal Pollution around coal Gangue dump. *Natural*  
 464 *Hazard and Earth System Science* 14:1559-1610.

465 Kenneth, V.A.R (2011). Evaluation of three cassava varieties for tuber quality and  
 466 yield. gladstone road agricultural centre crop research report no. 4.

467 Knowle and Watkins. (1974). A Practical Agricultural Chemistry Test. Culled  
 468 <http://soeagra.com/iaast/vol1/iaast3.pdf> Accessed 3<sup>rd</sup> may 2016

469 Krisadae, B., Metha, W., Ngarmnit, N. and Sadudee, W. (2009). Enriching Nutritive value of  
 470 Cassava Root by Yeast Fermentation. *Sci. Agric. (Piracicaba, Braz.)* 66(5): 629-

471 633.Culled from [http://www.scielo.br/scielo.php?script=sci\\_arttext&pid=S0103-](http://www.scielo.br/scielo.php?script=sci_arttext&pid=S0103-)  
472 [90162009000500007.html](http://www.scielo.br/scielo.php?script=sci_arttext&pid=S0103-90162009000500007.html)  
473 Lanham,S.A and Lambert,H. (2012). *Potassium.Adv. Nutr* 3:820 – 821.  
474 Lennon-Edwards,S., Allman,S.R., Schellhardt,T.A., Ferreira,C.R., Farquhar,W.B and  
475 Edwards,D.G. (2014). Lower potassium intake is associated with wave reflection in  
476 young healthy adults. *Nutrition Journal* 13:39.  
477 Marcus, A.A. and Adesina,B.S. (2001). The Effect of Cooking Time on Root Mealiness and  
478 Taste of Cassava Tuber.*Nigeria Journal of Food Science and Technology* 113.  
479 Murray, R.K., Granner, D.K., Mayes, P.A and Rodwell, V.W. (2000). Harper's Biochemistry,  
480 25th Edition, McGraw-Hill, Health Profession Division, USA.  
481 Madubunike, P.C., Onyema. C.T. Odinma., S.C and Sokwaibe, C.E .(2014). Evaluation of  
482 Elements,Cyanide and proximate composition of cassava (*Manihot Escuenta Crantz*)  
483 from Ebonyi state, Nigeria. *OSR Journal of Environmental Science, Toxicology and*  
484 *food technology*.8 (8):41-43  
485 Mingi, N., Paulter, N.H. and Rosling, H. (1992) an outbreak of acute intoxications from  
486 consumption of insufficiently processed cassava in Tanzania, *Nutr. Res.* 12, 677-687  
487 National Root Crop Research Institute, Umudike (NRCRI). 2006, Cassava  
488 programme.From [www.ncrci.gov.ng/pages/cassava.html](http://www.ncrci.gov.ng/pages/cassava.html).  
489 Ngiki, Y.K Igwebuike,J.U and Moruppa, S.M. (2014). Effect of replacing maize with cassava  
490 root- leaf mixture on the performance of broiler chickens. *International Journal for*  
491 *science and Technology* 3 (6):352-362  
492 Nhassico, D., Muquingue, H., Cliff, J., Cumbana, A and Bradbury, H.J. (2008). Rising  
493 African Cassava Production, Diseases Due To High Cyanide Intake and Control  
494 Measures.*J. Sci. Food Agric* 88: 2043-2049.  
495  
496 NRCRI, National Root Crop Research Institute. (2005). Proximate Analysis of fresh Cassava  
497 Root of Six Cultivars.  
498 Oboh, G and Akindahunsi, A.A. (2003) Biochemical changes in cassava products (flour  
499 & garri) subjected to *Saccharomyces cerevisiae* solid media fermentation. *Food*  
500 *Chemistry*, 82, 599-602.

501 Okafor, N, Umeh C, Ibenegbu, C, Obizoba, I. Nnam, N. (1998). Improvement of Garri  
 502 quality by the inoculation of microorganisms into cassava mash. *International Journal*  
 503 *Food Microbiology* 40: 43-49

504 Oluwole, O.B, Olatunji O.O., &Odunfa, S. A. (2004).A Process Technology for Conversion  
 505 of Dried Cassava Chips into Garri. *Nigeria Food Journal* 22, 65-73

506 Orjiekwe, C.L., Bolola, A., Iyen, E and Imade, S. (2013). Determination of  
 507 Cyanogenic Glucosides in Cassava product sold in Okada, Edo State, Nigeria. *African*  
 508 *Journal of Food Science* 7(12): 468-472.

509 Oshuntokun, B.O. (1994). Chronic cyanide intoxication of dietary origin and a degenerative  
 510 neuropathy in Nigerians. *Acta Horti* 375:311- 321.

511 Premium Times (2017). Cassava flour meal kills 9 in Kogi State. Culled from  
 512 [www.premiumtimesng.com/news/headlines/214443-cassava-flour-meal-kills-9-](http://www.premiumtimesng.com/news/headlines/214443-cassava-flour-meal-kills-9-kogi.html)  
 513 [kogi.html](http://www.premiumtimesng.com/news/headlines/214443-cassava-flour-meal-kills-9-kogi.html).

514 Priya, K.D., Pachiappan, C., Sylvia, J and Aruna, R.M. (2011). Study on the effects of hydrogen  
 515 cyanide exposure in cassava workers. *Indian J Occup Environ Med* 15(3):133-136.

516 Piracicaba (2009) Enriching nutritive value of cassava root by yeast fermentation. *Journal*  
 517 *of Sci. Agric.* 66(2)242-249 culled from  
 518 [http://www.scielo.br/scielo.php?script=sci\\_arttext&pid=S0103-90162009000500007](http://www.scielo.br/scielo.php?script=sci_arttext&pid=S0103-90162009000500007)

519 Rafael Marques Pereira Leal<sup>1,2</sup>; Lilian Pittol Firme<sup>1,2</sup>; Céla Regina Montes<sup>3,2</sup>;  
 520 Adolpho José Melfi<sup>2,4\*</sup>; Sonia Maria De Stefano Piedade  
 521

522 Rechardson, V.A.K. (2011). Evaluation of three cassava varieties for Tuber Quality and  
 523 Yield. Gladstone Road Agricultural Centre. Crop Research Report No. 4.

524 Seldin, D.W and Giebisch, G. (1990). The regulation of sodium and chloride balance. New  
 525 York: Raven Press.

526 Seri, S.G., Souleymane, T and Kouakou, B. (2013). Assessment of cyanide content in cassava  
 527 varieties and derived products from Senegal. *International Journal of Nutrition and*  
 528 *Food Sciences* 2(5):225 – 231.

529 Soetan, K.O., Olaiya, C.O and Oyewole, O.E. (2010). The importance of mineral elements for  
 530 humans, domestic animals and plants: A review. *African Journal of Food Science*  
 531 4(5):200-202.

532 Somendrika, M.A.D., Wickramasinghe, I., Wansapala, M.A.J and Peiris, S. (2017). Nutritional  
533 Composition of Cassava Cultivar "CARI-555". *Pakistan Journal of Nutrition*, 16: 216-  
534 220.

535 Strazzullo, D and Leclercq, C. (2014). Sodium. *Adv. Nutr* 5:188 – 190.

536 Twayongyere, R., Katongole, I. (2002.) Cyanogenic potential of cassava peels and their  
537 detoxification for utilization as livestock feed. *Veterinary and Human Toxicology*. 366-  
538 369,

539 Teles (2002) Processing Techniques to Reduce Toxicity and Antinutrients of Cassava for Use  
540 as a Staple Food. *Comprehensive Reviews in Food Science and Food Safety*. Culled  
541 from [www.10.1111/j.1541-4337.2008.00064.x](http://www.10.1111/j.1541-4337.2008.00064.x)

542 Ubwa, S.T., Otache, M.A., Igbum, G.O and Shambe, T. (2015). Determination of Cyanide  
543 Content in Three Sweet Cassava Cultivar in Their Local Government Area of Benue  
544 state, Nigeria. *Food and Nutrition Science* 6:1078-1085.

545 United State Department of Agriculture (USDA) (2016). Nutritional Nutrient Database for  
546 Standard Reference Release  
547 28. Culled from [www.ndb.nal.usda.gov/ndb/foods/show/2907?manu=&fgcd=&ds=html](http://www.ndb.nal.usda.gov/ndb/foods/show/2907?manu=&fgcd=&ds=html).

548 Volpe, S.L. (2015). Magnesium and Athlete. *Current Sports Medicine Reports* 14(4):279 – 283.

549 Weaver, C.M and Heaney, R.P. (2006). Calcium. In: Shils ME, Shike M, Ross AC, Caballero  
550 B, Cousins RJ, editors. *Modern nutrition in health disease*. 10th ed Baltimore  
551 (MD): Lippincott Williams & Wilkins;. p 194–210.

552 Weaver, C.M and peacock, M. (2011). Calcium. *Adv Nutr* 2:290 – 292.

553 Westby, A. (2002). Cassava utilization, storage and small scale processing. In: Hillocks RJ,  
554 Thresh JM, Belloti, A.C. (Eds.). *Cassava: biology, production and utilization*, p 281-  
555 300. CAB International Publishing, Wallingford.

556 Younoussa, D., Momar, G. Cheikh, N. Mama, S. Amadou, K., Jean, P. B. and Georges L.  
557 (2014). A New Method for the Determination of Cyanide Ions and Their Quantification  
558 in Some Senegalese Cassava Varieties. *American Journal of Analytical Chemistry*,  
559 5:181-187

560