

Comparative Study of the Effects of the Commonly-consumed Aqueous Extracts of *Hibiscus sabdariffa* (Zobo Drinks) on Hepatic and Renal Indices of Normal Wistar Albino Rats.

ABSTRACT

Aim: Investigating and comparing the effect of administration of different preparations of the commonly-consumed *Hibiscus sabdariffa* (Zobo) drinks on hepatic and renal indices.

Study design: Animal models (Wistar Albino Rats) with daily administration of the same concentration of different zobo drink samples.

Place and Duration of Study: University of Port Harcourt, Choba, Rivers State, Nigeria and its environs between November 2014 and February 2015.

Methodology: Thirty (30) Wistar albino rats were grouped into six (6) groups of five rats each. Group A served as the control and group B was administered an unblended zobo drink. Groups C – E were administered locally-produced zobo samples and group F was a National Agency for Food and Drug Administration and Control (NAFDAC)-branded zobo drink. A concentration of 200 mg/kg body weight of the samples was administered orally to the groups B – F for 21 days. Hepatic marker enzymes (ALT, AST and ALP), bilirubin (total and conjugated), albumin, total protein and renal indices were analyzed and compared

Results: The results indicate that the activities of ALT, ALP, Albumin and Unconjugated bilirubin were significantly lower ($P < 0.05$), while Total protein and Total bilirubin levels were significantly higher ($P < 0.05$) than the control (Group A). AST and Conjugated bilirubin showed no significant difference ($P > 0.05$) in all the groups. The levels of Urea, Creatinine, Sodium (in group B), Potassium (K^+), and Chloride (Cl^-) were significantly higher ($P < 0.05$) than the control. The Sodium (Na^+) level was significantly lower ($P < 0.05$) than the control in groups C, D and E.

Conclusion: The results of this study suggest that all the samples were not hepatotoxic. However, the aqueous extract of Zobo may have nephrotoxic and hepatotoxic effects if the concentration is increased over time.

Keywords: [*Hibiscus sabdariffa*; Zobo; Liver enzymes; Renal Indices; Hepatotoxic; Hepatoprotective; Nephrotoxic]

1. INTRODUCTION

Hibiscus sabdariffa belongs to the order of Malvales and the family of Malvaceae. It is believed to originate from East Africa [1]. *H. sabdariffa* plants are cultivated and consumed as vegetable and tea, whereas other *Hibiscus* varieties are planted for the fibres they produce. It is called different names like Roselle and Sorrel in English and it is locally called zobo and Isapa in Nigeria [2]. Various types of highly valued food and medicinal products are produced from parts of the *Hibiscus sabdariffa* including the seeds, leaves, fruits and roots. Among them, the fleshy red calyces are the most popular [3] as they are commonly used to prepare non-alcoholic drinks in form of beverage or juice.

The non-anthocyanin portions of the *Hibiscus sabdariffa* calyces also have hepatoprotective effects, as evidenced by a pigment free extract that is able to protect from acetaminophen induced hepatotoxicity [4]. A study in persons with metabolic syndrome noted that 31 days of 100 mg Roselle extract (19 mg Anthocyanin Sambubiosides) was able to reduce serum ALT and AST, suggesting Roselle attenuates hepatic damage [5].

In Nigeria, the dried calyces of this plant are processed into a refreshing non-alcoholic local beverage commonly called zobo, zaborodo or Isapa (pronounced Isakpa) [6].

Zobo beverage is made from different varieties of calyces of the flower *Hibiscus sabdariffa* by boiling and filtration [7, 8]. Zobo drink, which is usually prepared with the calyces of *Hibiscus sabdariffa* is

believed to be highly nutritive and has many medical potentials including reduction in blood pressure, anti-diabetic, reduction in weight, antihyperlipidemic, hepatoprotective, anti-cancer, as well as an antioxidant e.t.c. [9, 10, 11, 12, 13].

Following increased religious and health awareness against consumption of alcoholic beverages in Nigeria and the consequent decrease in the consumption of alcoholic beverages in certain areas, Zobo drink has great potential as a local alternative to imported red wines in particular and alcoholic beverages in general [14]. Moreover, production of this and similar local beverages has become the main source of income in many homes in the rural communities and more recently in the urban areas where these have grown to cottage business proportions following governmental interventions through the poverty alleviation schemes, thereby alleviating poverty among the people [15]. Zobo is mostly consumed by low and middle class people due to its relative low cost, because the *Hibiscus sabdariffa* calyces and the ingredients are cheap and easy to get. It has recently gained wider acceptance, being consumed by several millions of people from different socio-economic classes and background in the West Africa sub-region and in Nigeria particularly.

Within the University of Port Harcourt and Choba community, the rate of consumption of Zobo drink by students and staff alike is enormous. This is reflective of the perception by many, that, zobo drink is highly nutritious, medicinal and of course, cheaper (following the prevailing economic downturn in Nigeria) than other non-alcoholic beverages sold around and within the environment. However, many of the local producers of the drinks prepare the drinks under poor hygienic conditions, using different preparation methods and variants of synthetic flavours. Some of these synthetic flavours have been suspected of being toxic or carcinogenic and many have been banned whenever possible [16]. Some also believe that the consumption of zobo drink supplemented with flavour increased its antioxidant potentials and has no severe effect on the liver and kidney and thus, supplementation of zobo drinks with flavour additive maybe nutritionally beneficial [17].

Therefore, this study will be of importance in evaluating the effect of zobo drinks on hepatic and renal using normal Wistar albino rats. It will seek to find out and/or add to existing literature, whether or not, consumption of zobo drinks supplemented with synthetic flavours is beneficial to the body.

2. MATERIALS AND METHODS

2.1 Chemicals and Reagents

All chemicals and reagents used were of analytical grade. Alanine Transaminase (ALT) and Aspartate Transaminase (AST), Bilirubin, Total Protein and Albumin Test Kits by Randox Laboratories Ltd, Crumlin, England, UK were used.

2.2 Plant Material

Dried calyces of *Hibiscus sabdariffa* were bought from Choba market, Port Harcourt on 11th November, 2014. They were identified and authenticated in the Herbarium of the Department of Plant Science and Biotechnology (PSB), University of Port Harcourt where voucher specimens with code number 2435 were deposited.

2.3 Samples and Preparations

The unblended zobo drink sample was prepared using the method described by [17]. The dry calyces of *Hibiscus sabdariffa* (HS) were carefully sorted to remove dirt and other unwanted materials. Sixty grams (60 g) of the dry HS calyces were washed with cold water and added to two litres of boiling distilled water. It was allowed to boil for 15 minutes and then cooled. After cooling, the mixture was sieved with muslin cloth and filtered with Whatman No. 1 filter paper. The clear filtrate was covered with aluminium foil and stored in the refrigerator at 4 °C until use.

The ZOBO COLA was also used for comparison. This is a commercially sold branded 400 mL Zobo drink produced by Zobo Cola Company Ltd, Nigeria, certified by National Agency for Food and Drug Administration and Control (NAFDAC) and sold in shops within the community. The ingredients used to prepare the Zobo Cola include: Purified water, HS extracts, Aspartame, Sugar, Cola flavour, Ginger and Citric acid. Other drink samples were obtained from shops around the community.

2.4 Experimental Animals

Thirty (30) Wistar Albino rats weighing between 110 – 195 g were used for the study. The animals were obtained from the Animal House of the Department of Biochemistry, University of Port Harcourt, Choba, Rivers state. All the animals were housed in the animal house, University of Port Harcourt, Choba Campus – using plastic cages covered with wire gauze and given standard food pellets (Top Feeds' grower's mash) and water *ad libitum*. They were acclimatized for 2 weeks and marked for easy identification and monitoring, after their baseline weights were taken. All procedures and techniques in handling the animals were according to standard methods and complied with the guidelines of the National Institutes of Health of the United States [18].

2.5 Experimental Design and Administration of Samples

The acclimatized albino rats were sorted according to their weights into five groups of six rats each. Group A was fed the normal rat feed with water and served as the control. Groups B – F served as experimental groups and were administered 200 mg/kg body weight of the respective samples via oral intubation for a period of 21 days. Group B was administered the unblended zobo drink. Group F was administered the Zobo Cola, while groups C, D and E were administered with other drink samples obtained from shops around the community. All animals were allowed access to water and food for the 21 days.

2.6 Sacrificing of Animals and Collection of Blood Samples

All the animals from the groups were sacrificed at the end of the administration period. The animals were incapacitated with chloroform in a desiccator. Under this condition, the rats were dissected using dissecting tools and the blood was collected and put into lithium heparin and EDTA anticoagulant tubes (to prevent blood clotting) for biochemical analyses.

2.7 Determination of Biochemical Parameters

The plasma ALT and AST activities were determined by monitoring the concentration of pyruvate and oxaloacetate hydrazine respectively, formed with 2,4-dinitrophenylhydrazine according to [19] using Randox test kits. The plasma ALP activity was determined using the colorimetric end-point method according to Rec [20]. The plasma total and conjugated bilirubin concentration were determined using Randox Test Kits according to the Jendrassik and Grof [21] method. Total protein concentration was determined by Biuret Method using Randox Test Kits according to Tietz [22]. Albumin concentration was determined by the Bromocresol Green (BCG) Dye Binding Method by [23], using Randox Test Kits.

2.8 Determination of Renal Indices

The Urease-Berthelot method according to Fawcett and Scott [24] as provided by Randox diagnostics Ltd. UK was used. Creatinine was determined by the direct end method of [25]. Sodium concentration was determined using the method of Maruna [26] and Trinder [27]. Potassium and Chloride ions were determined using the Colorimetric method.

2.9 Statistical Analysis

All data obtained in this study were subjected to statistical analyses using One-way Analysis of Variance (ANOVA). Tukey's Multiple Range Test was used to test for differences between the administration groups. All analyses were done using Statistical Package for Social Sciences (SPSS) version 20.0 (IBM Statistics, UK). All the values were reported as means $\pm$ standard error of mean (SEM) and the results were considered significant at P-values of less than 0.05 ($P < 0.05$) i.e. at 95 % confidence level.

3. RESULTS AND DISCUSSION

Results of the comparative effect of different aqueous extracts of *Hibiscus sabdariffa* (Zobo drinks) on activities of liver marker enzymes (ALT, AST and ALP), Total protein (TP), Albumin (ALB), Total bilirubin, conjugated bilirubin and unconjugated bilirubin (TB, CB, UC respectively) are summarized in "Table 1". From the results, the plasma ALT level in groups B and C were lower than the control, though only group B was significantly lower ($P < 0.05$) when compared with the control (Group A). Meanwhile, the ALT level in groups D, E and F was slightly higher but not significantly ($P > 0.05$)

when compared with the control. Although the AST level was lower in groups B, C, E and F, they were not significantly different when compared with the control. The ALP level of groups B, C, D and E was significantly lower ($P < 0.05$) than the control, while there was no significant difference ($P > 0.05$) between group F and the control (Group A). The Total protein level of groups B, E and F was significantly higher ($P < 0.05$) than the control, while there was no significant difference ($P > 0.05$) in groups C and D when compared to the control (Group A). The Albumin level in groups C and D was significantly lower ($P < 0.05$) than the control and there was no significant difference ($P > 0.05$) in Groups B, E and F when compared with the control (Group A). The Total bilirubin level in groups C, D, E and F was significantly higher ($P < 0.05$) than the control, while it was almost significantly higher ($P > 0.05$) in group B when compared with the control (Group A). The level of Unconjugated bilirubin in groups D and F was significantly higher ($P < 0.05$) than the control, while it was also higher in groups B, C and E but not significantly ($P > 0.05$). The levels of AST and conjugated bilirubin showed no significant difference ($P > 0.05$) when compared with the control (Group A).

Table 1. Effect of 21 days administration of different zobo drinks on biochemical parameters of wistar albino rats

GROUP	ALT (U/L)	AST (U/L)	ALP (U/L)	TP (g/L)	ALB (g/L)	TB (μ mol/L)	CB (μ mol/L)	UB*
A	14.6 $\pm$	23.0 $\pm$	67.6 $\pm$	69.6 $\pm$	36.0 $\pm$	5.66 $\pm$	2.40 $\pm$	3.26 $\pm$
(CNTRL)	0.87 ^a	0.32 ^a	0.68 ^a	1.50 ^a	0.63 ^a	0.36 ^a	0.63 ^a	0.32 ^a
B (ZSTD)	8.0 $\pm$	17.8 $\pm$	56.0 $\pm$	81.0 $\pm$	37.2 $\pm$	6.74 $\pm$	2.24 $\pm$	4.50 $\pm$
	0.63 ^b	0.58 ^b	1.48 ^a	1.00 ^b	0.37 ^a	0.05 ^a	0.07 ^a	0.10 ^a
C (ZAP1)	10.0 $\pm$	19.6 $\pm$	43.0 $\pm$	65.6 $\pm$	31.8 $\pm$	7.40 $\pm$	3.80 $\pm$	3.60 $\pm$
	0.89 ^{ab}	1.57 ^a	2.55 ^a	1.47 ^a	1.02 ^b	0.07 ^b	0.47 ^a	0.43 ^a
D (ZAP2)	18.6 $\pm$	25.2 $\pm$	58.6 $\pm$	73.6 $\pm$	32.0 $\pm$	9.00 $\pm$	2.40 $\pm$	5.60 $\pm$
	2.27 ^b	0.66 ^b	2.16 ^a	2.06 ^a	1.41 ^b	0.40 ^b	0.03 ^a	0.14 ^b
E	14.8 $\pm$	19.6 $\pm$	30.6 $\pm$	79.0 $\pm$	36.8 $\pm$	6.76 $\pm$	4.86 $\pm$	5.10 $\pm$
(ZCHO)	1.16 ^a	1.21 ^b	0.40 ^a	1.00 ^b	0.73 ^a	1.02 ^b	0.85 ^a	0.70 ^a
F	15.4 $\pm$	22.4 $\pm$	61.0 $\pm$	77.0 $\pm$	34.2 $\pm$	9.46 $\pm$	2.36 $\pm$	7.10 $\pm$
(ZCOLA)	0.93 ^a	2.79 ^a	0.71 ^b	0.37 ^b	0.37 ^a	0.57 ^b	0.05 ^a	0.54 ^b

Values in the table are means $\pm$ Standard error of mean (SEM) and n = 5. At ($P < 0.05$), means with different superscripts in a column are significantly different. ALT = Alanine transaminase; AST = Aspartate aminotransferase; ALP = Alkaline phosphatase; TP = Total protein; ALB = Albumin; TC = Total bilirubin; CB = Conjugated Bilirubin; UC = Unconjugated bilirubin. * = UB values were calculated. **ZSTD** = Unblended Zobo drink; **ZAP1** = Zobo drink sold within Abuja Campus Park

(UNIPORT); **ZAP2** = Zobo drink sold outside Abuja Campus Park; **ZCHO** = Zobo drink sold in Choba; **ZCOLA** = Branded Zobo drink (Zobo Cola®)

To ascertain the state of living cells, AST, ALT and ALP are basic biochemical markers that are usually measured. These cytosolic enzymes are usually more concentrated in some organs like the liver, heart and in other parts of the body. These enzymes usually leak out of damaged tissues into the plasma, leading to their high level in the plasma. Such elevation of the enzymes in the blood is therefore an indication of tissue damage and altered membrane permeability [28]. Therefore, the results suggest that **all the samples were not hepatotoxic**. This is in line with Tseng *et al* [29], which reported a hepatoprotective effect of extracts of *Hibiscus sabdariffa*. It is worthy of note that group D showed higher levels of ALT and AST, though not significant. This is suggestive that if the concentration of the sample is increased over time, it may cause a physiological dysfunction or damage to the hepatocytes. The non-significant changes in the levels of ALT, AST and ALP suggest that the liver functions were not impaired in those groups. **Total protein is a rough measure of all the proteins found in the fluid portion of the blood, specifically looking at the total albumin and globulin [30]. Total protein may be required to get an overview information about nutritional status like when someone has undergone a recent weight loss and also to provide information (alongside other tests) if there is any symptom that suggests a liver disorder. An increase in total protein is usually the result of tissue damage [31]. Therefore, the significant higher levels of total protein in groups B, E and F suggests that those drink samples may be hepatotoxic and also, the excretion of protein by the kidney may be impaired. Meanwhile, the albumin (a major component of total protein) level in groups C and D were significantly lower than – suggesting liver function was not impaired in those groups. The elevated levels of total bilirubin (in groups C, D, E and F) and unconjugated bilirubin (in groups D and F) is an indication that there may hemolytic anemia, internal hemorrhage, reduced hepatocyte uptake, impaired conjugation of bilirubin, reduced hepatocyte secretion of bilirubin or obstruction of the bile ducts in those groups.**

Results of the Renal Function Test (Urea, Creatinine, Potassium, Sodium, Bicarbonate and Chloride) of groups A – F are summarized in “Table 2”. From the results, the level of Urea in groups B, D and E was significantly higher ($P < 0.05$) than the control. The level of Urea in group F was significantly lower ($P < 0.05$) than the control, while there was no significant difference ($P > 0.05$) in group C when compared with the control (Group A). The Creatinine level in groups B, D, E and F was significantly higher ($P < 0.05$) than the control, while the increase in group C was not significant ($P > 0.05$) when compared with the control. The Potassium level in group F was significantly higher ($P < 0.05$), while there was no significant difference ($P > 0.05$) in the groups B, C, D and E when compared with the control. The Sodium level in group B was significantly higher ($P < 0.05$), while the level in groups C, D and F was significantly lower than the control. There was no significant difference ($P > 0.05$) in group E when compared with the control. The chloride level was higher in all the groups (B – F), but this was only significant ($P < 0.05$) in Groups E and F when compared with the control. There was no significant difference ($P > 0.05$) in bicarbonate (HCO_3^-) level of all the groups.

Table 2: Effect of 21 days administration of different zobo drinks on renal indices of wistar albino rats

GROUP	Urea	Creatinine	K^+	Na^+	HCO_3^-	Cl^-
	(mmol/L)					
A (CNTRL)	6.04 ± 0.70 ^a	330.00 ± 0.71 ^a	4.22 ± 0.23 ^a	60.0 ± 0.71 ^a	25.00 ± 1.48 ^a	73.60 ± 7.58 ^a
B (ZSTD)	6.70 ± 0.08 ^b	397.00 ± 5.64 ^b	3.95 ± 0.14 ^a	75.60 ± 3.37 ^b	26.00 ± 0.32 ^a	87.00 ± 1.10 ^a
C (ZAP1)	6.30 ± 0.17 ^a	336.60 ± 11.42 ^a	4.45 ± 0.09 ^a	10.00 ± 0.32 ^c	28.00 ± 0.71 ^a	87.20 ± 1.20 ^a

D (ZAP2)	6.75 ± 0.11 ^b	418.60 ± 11.03 ^b	4.15 ± 0.06 ^a	10.00 ± 0.32 ^c	27.00 ± 1.00 ^a	87.80 ± 1.39 ^a
E (ZCHO)	7.65 ± 0.13 ^b	434.60 ± 5.45 ^b	4.10 ± 0.11 ^a	56.00 ± 10.07 ^a	27.80 ± 0.20 ^a	92.60 ± 2.73 ^b
F (ZCOLA)	4.80 ± 0.04 ^c	382.60 ± 2.60 ^b	4.80 ± 0.03 ^b	10.20 ± 0.20 ^c	24.40 ± 0.24 ^a	91.20 ± 0.58 ^b

206 Values in the table are means ± Standard error of mean (SEM) and n = 5. At (P < 0.05), means with
207 different superscripts in a column are significantly different.

208 K⁺ = Potassium ion; Na⁺ = Sodium ion; HCO₃⁻ = Bicarbonate ion

209 The higher levels of renal indices in the groups (B, D, E and F) suggest that the zobo drinks may have
210 nephrotoxic effects (kidney impairment) on the experimental animals. The higher levels of sodium,
211 potassium and creatinine, and no significant difference in bicarbonate, are in line with [32], which
212 obtained the same trend in those renal indices. The likely kidney impairment is in line with Ogundapo
213 [17], which implied that nephrotoxic effects of zobo drinks supplemented with commercial flavour
214 additives may have been due to the presence of sodium cyclamate present in the flavour, which
215 causes kidney damage and other effects [33]. The high levels of sodium in group B, chloride in groups
216 A – F and potassium in group F is in line the study carried out by [34] on the diuretic property of
217 *Hibiscus sabdariffa*. The study observed that rats that consumed aqueous *Hibiscus sabdariffa* extract
218 had elevated plasma sodium, chloride and potassium. Aqueous extract of *Hibiscus sabdariffa* had
219 also been shown in another study to be rich in Na⁺ [35]. By implication, the consumption of *Hibiscus*
220 *sabdariffa* by the rats may have caused an increase in the oral Na⁺ load which consequently
221 increased plasma Na⁺. A toxic insult to the kidney therefore, could have profound effect on total body
222 metabolism [36]. This is because; the proper functioning of the kidney ensures proper excretion of
223 metabolic wastes, maintenance of acid-base balance, electrolyte composition and regulation of
224 intracellular fluid volume [37].

225

226 4. CONCLUSION

227 This study aimed at investigating and comparing the effects of 21 days administration of different
228 preparations of *Hibiscus Sabdariffa* (Zobo) drink consumed around the University of Port Harcourt
229 community **hepatic and renal indices** using normal albino rats. All the samples of zobo drink **were not**
230 **hepatotoxic**. The non-significant changes in the levels of ALT, AST and ALP suggest that the liver
231 functions were not impaired in those groups.

232 The higher levels of renal suggest that the zobo drinks had nephrotoxic effects (kidney impairment) on
233 the experimental animals.

234 Despite the enormous nutritive, health and medical potentials of the aqueous extract of *Hibiscus*
235 *sabdariffa* (Zobo drink), this study showed that the drink may have has nephrotoxic properties. Dose
236 and time-dependent studies of the effects of different samples of zobo drink on biochemical and renal
237 indices is strongly recommended. Also, histological analysis should be done on the vital organs of
238 animals administered with Zobo drinks.

239

240

241 ETHICAL APPROVAL

242 All authors hereby declare that "Principles of laboratory animal care" (NIH publication No. 85-23,
243 revised 1985) were followed, as well as specific national laws where applicable. All experiments have
244 been examined and approved by the appropriate ethics committee.

245

246

247 REFERENCES

248

249 1. Ilondu EM, Iloh AC. Inhibition of Three Fungal Isolates from Sorrel Drink (Zobo) using Hurdle
250 Technique. *World Journal of Agricultural Science*. 2007;3(3):339-343.

251

252 2. Adebayo-Tayo BC, Samuel UA. Microbial Quality and Proximate Composition of dried *Hibiscus*
253 *sabdariffa* Calyces in Uyo, Eastern Nigeria. *Malayas. Journal of Microbiology*. 2000;5(1):13-18.

3. Yadeng Q, Chin KL, Malekian F, Berhane M, Gager J. Biological Characteristics Nutritional and Medicinal values of Roselle, *Hibiscus sabdariffa*. Circular-Urban Forestry Natural Resources and Environmental No. 604, 2005.
4. Lee CH, Kuo CY, Wang CJ, Wang CP, Lee YR, Hung CN. A polyphenol extract of *Hibiscus Sabdariffa* L. ameliorates acetaminophen-induced hepatic steatosis by attenuating the mitochondrial dysfunction in vivo and in vitro. *Bioscience, Biotechnology, and Biochemistry*. 2012;76(4): 646–651.
5. Gurrola-Diaz CM, Garcia-Lopez PM, Sanchez-Enriquez S, Troyo-Sanroman R, Andrade-Gonzalez I, Gomez-Leyva JF. Effects of *Hibiscus sabdariffa* extract powder and preventive treatment (diet) on the lipid profiles of patients with metabolic syndrome (MeSy). *Phytomedicine: International Journal of Phytotherapy and Phytopharmacology*. 2010;17(7):500 – 505.
6. Adesokan IA, Abiola IP, Adigun MO, Anifowose OA. Analysis of Quality attributes of *Hibiscus sabdariffa* (zobo) drinks blended with aqueous extract of ginger and garlic. *African Journal of Food Science*. 2013;7(7):174 -177.
7. Ogiehor IS, Nwafor OE. Associated microbiological, biochemical and chemical quality changes in zobo beverage produced from *Hibiscus sabdariffa* Linn. *Nigerian Annals of Natural Sciences*. 2004;5: 1-10.
8. Kolawole JA, Okeniyi SO. Quantitative Mineral Ion Content Of A Nigerian Local Refreshing Drink – "Zobo" – (Water Extract of *Hibiscus sabdariffa* Calyx). *Research Journal of Pharmacology*. 2007;1(1): 23-26.
9. Hirunpanich V, Utaipat A, Morales NP. Hypocholesterolemic and antioxidant effects of aqueous extracts from the dried calyx of *Hibiscus sabdariffa* L. in hypercholesterolemic rats. *Journal of Ethnopharmacology*. 2006;103(2):252-260.
10. Carvajal-Zarrabal O, Waliszewski SM, Barradas-Dermitz DM. The consumption of *Hibiscus sabdariffa* dried calyx ethanolic extract reduced lipid profile in rats. *Plant Foods for Human Nutrition*. 2005;60(4):153-159.
11. Farombi EO, Ige OO. Hypolipidemic and antioxidant effects of ethanolic extract from dried calyx of *Hibiscus sabdariffa* in alloxan-induced diabetic rats. *Fundamentals of Clinical Pharmacology*. 2007;21(6):601-609.
12. Alarcon-Aguilar FJ, Zamilpa A, Perez-Garcia MD. Effect of *Hibiscus sabdariffa* on obesity in MSG mice. *Journal of Ethnopharmacology* . 2007;114(1):66-71.
13. Liu JY, Chen CC, Wang WH, Hsu JD, Yang MY, Wang CJ. The protective effects of *Hibiscus sabdariffa* extract on CCl4-induced liver fibrosis in rats. *Food and Chemical Toxicology*. 2006;44(3):336-343.
14. Egbere OJ, Anuoye PF, Chollom PF, Okpara, PV. Effects of Some Preservation Techniques on the Quality and Storage Stability of Zobo Drink (Nigeria, Non- alcoholic Beverage from *Hibiscus sabdariffa*). *Journal of Food Technology*. 2007;5(3):225-228.
15. Essien E, Monago C, Edor EA. Evaluation of the Nutritional and Microbiological Quality of Kunun (A Cereal Based Non-Alcoholic Beverage) in Rivers State, Nigeria. *The Internet Journal of Nutrition and Wellness*. 2011;10(2): 1-10.
16. Hirschbruch MD, Torres EA. Toxicology de Alimentos: *Uma Discussao. Hig-Alim.*, 1998;12(53):21-25.
17. Ogundapo SS, Onuoha JC, Olekanma CN, Okon AB, Soniran OT, Omoboyowa DA. Alteration in biochemical parameters of albino Rats treated with aqueous extract of *Hibiscus sabdariffa* calyces

313 (zobo) supplemented with commercial flavor additive. *Journal of Natural Products*. 2014;7(2014):116-
314 123

315

316 18. National Research Council. Guide for the Care and Use of Laboratory Animals. A report of the
317 Institute of Laboratory Animal Resources Committee on Care and Use of Laboratory Animals. NIH
318 Pub. No. 85-23; 1985. Washington, D.C.:U.S. Department of Health and Human Services.

319

320 19. Rietman, S. and Frankel, S. Colorimetric Assay of Alanine, Aspartate Aminotransferase. *American*
321 *Journal of Clinical Pathology*. 1957;28:56.

322 20. Rec GS. Colorimetric method for serumalkaline phosphatase determination. *Journal of Clinical*
323 *Biochemistry*. 1972;10(2):182

324 21. Jendrassik and Grof. Clinical Chemistry Method of Bilirubin Estimation. 2nd Edition, Chang and
325 Hall London. 1999;358 – 62.

326 22. Tietz NW. Fundamentals of Clinical Chemistry. Saunders, W. B. Company Philadelphia, PA. 1995;
327 pp 874.

328 23. Doumas BT, Watson W, Biggs HG. Albumin Standards and Measurement of Serum albumin with
329 Bromocresol Green. *Clin.Clim. Acta*, 1971;31:87-96

330 24. Fawcett JK, Scott JE. A Rapid and Precise method for the Determination of Urea. *Journal of*
331 *Clinical Pathology*. 1960;13(2):156-9.

332 25. Butler AR. The Jaffe's reaction identification of coloured species. *Clin. Chem. Acta*. 1975;59:227-
333 232.

334 26. Maruna RFL. Determination of Sodium in Serum. *Clinical Chemistry, Acta*. 1958;2(1):281.

335 27. Trinder P. Determination of Sodium in Serum. *Analyst*. 1951;76:596.

336 28. Satpal, Jain SK, Punia JS. Studies on biochemical changes in subacute thiodicarb toxicity in rats.
337 *Toxicology International*. 2010;17: 30-32

338 29. Tseng TH, Kao TW, Chu CY, Chou FP, Lin WL, Wang CJ. Protective effects of dried flower
339 extracts of *Hibiscus sabdariffa* L. against oxidative stress in rat primary hepatocytes. *Food and*
340 *Chemical Toxicology*. 1997;35(12):1159-64.

341 30. Renee JK. Interactions of Biomolecules with Selected Proteins. 6th Edition. McGraw, London.
342 2002:591 – 3.

343 31. Kolawole OT, Akiibinu MO, Akanji MA. Assessment of the effect of aqueous extract of calyx of
344 *Hibiscus sabdariffa* on some biochemical indices of renal function in rats. *International Journal of*
345 *Pharma Sciences*. 2014;4(3):587-590.

346 32. Emelike CU, Dapper DV. Biochemical Effects of Oral Administration of Aqueous Extract of
347 *Hibiscus sabdariffa* on Wistar Albino Rats. *IOSR Journal of Dental and Medical Sciences*. 2013;9(1):
348 35-39.

349 33. Gosselin RE, Hodge HC, Smith RP, Gleason MN. Clinical Toxicology of Commercial Products. 4th
350 Edition. Baltimore: Williams and Wilkins. 1976;11-243.

351 34. Mojiminiyi FBO, Adegunloye BJ, Egbeniyi YA, Okolo RU. An investigation of the diuretic effect of
352 an aqueous extract of the petals of *Hibiscus sabdariffa*. *African Journal of Medicine and Medicinal*
353 *Sciences*. 2000;2(1):77-80.

354 35. Adigun MO, Ogundipe OD, Anetor JI, Odetunde AO. Dose-dependent changes in some
355 haematological parameters during short-term administration of *Hibiscus sabdariffa* calyx aqueous

- 356 extract (zobo) in Wistar albino rats. *African Journal Medicine and Medicinal Science*. 2006;35(1):73-
357 77.
- 358 36. Goldstein RS, Schenellmann RG. Toxic responses of the kidney: in Casarett and Doull's toxicity.
359 McGraw-Hill Co. London, 5th edition. 1999;417-422.
- 360 37. Orisakwe OE, Husaini DC, Afonne OJ. Testicular effects of sub-chronic administration of *Hibiscus*
361 *sabdariffa* calyx aqueous extract in rats. *Reproductive Toxicology*. 2003;18(2):295-8.