

PROXIMATE COMPOSITION, MINERAL AND PHYTOCHEMICAL CONTENTS OF SOME LEAFY VEGETABLES NATIVE TO IGALA KINGDOM KOGI STATE NIGERIA.

+-

ABSTRACT

The proximate composition, minerals and phytochemical contents of some leafy vegetables native to Igala land were determined. It was designed to establish their nutritional value and inclusion in food composition table for use by nutritionists and dieticians. Fresh vegetable samples (cocoyam, egg plant leaf, sweet potato leaf, fluted pumpkins, camwood, spinach, drum stick, jute mallow, basil plant leaf and bitter leaf) were obtained from gardens and vegetable shops at Anyigba. They were washed dried and standard methods were used for the analysis. Triplicate values were obtained in each case, expressed on dry weight basis. All data obtained were analyzed statistically using SPSS version 17.0 package. Means and standard deviation were calculated at a significant level of $p \leq 0.001$. Among the vegetable samples, crude fibre contents ranged from (26.7 to 8.0 %). Ash (15.2 to 5.1 %), moisture (5.3 to 1.52 ± 0.02 %), fat 4.1 to 1.3 %, protein (25.1 to 9.2 %) and carbohydrates, (67.2 to 8.04, Phytochemical contents (mg/100 g) included; Phenol (312.6 to 10.1) flavonoid was (420.7 to 2.6,) alkaloid (9.7 to 2.6), oxalate 0.9 to 0.003. Tannin (36.20 to 0.03), Saponin (5.20 to 0.0), Phytates (2.02 to 0.06). Cyanides 0.5 to 0.05. Mineral contents mg / 100 g included: sodium (0.08 to 0.001) potassium (0.3 to 0.002) calcium (0.06 to 0.02) magnesium (0.06 to 0.00) and phosphorus (0.06 to 0.02). Drum stick contained highest crude fibre, in protein, Potassium, Magnesium and Phytate contents while Ash content was higher in fluted pumpkins. For all the leafy vegetables considered, they contain Phytochemicals within allowable level for safety.

KEY WORDS: chemical analysis, proximate, phytochemical, minerals, leafy vegetables.

INTRODUCTION

Leafy vegetables are low in fat and high in protein and dietary fibre, they are rich sources of minerals (Calcium, Potassium and Magnesium) vitamins and high in phytochemicals [1]. Vegetables contain bioactive compounds that protect the body from nutritional deficiency diseases [2] and free radicals that cause oxidative damage to cells. Vegetables are important food both from economic and nutritional stand points. Their nutritive significance is their richness in minerals which is very essential in the maintenance of human health [3], vegetables are an important component of healthy diet if consumed daily in sufficient amounts, could help prevent major diseases such as cardiovascular diseases and certain cancers [4]. They are at their best when tender or succulent, many of them can easily be grown in home gardens and thus provide economic access and convenient sources of fresh, unprocessed vegetables with active nutrients [5]. In Nigeria; vegetables are used in dishes for soups (melon or egusi, ogbono (*Trvingia gabonensis*) and vegetable soups for the consumption of the main dishes (fufu, pounded yam, garri and akpu). Other main dishes include; white rice, bean, corn meal, yam porridge, semolina and others.

Most of these are potent bioactive compounds found in medicinal plant parts that can be used for therapeutic purpose or which are precursors for the synthesis of useful drugs [6]

The feeling of fullness produced by intake of vegetables helps in controlling overweight and obesity since they are low in calories and high in fibres. From the economic point of view, vegetables quench hunger due to their satiety factor [7]. Some of the green leafy vegetables native to Igala land are cocoyam leaf (*Colocacia esculentus*), egg plant leaf (*Solanum incamum*), bitter leaf (*Vernonia amygdalina*), sweet potato leaf (*Ipomoea batatas*), fluted Pumpkin leaf (*Telferia occidentalis*) others include; camwood leaf (*Pterocaptus soyanxii*), +spinach (*Amaranthus chlosostachys*), drumstick (*Moringa oleifera*) jute mallow (*Cochorus olitouis*) and basil plant leaf (*Occimum bacilicum*) [8]. Therefore, this work aimed at establishing the quantitative nutrient contents, Phytochemical and minerals in the leafy vegetables available in Igala land, some of which are often neglected.

Materials and methods

Sources of vegetables

Ten leafy vegetable samples were harvested and purchased from gardens and shops of Igalas, of which drum sticks, sweet potato leaf, basil plant leaf, cocoyam leaf and jute mallow were harvested from gardens adjacent to various homes of Igalas, others (egg plant leaf, camwood leaf, spinach, fluted pumpkins, and bitter leaf) were procured from only Igala vegetable shops. Both harvested and purchased samples were identified by experts.

Preparation of samples

Samples were trimmed, washed and drained as described by [9] and air dried to prevent loss in nutrients and color. Each sample was milled into fine powder using type 8 k 31 Kenwood blender and packaged in plastic bags to prevent moisture absorption, contamination and also labeled for easy identification and kept refrigerated at $10 \pm 2^{\circ}\text{C}$; that means the temperature of the storage could be less than 10 by 2°C which is 8°C or greater than 10 by 2°C which is 12 , it mean the range of the temperature was between 8°C to 12°C for not longer than two weeks prior to proximate, phytochemical and minerals analysis, using standard methods.

Analytical methods:

Proximate composition analyses

Ash content determination

Ash content of the finely sieved sample was determined by incineration at 600°C in muffle furnace as described by [10]. Triplicate values were obtained for each sample, two grams (2.0 g) of each sample was weighed into a weighed and ignited tarred crucible (w_1).

The crucible and weighed sample were placed on a hot plate inside a fume cupboard to prevent smoke accumulation, the remaining residue was transferred to a pre - heated muffle furnace and maintained at 600°C for 6 hours to ash until the sample was reduced to a light ash, the crucible was removed, placed in the desiccators, cooled and weighed (w_2) and the ash content was calculated:

$$\% \text{ Ash (on dry basis)} = \frac{W_2 - w_1 \times 100}{2.0\text{ g}}$$

Fat content Determination

Fat content was estimated using soxhlet extraction method; according to [11] 2.0 g of sample was weighed into a thimble (w_1). Dried and cooled boiling flask was weighed, filled with 300 ml (w_2) petroleum ether and boiled at 60°C with the extraction thimble in soxhlet apparatus which was allowed to reflux for 6 hours. The thimble was carefully removed, while the extracted oil in the petroleum ether flask was dried at $105\text{--}110^{\circ}\text{C}$ for 1hour. It was then transferred from the oven to the desiccators, cooled, weighed and calculated as below;

$$\% \text{ fat} = \frac{w_2 - w_1 \times 100}{\text{Weight of sample (2.0 g)}}$$

Crude Protein Determination

The crude protein of the sample was determined using the micro-Kjeldahl method described by [12] 2 g was weighed along with 20 cm^3 of distilled water into a micro - Kjeldahl digestion flask. It was shaken and allowed to stand for some time. One tablet of selenium catalyst and 20 cm^3 tetra oxo sulphate (vi) acids were added. The flask was heated on the digestion block at 100°C for 4 hours until the digest became clear. The flask was removed from the block and allowed to cool. Content was transferred into 50 cm^3 volumetric flask and diluted to the mark with water.

An aliquot of the digest (10 cm^3) was transferred into another micro-Kjeldahl flask along with 20 cm^3 of distilled water, placed in the distilling outlet of the micro - Kjeldahl distillation unit. A conical flask containing 20 cm^3 of boric acid indicator was placed under the condenser outlet. Sodium hydroxide solution (20 cm^3 , 40 %) was added to the content in the Kjeldahl flask by opening the funnel stopcock. The distillation started and the heat supplied was regulated to avoid sucking back. Distillate was

collected in 20 cm³ boric acid and the distillation was stopped. Nitrogen in the distillate was determined by titrating with 0.014M of H₂SO₄; the end point was obtained when the color of the distillate changed from green to pink.

Calculation;

% Crude protein = % N x 6.25 (conversion factor)

The nitrogen content of the sample is given by the formula below.

% N = TV x Na x 0.014 x V1 x 100 x G x V2

Where TV = Titre value of acid (cm³)

Na = Normality of acid

V1 = Volume of distilled water used for distilling the digest (50 cm³).

V2 = Volume of aliquot used for distillation (10 cm³)

G = Original weight of sample used, g

Determination of moisture contents

Moisture content of each sample was determined as described by [13] using vacuum oven method; 2.0 g of the sample was rapidly weighed into a pre-weighed dried dish (w₁), and weighed with the dish (W₂) It was dried to a constant weight at 95-100°C at a pressure that was not exceeding 100 mHg for 5 hours when the drying procedures was completed, the dish was placed in the desiccators to cool and re-weighed (W₃) and the recorded loss in weight, was the moisture. The percentage moisture was calculated as below;

$$\% \text{ moisture} = \frac{W_2 - W_1 \times 100}{W_3 - W_1}$$

Where; W₁ = initial weight of the empty crucible

W₂ = weight of the crucible plus (+) the sample before drying.

W₃ = final weight of crucible + sample after drying

% total solid (dry matter) = 100 - % moisture.

Crude Fiber Determination

A non-enzymatic method of [14] was used to determine crude fiber content, 2.0 g of the dried sample was defatted with petroleum ether and boiled under reflux for 30 minutes with 200 ml of solution containing 1.5g Of H₂SO₄ per 100 ml of the solution. The solution was filtered through linen on a fluted funnel and washed with boiling water until the washing was no longer acidic. The residue was transferred to a beaker and boiled for 30minutes in 200 ml of solution containing 1.25 g of carbonate-free NaOH per 100 ml. Final residue was filtered through a thin but closed pad of washed and ignited asbestos in a gooch crucible. It was dried in electric oven, weighed, incinerated, cooled and re-weighed;

The loss in the weight after incineration x 100 was calculated as the percentage (%) of the crude fiber.

$$\% \text{ crude fiber} = \frac{\text{Loss in weight (g)} \times 100}{\text{Original mass (2.0)}}$$

Carbohydrate content determination

Carbohydrate content was determined by difference as described [15] where the total proportion of carbohydrate in the leaves sample was obtained by calculation, using the percentage weight method by subtracting the % sum of food nutrients: (% protein, % crude fiber, fat % and % ash %) from 100 %.

Where, % percentage of carbohydrates (=) (CF + CP + F + A + M – 100 %) where; CF = Crude fibre, CP= Crude Protein, M = Moisture, F = Fat and A = Ash.

Note: Triplicate values were obtained for each sample.

Phytochemical analysis procedures

Phytic acid Determination

Phytic acid was determined using the procedures described by [11]. 2.0 g of each sample was weighed into 250 ml conical flask. 100 ml of 2 % concentrated hydrochloric acid was used to soak the sample in the conical flask for 3 hours. It was filtered through a double layer of hardened filter papers. 50 ml of

each filtrate was placed in 250 ml beaker and 107 ml of distilled water was added in each case to give proper acidity. 10 ml of 0.3 % Ammonium thiocyanate solution indicator was added. It was titrated with standard iron chloride solution, which contained 0.00195 g iron per ml; the end point was slightly brownish-yellow, this persisted for 5 minutes. The percentage (%) phytic acid was calculated using the formula;

% phytic acid = r. x 1.19x100 where x = titer value x 0.00195 g.

Oxalate Estimation.

Oxalate determination was carried out using methods described by [11] 2.0 g of each sample was boiled in 40 ml distilled water for 30 minutes in a reflux condenser. A total of 10 ml of 20 % Na_2CO_3 was added and boiled for another 30 minutes. The resulting mixture was filtered and the residue was washed severally with hot water until the p^{H} was neutral, the residue was washed with hot water until the wash water stopped showing any alkaline reaction. The filtrates were concentrated to a small volume and cooled with constant stirring. Hydrochloric acid (HCl) was added, (1:1) drop wise until the final acid concentration after neutralization was 1 %. The precipitate was allowed to flocculate and the extract carefully filtered into a 250 ml flask and made-up to mark. It was kept overnight and supernatant liquid was filtered through a dry filter paper in a dry beaker. An aliquot of this filtrate was taken into a 400 ml beaker, diluted with water to 200 ml and made just ammonia cal to re-acidify with acetic acid. In the cold medium, 10 ml of 10 % calcium chloride solution was added and stirred well to induce calcium oxalate precipitate to appear, it was allowed to stay overnight. The clear supernatant liquid was decanted off through whatman No 42 filter paper; the precipitate was dissolved in HCl acid (1:1). Oxalic acid was re-precipitated by adjusting the pH with ammonia hydroxide solution. Content was boiled and allowed to settle overnight. Oxalic acid was determined by titrating against 0.05N KMnO_4 solution.

Calculation;

1ml of 0.05N KMnO_4 = 0.00225 g anhydrous oxalic acid.

% Oxalic acid = $\frac{\text{Titre value} \times 0.00225}{2} = \text{T.U} \times 0.1125$.

Where, T. U = $\frac{\text{Titre value}}{2}$

Tannic Acid Determination

The total tannin content was determined according to method described by [16]. This involved the use of butanol /HCl reagent (95.5 w/v and 37 % HCl) and ferric reagent (2 % ferric ammonia sulphate in NH_4Cl) where 0.5 ml of extract was diluted with 70 % acetone to make absorbance not to exceed 0.6 in a tube of 30 ml. and incubated at 97-100 °C for 60 minutes. The tube was cooled and read absorbance at A550 nm using unheated suitable blank and condensed tannin (96 % of dry matter) and calculated using; $A_{550\text{nm}} \times 78.26 \times \text{dilution factor}$.

Where; 1cm, 550 nm of leucocyanidin = 406. (The formula assumed that 1 cm, 550 nm of leucocyanidin is equal to 460) and A = absorbance

Flavonoid determination

Flavonoid was determined using spectrophotometer (Aluminum chloride method) as described by [17], 9 ml of the sample was weighed and treated with ethyl acetate and mixed with 1 ml of Aluminum chloride (AlCl_3) and methanolic solution (2 % w/v) incubated at a room temperature for 15 minutes, the absorbance was read at 430 nm. The amount of the TFC (total flavonoid contents) was estimated from the standard calibration curve of 10-100 $\mu\text{g ml}^{-1}$ quercetin.

Saponin Contents Determination

Saponin content was determined as described by [18]. 5 g of each sample was dispersed in 50 mL of 20 % ethanol. The suspension was heated over a hot water bath for 4 hours with continuous stirring at 55 °C. The mixture was filtered and the residue re- extracted with another 200 ml of ethanol prepared in distilled water (20 % v/v). The combined extracts were reduced to 40 ml over water bath at 90 °C and

the concentrate was transferred into 250 ml separator funnel, 20 ml of diethyl ether was added and shaken vigorously. The aqueous layer was recovered while the ether layer was discarded. The purification process was repeated and 60 mL of n-butanol was added and the mixture was washed twice with 10 ml of 5 % aqueous sodium chloride. The remaining solution was heated in a water bath at 90 °C. Samples were dried in an oven at 100 °C until a constant weight was obtained and Saponin content was calculated in percentage.

$$(\%) \text{ Saponin} = \frac{\text{absorbance of sample} \times \text{gradient factor} \times \text{Dilution factor}}{\text{Wt of sample} \times 10,000}$$

Determination of phenolic acid

Total phenol contents was determined using the procedures described by [19] 1 ml aliquot of the extracts were taken into 10 ml glass tube and made up to a volume of 3 mL with distilled water, 5mL of folin ciocalteau reagent (1:1) with water and 2 ml. 20 % of Na₂CO₃ were added sequentially in each tube. Blue color was developed in each tube because the phenols undergo a complex redox-reaction with phosphomolibdic acid in folin ciocalteau reagent in alkaline medium, which resulted in blue colored complex molybdenum blue. The test solutions were warmed for 1 minute, cooled and absorbance was measured at 650 nm against the reagent, used as blank. A standard calibration plot was generated at 650nm using a concentration of catechol. The concentration of phenol in the test samples were calculated from the calibration plot and expressed as mg catechol equivalent of phenol/100 g of sample.

Alkaloid Determination

Alkaloid was determined using gravimetric method, [20], 2.0 g of sample was weighed (w₁) and dispensed into 50 ml of 10 % acetic acid solution in ethanol. The mixture was well shaken and allowed to stand for 4 hours and filtered. The filtrate was evaporated to ¼ of its original volume. Concentrated NH₄OH was added and alkaloid precipitated. The filter paper was weighed (w₂) and the precipitate was scrapped off and washed with 1 % NH₄OH solution. This was weighed with filter paper (w₃); the precipitate was dried in a filter paper in the oven at 60 °C for 30 minutes and reweighed (w₄). By weight difference, the weight of the alkaloid was determined and expressed as a percentage of sample weight analyzed, given by formula

$$\% \text{ Alkaloid} = \frac{W_1 - W_2 \times 100}{W_3 - w_4}$$

Where; W₁ = Weight of sample

W₂ = Weight of empty filter paper

W₃ = Weight of filter paper+ precipitate before drying.

W₄ = weight of filter paper+ precipitate after drying

Triplicate values were obtained for each sample

Cyanide determination

Total cyanide was determined by the pH-differential method as described by [21] two buffer system: potassium chloride buffer pH 1.0 M and sodium acetate buffer p^H 4.5 M . 0.4 ml of the extract was mixed with 3.6 ml of corresponding buffers and read against a blank at 510nm and 700 nm, Absorbance (A) was calculated as follows; A = (510 – A700) pH 1.0 – (510 – A700) pH 4.5 Cyanide (mg/l) = A x MW x DF x 1000 (Mn x 1.0).

Where, A = absorbance

MW = molecular formula (449.2)

DF = dilution factor

MA = molecular absorptive (26, 900)

Mineral Analysis;

Sodium, Potassium, Calcium and Magnesium Mineral contents of the samples were determined using triple acid digestion method of [22]. 2.0 g of the samples were weighed into a micro-Kjeldahl digestion flask to which 24 cm³ of mixture of concentrated HNO₃, H₂SO₄, and 60 % HClO₄ (9 : 2 : 1 v/v) were

added. The flask was put on the heating block and digested to a clear solution, cooled and the content was transferred into 50 cm³ volumetric flask and made-up to the volume mark with water.

Phosphorus Determination

Phosphorous was determined using [23] methods. Digested supernatant solution (2 cm³) was placed into 50 cm³ volumetric flask, 2 cm³ of sample and 2cm³ of ammonium molybdate solution were added with distilled water to make-up to 48 cm³, the content was mixed and 1 cm³ of diluted stannous chloride solution was added and mixed, then 1cm³ of distilled water was added to make-up to 50 cm³ marked and left to stand for 5 minutes. The % absorbance on spectrophotometer at 660 nm wavelength was used to determine the concentration of phosphorus.

Triplicate values were obtained for each sample

Statistical analysis

All data obtained were analyzed statistically, using SPSS version 17.0 package. Means were separated using [24] multiple comparison test at P < 0.05.

Results

Proximate composition of the studied leafy vegetables, Table 1 represents the proximate composition of (moisture, ash, crude fibre, protein and carbohydrates) contents of ten (10) leafy vegetables native to Igala kingdom Kogi state. Ash contents ranges between (15.2 to 5.1 %), Crude fiber contents (26.7 g to 8.0 %), moisture (5.3 to 1.52 %), fat (4.1 to 1.3 %), proteins (25.1 to 9.2 %) and carbohydrates, (69.49 to 40.2.04 g)

Table 1: Proximate composition of ten vegetables native to Igala land (%) dry basis

SAMPLE CODE	MOISTURE	ASH	CRUDE FIBER	FATS	PROTEIN	CABOHYDRATE
212	2.7±0.03 ^a	5.1±0.01 ^a	8.05 ± 0.05 ^a	2.2±0.17 ^b	9.2±0.1 ^b	69.48±0.008 ^a
213	5.3±0.2 ^a	12.1 ± 0.2 ^a	11.2 ± 0.8 ^b	2.1 ± 0.0 ^b	20.1 ± 1.0 ^b	49.2 ± 0.11 ^a
214	1.8±0.04 ^a	9.2±0.11 ^a	12.1 ± 0.08 ^b	2.1±0.06 ^b	20.10± 5.00 ^b	54.5±0.3 ^a
215	2.7±0.17 ^a	6.7±0.2 ^b	15.1 ± 0.1 ^a	1.9±1.6 ^b	9.11±0.1 ^b	64.1±0.3 ^a
216	1.52±0.02 ^b	3.4±1.05 ^a	26.7±0.2 ^a	3.02±0.0 ^a	25.1±1.1 ^b	40.2±0.9 ^a
217	1.77±0.3 ^a	7.2±0.3 ^a	12.18±0.2 ^a	1.5±0.07 ^b	10.10±0.2 ^b	67.3±0.1 ^a
218	2.5 ± 0.01 ^a	15.2±0.1 ^a	12.2±0.06 ^a	1.80±0.07 ^b	14.5±0.2 ^b	53.70±0.05 ^a
219	4.6±0.01 ^a	7.5±0.03 ^b	18.1±0.04 ^a	4.1±0.0 ^a	11.30 ± 0.2 ^b	54.1±0.2 ^a
220	5.1±0.1 ^a	5.9±0.1 ^c	11.8±0.9 ^b	1.3±0.09 ^b	13.4±0.01 ^b	62.5±0.02 ^a
221	1.6±0.04 ^b	11.2±0.05 ^a	11.6±0.3 ^b	1.9±1.59 ^b	23.4±0.06 ^b	50.1±0.2 ^a

Values represent means of triplicate values ± sd (standard deviation)

Sample means with the same superscripts in a Column are not significantly different p≥0.05

SAMPLE CODE

SAMPLE CODE

SAMPLE CODE

212 = (BASIL PLANT LEAF)

218 = (FLUCTED PUMPKINS)

a = (significant ≥0.01)

213 = (BITTER LEAF)

219 = (JUTE MALLOW)

b = (not significant ≥0.05)

214 = (CAMWOOD PLANT LEAF)

220 = (SPINACH)

c = weak significant

215 = (COCOYAM LEAF)

221 = (SWEET POTATOE PLEAF)

± = Data expressed as mean ±

216 = (DRUMSTICK)

217 = (EGG PLANT LEAF).

276 Table 2: Phytochemical contents of ten vegetables common in Igala land (mg/100) powdered
277 Sample

278	Values represent means of triplicate values $\pm$ sd (standard deviation)		
279	Sample means with the same superscripts in a Column are not significantly different $p \geq 0.05$		
280	SAMPLE CODE	SAMPLE CODE	SAMPLE CODE
281	212 = (BASIL PLANT LEAF)	218 = (FLUCTED PUMPKINS)	a = (significant ≥ 0.01)
282	213 = (BITTER LEAF)	219 = (JUTE MALLOW)	b = (not significant ≥ 0.05)
283	214 = (CAMWOOD PLANT LEAF)	220 = (SPINACH)	c = weak significant
284	215 = (COCOYAM LEAF)	221 = (SWEET POTATOE PLEAF)	$\pm$ = Data expressed as mean $\pm$
285	216 = (DRUMSTICK)		
286	217 = (EGG PLANT LEAF).		

290

Sample code	Sodium mg	potassium mg	Calcium mg	Magnesium mg	Phosphorous mg
212	0.02±0.005 ^b	0.1±0.01 ^b	0.06±0.01 ^c	0.04±0.30 ^b	0.043±0.03 ^a
213	0.006±0.03 ^b	0.06±0.02 ^b	0.02±0.006 ^b	0.03±0.006 ^b	0.02±0.01 ^b
214	0.002±0.04 ^b	0.02±0.03 ^b	0.02±0.003 ^b	0.002±0.03 ^b	0.04±0.015 ^a
215	0.080±0.01 ^b	0.08±0.015 ^b	0.02±0.003 ^c	0.02±0.006 ^b	0.03±0.01 ^b
216	0.009±0.01 ^b	0.27±0.11 ^b	0.02±0.005 ^c	0.06±0.02 ^b	0.03±0.20 ^b
217	0.001±0.003 ^b	0.3±0.06 ^b	0.05±0.005 ^b	0.05±0.0005 ^b	0.03±0.005 ^b
218	0.001±0.003 ^b	0.08±0.06 ^b	0.02±0.006 ^c	0.02±0.01 ^b	0.30±0.06 ^a
219	0.01±0.04 ^b	0.07±0.01 ^b	0.03±0.07 ^b	0.04±0.02 ^b	0.03±0.01 ^a
220	0.002±0.005 ^b	0.1±0.01 ^b	0.02±0.01 ^c	0.02±0.005 ^b	0.06±0.01 ^b
221	0.003±0.01 ^b	0.1±0.01 ^b	0.02±0.003 ^b	0.04±0.05 ^b	0.02±0.01 ^a

Values represent means of triplicate value± sd (standard deviation)

Sample means with the same superscripts in a Column are not significantly different p≥0.05

SAMPLE CODE

SAMPLE CODE

SAMPLE CODE

212 = (BASIL PLANT LEAF)

218 = (FLUCTED PUMPKINS)

a = (significant ≥0.01)

213 = (BITTER LEAF)

219 = (JUTE MALLOW)

b = (not significant ≥0.05)

214 = (CAMWOOD PLANT LEAF)

220 = (SPINACH)

c = weak significant

215 = (COCOYAM LEAF)

221 = (SWEET POTATOE PLEAF)

± = Data expressed as mean ±

216 = (DRUMSTICK)

217 = (EGG PLANT LEAF).

Discussion

Proximate composition of leafy vegetable samples

Proximate composition of ten (10) studied leafy vegetables (%) on dry basis, showed drumstick [coded sample 216] contained highest amount of crude fiber and protein, with (26.7 ±0.2) and (25.1±1.1) values respectively. The fiber content signified that the leaves of drumstick can be used to reduce incidence of duodenal ulcer. The findings agree with the results of [25] in a study where intake of high fiber diet of 28.2 g / day for 6 months lowered incidence of duodenal ulcer recurrence. Also, [26] reported that high crude fibre in jute mallow decreased blood sugar level and cardiovascular diseases. [27] Reported that a total dietary fibre of 20 g to 25 g / day can prevent obesity, diabetes and cancers, the findings in this study showed that drumstick contained the highest amount of crude fiber therefore, it could be suitable for prevention of duodenal ulcer, hyperglycemia, obesity, diabetes, cancers and other related disorders. The significant difference at p≤0.001 for fibre content at various levels indicated that the samples are good source of vegetable exchange list that can be used in cases of allergy vegetables; vegetables studied were classified based on macro/micro nutrients and chemical composition, which establishes that 100 g of some of the vegetables studied contain nutrients (ash, flavonoid, phenolic acid, phytate, Saponin, carbohydrates and fibre) and were significant at p ≥ 0.01, while others contains (protein, oxalate, sodium and alkaloid) which were not significant at p ≥ 0.05, based on the nutrient components of the studied vegetables at various level, that are significant at p ≥ 0.01 can be alternated with the vegetables that are not significant at p ≥ 0.05 if reactions from allergen occur on consumers. The Recommended Dietary Allowance (RDA) for protein is in the range of 28-65 g

for children, lactating mothers, pregnant women and adults [28] also [29] reported that high protein of 1.8 g / kg is required in postoperative nutrition, the protein content of [25.1±1.1] in drumstick in this current study is reasonably high and adequate. Therefore, it could be referenced for use in disease condition and for management of other nutritional related diseases. Fluted pumpkin has the highest amount of Ash [15.2] and drumstick [3.4±1.05] the least. The highest moisture content occurred in bitter leaf [5.3±0.2] and lowest in drumstick (1.52±0.02.). Vegetable studied generally have low moisture which is an index of extended shelf life also according to [30] and [31] Low moisture content after dehydration provides concentrated nutrients while high moisture enhances water activities that can increase spoilage therefore; the low moisture content of the leafy vegetables studied can encourage the keeping quality through prevention of the growth of microorganisms, and conservation of nutrients that can prevent/ reverse nutritional related diseases. Fat contents for all vegetable samples were all low; jute mallow contained the highest amount [4.1±0.0 %] and least in spinach (1.3±0.09 %). The low fat content of the vegetables revealed in this study can be referenced in prevention of chronic diseases such as liver cirrhosis, diabetes and cardiovascular diseases [32] and [33]. Carbohydrates in basil [sample coded 212] is [69.48±0.008 %] which is reasonably high could be suitable for active man of 36-40 years with total intake of 2,800 Kcl /day, the findings in this study is evidenced and also in line with [34].

Phytochemical contents of leafy vegetables;

Phytochemical contents of ten vegetables, showed that bitter leaf has the highest amount of flavonoid [420.7 ± 0.1] but lowest in camwood, phenol in drumstick was the highest [312.6] with spinach being the least while values for all the vegetables were extremely significant at $p \geq 0.01$. Alkaloid in spinach [9.7±0.03] was found to be the highest and camwood leaf the least. However, there existed significant difference at $p \geq 0.01$. Oxalate [0.9 ± 0.02] was highest in camwood but, lowest in spinach. The result showed that tannin was highest in bitter leaf [36.20±0.1] with the least amount in sweet potato but, were not significantly different at $p \leq 0.05$, Saponin [5.20 ± 0.91] was high in bitter leaf while both cocoyam and fluted pumpkins were the lowest, with 0.00 value though significant at $p \leq 0.001$. Drumstick contained highest amount of Phytate [2.02 ±0.01] of which sweet potato was the least; cyanide was [0.5 ± 0.9] highest in egg plant and least amount was found in jute mallow. The contents of Phytochemicals indicated the presence of high anti-nutrients at various concentrations which could be harmful to human if consumed raw but, could be reduced to safety level through processing of the vegetables by various blanching and abrasion [35]. According to [36] in the study of effect of processing methods on the nutritional contents of bitter leaf it was reported that boiling, cooking and other heat processing techniques reduced anti-nutrient (Saponin, tannin, Phytates and alkaloids, oxalate and cyanide) in vegetables. This study showed that other phytochemical with minimal amount are negligible and could not course any harmful effects if consumed by humans. Hypoglycemic effect of bitter leaf on experimental broilers at various levels (5 %, 10 % and 15 %) for 28 days was observed [37]. Sample studied, showed that flavonoid in bitter leaf, sweet potato and spinach high amount of 346.5±0.03 and 350.6±0.4 which could be referenced to reduce the occurrence of diabetes. Consumption of purple sweet potato leaf of 200 g/day (902 mg) of phenol compound for 7 and 14 days decreased lipid per-oxidation and DNA damage in human [38]. In this study, high amount of phenol in drumsticks, basil, fluted pumpkins, cocoyam leaf and egg plant leaf was observed therefore; they are good source of dietary antioxidant defense and can decrease oxidative stress in human.

Mineral contents of the leafy vegetables

Table 3 showed the sodium, potassium, calcium, magnesium, and phosphorous contents of the common leafy vegetables in Igala land. Sodium ranged from [0.08 - 0.001] of which drumstick contained the highest value, potassium (0.3±0.06) with egg plant leaf having the largest amount, calcium (0.06±0.01), basil with the largest contents and spinach the least. While drumstick has the highest magnesium contents (0.06±0.02), comwood plant leaf the least, phosphorus in spinach (0.06±0.0.01)

with highest amount, bitter leaf and sweet potato samples had least values. [39] Reported that reduced sodium intake has a reduced risk ratio on cardiovascular, coronary diseases and stroke, The result of this study produced a low sodium content (0.08/100 mg) at $p \leq 0.001$ if compared to [40] recommended dietary allowance of $\leq 2 \text{ g/day}$ (1.3 g) therefore, the findings in this current studies is recommendable for low/ sodium restricted diets, hypertension and for composition table and in vegetable exchange list. [41] Found that the major component of bitter leaf include; calcium, potassium and sodium and that are the major factors sustaining strong bones therefore, reduces the risk of arthritis. High amount of micronutrients in this study is recommendable in building strong healthy bones and teeth that could reduce and prevent arthritis.

CONCLUSION

The study revealed that most of the vegetables available in Igala kingdom especially drumstick, jute mallow, cocoyam, eggplant leaf, fluted pumping leaf are good sources of crude fiber, phytochemical, minerals and macronutrients which could prevent metabolic and chronic nutritional deficiency disorders and for alleviation of risk factors of chronic diseases.

REFERENCES

1. Oluwalana I.B. (2013), effect of processing methods on the nutritional value of amaranthus hybridus L, Department of Food Science and Technology, Federal University, Akure, P.M.B 704 Ondo State, Nigeria. IJAFS 4, 15: Page 523 - 529
2. Momoh-Aliu. Berikisu (2007) Foods and herbs for Radiant Health Ascent, Bookshops Limited, bookshop House 50/52 broad Street P.O.Box 174 Lagos Nigeria 122:8, Published by CSS Press (A Division of CSS OD Limited),15 Broad Street, Lagos Nigeria.
3. Arogba S.S, (2008) Phenolics: A class of Nature's Chemical Weapons of self preservation inaugural lecture, K. S. U Anyigba.
4. Bolaji, P.T, Komolafe G.O and Allie (2008) *Drying Characteristics of Selected local vegetables. Nigeria Food Journal, Vol. 2 No. 1 PP138-143.*
5. Lydia Bazzano (2004), Fruit and Vegetables for Health Joint FAO/WHO Workshop Kobe, Japan Fruit and vegetables for health: Report of a Joint FAO/WHO Workshop, Kobe, Japan.
6. Fayemi P.O (1999) Nigerian Vegetables, Heinemann Educational Book (Nigerian) Plc. Igbodaro road Jerico P.M.B5205 Ibadan
7. Edeoga, H.O, Omosun G, Ushe L.C (2006) Chemical Composition of Hyptis Suaveolens and Ocimum Gratissimum Hybrids from Nigeria, African Journal of Biotechnol, 5 (10) 892-893.
8. Nzelu I.C (2006) Identification, Composition and Processing of Tropical Food Commodities, Fergu Nwankwo printing services, Enugu Nigeria.
9. Gafa, MK. and Itodo. A.U (2011) proximate composition and mineral contents of hairy indigo leaves, Journal of Pharmaceutical Biological and Chemical Science volume 2 issue 1. department of chemistry, Kebbi state university of science and technology Allero
10. Association of Official Analytical Chemist A.O.A.C (20002) Ash and protein contents Determination.
11. Akinmutimi A.H (2006), *Chemical Analysis of Nutritive Value of Raw and Processed Jack Fruit seeds (Artocaptus heterophyllus), Agricultural journal, Vol 1 page No: 266-271*
12. Association of official Analytical Chemist A.O.A.C (1995), Non-enzymatic methods of Crude protein contents determination official methods of analysis international, 16th edition method 4.6.01(962.09), international, Maryland, USA.
13. Onwuka, G.L. (2005) Food Analysis and Instrumentation Theory and Practice Napthali Prints, Lagos, Nigeria pp 124-125

14. Association of official Analytical Chemist A.O.A.C (1995), official methods of analysis of international, 16th edition method 4.6.01(962.09), international, Maryland, USA. Non-enzymatic methods of Crude Fiber determination.
15. AOAC. (1999). Official methods of analysis. 21st Edition, Association of official analytical chemists. Washington D. C. USA.
16. Okwu, D.E and Emenike I. N (2006) Evaluation of Phytonutrients and Vitamin contents of citrus fruits, international journal of molecular medicine and advance science 2 (1): 1-6
17. Vukovi J, Renate J.G, Darink and Santa V, (2007) flavonoid Assay; pre-validation and application on plantago L species. Acta chim slov 54, page 397-406.
18. Obadoni B.O and Ochuko P. O. (2001). Phytochemical studies and Comparative efficacy of the crude extracts of some homeostatic plants in Edo and Delta States of Nigeria. Global Journal, Pure Appl. Science. 8: page 203-208
19. Khatiwora¹ Elija , Vaishali B. Adsul², Manik M. Kulkarni¹, N.R. Deshpande¹ and R.V Kashalkar¹ (2010) Spectroscopic determination of total phenol and flavonoid contents of Ipomoea carnea, International Journal of ChemTech Research CODEN (USA): IJCRGG ISSN : 0974-4290 Vol.2, No.3, pp 1698-1701
20. Harborne. J.B (1980) Phytochemical method, a guide to modern techniques of and plants analysis London New York chapman and hall.
21. Trenerry V.C, Laco. J and Rochford. S (2008), routine analytical method for use in South Pacific regional laboratories for determining naturally occurring antioxidants in foods,
22. Sahrawat, K. L., Kuaar, G. R. and Rao, J. K. (2002). Evaluation of tarciid and dry ashing procedures for detecting potassium, calcium, magnesium, iron, zinc, manganese and copper in plant materials. Communication of Soil Sciences and Plant Analysis, 33 (1and 2):95 – 102.
23. Eno Jumbo Udo and Trenchard .O. Ibia, (2009) mineral element analysis in plant samples, manuals of soil, plant and water analysis Pgs 101- 120 . Publishers Sibon books limited, flat 15 block 6 fourth avenue festac Lagos
24. (Bonferroni .H. Abdi and Šidák. Z, (2007). Corelation for multiple comparisons" (<http://www.utdallas.edu/herve/Abdi-Bonferroni-pretty>). In N.J. Salkind (ed.). Encyclopedia of Measurement and Statistics. Thousand Oaks, CA: Sage
25. Harshman Miily Ryan and Aldoori Walid (2004) how diet and lifestyle affect Duodenal ulcer, review of evidence. Le College Des College of medicine of Family of Canada.
26. Onwordi Chionyedua .T, Anyoluwa .M and Adedojo D. Wusu (2009) the proximate analysis And mineral composition of three leafy vegetables commonly consumed in Lagos. Nigeria African journal of pure and applied chemistry vol 3 (6) pp 102-107. Tools and Methods for Data Collection in Ethnobotanical Studies of Homegardens
27. World Health Organization (2002) Estimated Calorie Needs per Day by Age, Gender, and Physical Activity Level.
28. Adinortey, M.B., Sarfo, J.K., Quayson, E.T., Weremfo, A., Adinortey, C.A., Ekloh, W., and Ocran, J. (2012). Phytochemical screening, proximate and mineral composition of Launaea taraxacifolia leaves. Res. J. Med. Plant 6: 171-9.
29. Han-Geurts, I.J, Jeekel, J., Tilanus H.W, Brouwer, K.J. (2001) Randomized clinical trial of patient-controlled versus fixed regimen feeding after elective abdominal surgery. British Journal of Surgery; 88(12):1578-82
30. Dewole, A.E., Dewuni, D.F.A., Alabi, J.Y.T. and Adegoke, A. (2013). Proximate and phytochemical of Cola nitida and Cola acuminata. Pakistan Journal of Biological Sciences 16: 1593-1596.

31. (Pallavi Joshi and Dipika Mehta (2010) Effect of dehydration on the nutritive value Drumstick leaves Journal of Metabolomics and Systems Biology Vol. 1(1), pp. 5 -9, Department of Home Science, International College for Girls, Jaipur, Rajasthan India.
32. (Chizuru Nishida¹, Ricardo Uauy, Shiriki Kumanyika and Prakash Shetty (2002) Joint WHO/FAO Expert Consultation on diet, nutrition and the prevention of chronic diseases: process, product and policy implications Department of Nutrition for Health and Development, WHO, Geneva, Switzerland:
33. World Health Organization/Food and Agriculture Organization (2004), Expert Consultation on Diet, Nutrition and the Prevention of Chronic Diseases Report of a Joint WHO/FAO Technical Report Series. No 916 Geneva
34. World Health Organization 2002 Estimated Calorie Needs per Day by Age, Gender, and Physical Activity Level
35. (Yusufu. M and Obiegbuna J .E (2012) effect of processing of methods on the phytochemical Content consumer acceptability of two selected green leafy vegetables , Agricultural Science Research Journal, Department of Food Science and Technology, Federal Polytechnic, Idah Nigeria Department of Food Science and Technology, Nnamdi Azikiwe University, Awka, Nigeria .
36. Yakubu, N., Amuzat, A.O. and Hamza, R. U. (2012) Effect of processing methods on the nutritional contents of bitter leaf (*Vernonia amygdalina*) Department of Biochemistry, IBB University, Lapai. Niger State, Nigeria. American Journal of food and Nutrition Volume 1, Issue 3 (Sep-Oct. 2012).
37. Owen O.J, A.O. Amakiri and Takaribi Botoye ,(2011) sugar lowering effect of bitter leaf (*Vernonia amigdalina*) in experimental broilers chicken. Department of animal science, River state university of science and Technology P.M.B PrtHarcourt – Nigeria. (Department of chemical Pathology) University of Port Harcourt Teaching hospital. Asian clinical Resear
38. Chiao CM, YL Lin, CYO Chen, CY Hsu, MJ shiel Liu JF (2005) antioxidant activities of Purple Sweet potato leaves in human immune response lymphocytes function lyric activity of natural killer cell and antibody production and Gastro enterol, 11(37): page 577-581.
39. World Health Organization (2012) Effect of reduced sodium intake on cardiovascular disease, coronary heart disease, and stroke.
40. World Health Organization,(2012) Guideline: Sodium intake for adults and children Geneva
- 41 . (Mensah. J.K, R.I. Okoli, J.O. Ohaju-Obodo and K. Eifediyi (2011) Phytochemical, nutritional and medicinal properties of some leafy, vegetables consumed by Edo people of Nigeria Department. of Botany, Department of Pharmacology and Therapeutics, and Department of Agronomy, Ambrose Alli University, Ekpoma, Edo State, Nigeria.