

**Influence of Harvesting Stage and Storage Temperature on Nutritional Quality of
Tomato (*Lycopersicon esculentum* Mill) cv. PKM-1**

ABSTRACT

Premise of research. It is found that the nutritional quality of fruits varies with the harvesting stage and the storage temperature. Hence, a study was conducted to evaluate the nutritional quality of a popular tomato cultivar in South India, PKM-1 harvested at different maturity stages and during storage.

Methodology. Tomato fruits were harvested at mature green, yellow-green and red ripe stages and the nutritional quality of fresh tomatoes were determined. The fruits were stored at room temperature and in refrigerator at 6°C and the changes in biochemical parameters were estimated.

Pivotal results. Ascorbic acid, total sugars and lycopene contents increased during ripening and were found to be highest in red ripe stage. Titrable acidity and total phenols were highest in mature green tomatoes and decreased during ripening. Storage studies revealed that ascorbic acid, titrable acidity and total phenols decreased during storage. Total sugars and lycopene contents increased during storage.

Conclusions. Tomato fruits stored at 6°C showed better nutritional value due to lesser change in nutrient content. The study revealed that the nutritional quality of tomato harvested at red ripe stage had the highest nutritional value and a low temperature of 6°C favoured better shelf-life and is ideal for storage of harvested tomato.

KEYWORDS

Tomato, Harvesting stage, Ascorbic acid, Sugars, Phenolics

1. INTRODUCTION

Tomato (*Lycopersicon esculentum*) is one of the most consumed foods throughout the world. The area under tomato cultivation is increasing day by day as it is a short duration crop giving high yield and income to the farmers (Pedapati et al. 2013). Tomato is consumed raw, in salads, or after cooking. Dietary intake of tomato and tomato products containing lycopene has been shown to be associated with a decreased risk of chronic diseases such as cancer and cardiovascular diseases

31 (Agarwal and Rao 2000; Willis and Wians 2003). Tomato also contains other natural antioxidants,
32 including ascorbic acid and phenolic compounds. Antioxidants present in food can largely inhibit the
33 harmful effects of free radicals in humans. Recently, enormous evidence is available to affirm that
34 fruits and vegetables contain high concentrations of bioactive compounds including antioxidants
35 which may be beneficial to health (Kaur and Kapoor 2001; Parashar et al. 2014). Several important
36 changes occur in the ultra-structure of tomatoes during ripening, such as, synthesis of lycopene,
37 production of flavour and aroma compounds, and increase in the ratio of citric to malic acid (Grierson
38 and Kader 1986).

39 The maturity of harvested perishable commodities and the way in which they are handled,
40 transported and marketed affect their storage life and quality (Reid 1985). Immature fruits are more
41 subjected to shrivelling and mechanical damage, and are of inferior quality when ripe. Any fruit picked
42 either too early or too late in its season is more susceptible to physiological disorders and has a
43 shorter life than fruit picked at the proper maturity (Kader 1994). The most important quality criteria for
44 tomatoes are red colour, firm but juicy texture, and good flavour. The metabolism of tomatoes
45 continues, even after their detachment from the plant, when fruits have reached their red stage. They
46 continue to ripen and finally deteriorate to a point where they become valueless (Yanuriati et al.
47 1999). It was observed that total soluble solids (TSS) increased during the initial days of storage,
48 attained a peak value and then decreased (Mohammed et al. 1999; Tigist et al. 2013). Earlier studies
49 reported an increase in sugar-acid ratio during storage of tomato fruits at ambient conditions (Tigist et
50 al. 2013; Missio et al. 2015). During post-harvest storage, starch present in tomato fruits are
51 hydrolyzed to soluble sugar such as glucose, fructose and sucrose (Sood et al. 2011).

52 Normally tomatoes are harvested at light red or breaker stage for distant transportation.
53 Several studies have been conducted on storage behaviour and shelf life of tomatoes at varied
54 temperature and conditions (Goren et al. 2010; Anthon et al. 2011; Missio et al. 2015). However,
55 **information on** the changes in overall nutrients on storage of tomato fruits harvested at different
56 ripening stages on PKM-1 tomato cultivar is scanty. This study was thus, conducted to determine the
57 changes in the nutritional parameters on storage of PKM-1 tomato fruits harvested at mature green,
58 yellow-green and red ripe stages. The objective of this work was to identify the optimum harvest
59 maturity resulting in better quality and longer marketability of tomato.

60 **2. MATERIAL AND METHODS**

2.1. Fruit sampling and storage treatments

PKM-1 tomato variety is cultivated throughout the Southern States of India especially in Tamil Nadu. This tomato variety was released by the Tamil Nadu Agricultural University during the year 1978. It is an induced mutant from a local variety called Annanji. It yields on an average, 32 t/ha in a duration of 135 days. The fruits are flat round with capsicum red color with prominent green shoulders even after ripening and weigh about 60-70 g. Fruits are firm and suitable for long distance transport. Though various research work on agricultural production have been conducted with this cultivar, the changes in biochemical parameters during the ripening process and storage are not studied in detail. Hence, this variety was selected for this study.

A field experiment was conducted during 2009-10 at the orchard in Agricultural College and Research Institute, Madurai, Tamil Nadu, India. The field is situated at an elevation of 152 m (above sea level) with 9°97' N latitude and 78°20' E longitude. A raised nursery bed was prepared by applying 10 kg farmyard manure (FYM), 1 kg neem cake, 50 g VAM, 100 g superphosphate and 10 g Furadon per square metre. Certified seeds of tomato cv. PKM-1 were treated with 10 g *Pseudomonas fluorescens* and 2 g Carbendazim per kg of seeds 24 hours before sowing. Just before sowing, the seeds were treated with *Azospirillum* @ 10 g / 100 g of seeds. Seeds were sown in lines at 10 cm apart in nursery beds and covered with sand. Field was ploughed to fine tilth and was prepared with the addition of FYM 25t/ ha and formed ridges and furrows at a spacing of 60 cm. *Azospirillum* (2 kg/ha) and Phosphobacteria (2 kg/ha) were also applied along with FYM. A basal dose of chemical fertilizers, NPK 75:100:50 Kg/ha, Borax 10 kg and Zinc sulphate 50 kg/ha were applied. **Twenty five days old seedlings of tomato** were transplanted on the sides of ridges at a distance of 45 cm. After 30 days of planting 75 kg N/ha was applied. Since no visible pest attack or disease was noticed, plant protection measures were not taken during the experimental period.

Tomato fruits were harvested at three maturity stages including mature green (turning), yellow-green (half ripen) and red ripe (full ripen) from the trial. Fruits of uniform size, free from pest and disease injuries, bruises and blemishes were selected, washed with water and dried with soft cloth. Normally, for household purpose, the purchased tomato fruits are stored in open condition at room temperature or wrapped in polythene bags and stored in refrigerator. Hence, samples harvested at different maturity stages were stored at two experimental temperatures. One part (1kg) of tomatoes were placed as a single layer in a paper carton and kept open at ambient conditions prevailing in the

laboratory temperature ($28^{\circ}\text{C} \pm 1$) and the other part (1kg) was wrapped with polythene bag and stored in a refrigerator maintained at cold temperature, $6^{\circ}\text{C} \pm 1$ (Rai et al. 2012).

2.2. Measurement of nutrients

To study the effect of harvesting stage and storage temperature on nutrient contents, three fruits were taken randomly from each treatment on the day of harvest and subsequently on the 3rd, 6th, 9th and 12th days after harvest. Each fruit from a treatment represented a replicate. The different nutrients were assessed in each fruit.

Titration acidity, ascorbic acid and lycopene contents were estimated following the method of Ranganna (1977). To determine titration acidity, 10 g of the fruit sample was homogenised with about 25 ml of distilled water and filtered. The filtrate was collected and made up with distilled water. 10 ml of fruit extract was titrated against 0.1 N NaOH and the acidity was calculated and expressed as citric acid percentage.

Another 10 g of the fruit sample was blend with about 50 ml of 4% oxalic acid in a glass mortar and filtered. The filtrate was transferred to a 100 ml volumetric flask and made up with 4% oxalic acid. Known volume (10 ml) of this fruit extract was titrated against 2,4-dichlorophenol indophenol dye and the amount of ascorbic acid was calculated as mg (100g)^{-1} sample.

The lycopene content was measured in 5 g of fruit sample after extracting with acetone, till the pulp became colourless. Then, the acetone extract was mixed with 50 ml petroleum ether and 50 ml water in a separating funnel. Lycopene got transferred to the petroleum ether layer which was separated, dehydrated using anhydrous sodium sulphate and made up to 100 ml with petroleum ether. The absorbance of this solution was measured at 503 nm using a spectrophotometer.

Total sugars and phenols were extracted using ethanol. About 500 mg of the sample was homogenized in 5 ml of 80% ethanol using mortar and pestle. The homogenate was centrifuged and the supernatant was collected in a beaker. The residue was re-extracted with fresh 80% ethanol and centrifuged. The supernatant was pooled and evaporated to dryness. The residue was dissolved in distilled water (10 ml) and used for sugar and phenol estimation. Total soluble sugar was determined by Anthrone method (Hedge and Hofreiter 1962). The absorbance was read at 620 nm using spectrophotometer. Total sugar content was calculated using a standard graph and expressed as g (100g)^{-1} sample. Total phenol content was determined with the Folin-Ciocalteu reagent (Malik and

Singh 1980).The absorbance was measured at 650 nm using spectrophotometer. Total phenol content was calculated using a standard graph and expressed as **mg (100g)⁻¹** sample.

The experiment was conducted in a completely randomized design with three replications and the data were analysed by using three factorial ANOVA at 5% LSD. All statistical **analyses** were conducted using AGRESS software.

3. RESULTS AND DISCUSSION

Harvesting stage and storage temperature play important role in increasing the shelf-life and nutritional quality of fresh produce. The effect of temperature on shelf-life and nutritional quality of several tomato varieties have been studied earlier (Moneruzzaman et al. 2009; Nour et al. 2014). This experiment was conducted to study the effect of harvesting stage and storage temperature on nutritional quality of the most popular tomato variety PKM-1 cultivated in South India, especially in Tamil Nadu.

3.1. Titrable acidity

Tomato cv.PKM-1 fruits were harvested at three different ripening stages (mature-green, half ripe and red-ripe) to assess the nutritional quality at harvest and on storage. The study revealed that titrable acidity decreased gradually from mature green to red ripe stage (Table 1). Mature green, half ripe and red ripe tomatoes had 0.91, 0.84 and 0.77% titrable acidity respectively. Generally, the level of acidity in the fruits decreases during ripening process (Kandasamy 2016). The predominant acid in tomato is malic acid and is utilised as substrate for respiration during respiratory burst and further ripening, which is the reason for the decreasing trend in acidity during ripening (Helyes and Lugasi 2006). Storage study showed that titrable acidity decreased gradually on storage up to 12 days. This is in corroboration with the findings of Tigist et al. (2013). Least acidity of 0.51% was observed in tomato harvested at red ripe stage and stored at 28°C.

Table 1. Effect of harvesting stage and storage temperature on titrable acidity (%) of tomato fruits

Treatment	Days					Mean
	0	3	6	9	12	
M1T1	0.91	0.81	0.72	0.59	0.58	0.72

M1T2	0.91	0.85	0.75	0.67	0.65	0.77
M2T1	0.84	0.74	0.6	0.57	0.53	0.66
M2T2	0.84	0.8	0.7	0.67	0.62	0.73
M3T1	0.77	0.72	0.61	0.54	0.51	0.63
M3T2	0.77	0.74	0.71	0.63	0.59	0.69
MEAN	0.84	0.78	0.68	0.61	0.58	0.7

Factor	M	T	D	MT	TD	MD	MTD
SED	0.003	0.003	0.005	0.005	0.006	0.008	0.011
CD (0.05)	0.007	0.006	0.009	0.010	0.013	0.0156	0.022

145 M- Maturity stage, M1- Mature green, M2- Yellow green, M3- Red ripe.

146 T- Storage temperature, T1- 28°C, T2- 6°C.

147 D- No. of days of storage.

148 SED-Standard Error of Difference between two means, CD- Critical Difference

149 3.2. Ascorbic acid

150 Ascorbic acid content significantly increased with advancing in harvest maturity (Table 2).
151 Maximum ascorbic acid [21.54 mg (100g)⁻¹] was found in red ripe and the least in mature green [10.2
152 mg (100g)⁻¹] fruits at harvest. During storage, all treatments showed drastic decline in ascorbic acid
153 content both at room temperature and in refrigerator. Highest [14.64 mg (100g)⁻¹] ascorbic acid
154 content at twelfth day of storage was observed in red ripe fruits stored at 6°C and lowest [6.05 mg
155 (100g)⁻¹] in mature green fruits stored at room temperature. Ascorbic acid content was comparatively
156 higher when stored in refrigerator than in room temperature. Similar trend was observed in cherry
157 tomato also (Tsaniklidis et al. 2014). In tomato fruit, ascorbic acid content increases with maturity and
158 stage of ripening, however once fruit reach the full ripe stage, ascorbic acid content starts to decline
159 (Mathooko 2003). An increase in ascorbic acid content in fruit is thought to be an indication that the
160 fruit is still in the ripening stage, while a decrease indicates a senescent fruit (Sammi and Masud
161 2007).

Table 2. Effect of harvesting stage and storage temperature on ascorbic acid content [mg (100g)⁻¹] in tomato fruits

Treatment	Days					Mean	
	0	3	6	9	12		
M1T1	10.2	9.07	8.17	6.57	6.05	8.01	
M1T2	10.2	9.67	8.76	7.57	7.08	8.66	
M2T1	15.43	13.91	12.48	9.95	9.32	12.22	
M2T2	15.43	14.52	13.51	11.65	10.67	13.16	
M3T1	21.54	19.76	17.57	14.32	12.65	17.17	
M3T2	21.54	19.57	18.25	16.24	14.64	18.05	
MEAN	15.72	14.42	13.12	11.055	10.07	12.88	
Factor	M	T	D	MT	TD	MD	MTD
SED	0.122	0.100	0.158	0.173	0.223	0.273	0.386
CD (0.05)	0.244	0.199	0.315	NS	0.446	0.546	NS

M- Maturity stage, M1- Mature green, M2- Yellow green, M3- Red ripe.

T- Storage temperature, T1- 28°C, T2- 6°C.

D- No. of days of storage.

SED-Standard Error of Difference between two means, CD- Critical Difference

3.3. Total soluble sugars

Total sugar content of tomato fruits varied significantly at different ripening stages. It was found that total sugar content was increased with the advancement of ripening of fruits irrespective of maturity condition and storage temperature (Table 3). The highest quantity of total sugar [5.45 g (100g)⁻¹] was recorded in red ripe tomatoes stored at room temperature; while it was lowest [3.84 g (100g)⁻¹] in mature green tomatoes stored at 6°C at twelfth day of storage. The gradual increase in

total sugar content found in this experiment is agreement with previous studies with tomato and mango (Sood et al. 2011; Islam et al. 2013).

Table 3. Effect of harvesting stage and storage temperature on total sugar content [g (100g)⁻¹] in tomato fruits

Treatment	Days					Mean	
	0	3	6	9	12		
M1T1	3.11	3.52	4.24	4.52	4.63	4.00	
M1T2	3.11	3.19	3.49	3.76	3.84	3.48	
M2T1	3.59	4.06	4.53	4.72	4.96	4.37	
M2T2	3.59	3.76	3.92	4.06	4.16	3.90	
M3T1	4.08	4.52	5.16	5.31	5.45	4.90	
M3T2	4.08	4.13	4.23	4.30	4.37	4.22	
MEAN	3.59	3.86	4.26	4.45	4.57	4.15	
Factor	M	T	D	MT	TD	MD	MTD
SED	0.032	0.026	0.041	0.045	0.058	0.071	0.100
CD (0.05)	0.063	0.052	0.082	0.090	0.115	0.141	NS

M- Maturity stage, M1- Mature green, M2- Yellow green, M3- Red ripe.

T- Storage temperature, T1- 28°C, T2- 6°C.

D- No. of days of storage.

SED-Standard Error of Difference between two means, CD- Critical Difference

3.4. Lycopene

Lycopene is the most important antioxidant compound of the ripe tomato fruit and the one which determines its red colour. Lycopene content (Table 4) rapidly increased from half ripened stage [0.66 mg (100g)⁻¹] to red ripe stage [2.04 mg (100g)⁻¹]. Lycopene also increased during storage. Increased levels of lycopene in tomato during storage is due to ripening advancement of tomato fruits

and conversion of chloroplasts to chromoplasts. Ghosh et al. (2014) also reported the increase in lycopene contents during storage. The increasing in redness of tomatoes during ripening is due to lycopene accumulation (Rai et al. 2012). Red ripe fruits stored for 12 days at 28°C had the highest lycopene [2.98 mg (100g)⁻¹]. Fruits stored at 6°C showed lesser accumulation of lycopene compared to those stored at 28°C. Previous studies on storage at low temperature also reported lesser accumulation of lycopene in tomato (Aljouni et al. 2001).

Table 4. Effect of harvesting stage and storage temperature on lycopene content [mg (100g)⁻¹ in tomato fruits

Treatment	Days					Mean
	0	3	6	9	12	
M1T1	0.35	0.41	0.93	1.32	1.34	0.87
M1T2	0.35	0.37	0.43	0.57	0.68	0.48
M2T1	0.66	0.73	1.67	1.84	2.11	1.4
M2T2	0.66	0.67	0.96	1.12	1.18	0.92
M3T1	2.04	2.13	2.24	2.62	2.98	2.4
M3T2	2.04	2.09	2.12	2.18	2.26	2.14
MEAN	1.02	1.07	1.39	1.61	1.76	1.37

Factor	M	T	D	MT	TD	MD	MTD
SED	0.010	0.008	0.013	0.014	0.018	0.022	0.031
CD (0.05)	0.020	0.016	0.026	0.282	0.036	0.045	0.063

M- Maturity stage, M1- Mature green, M2- Yellow green, M3- Red ripe.

T- Storage temperature, T1- 28°C, T2- 6°C.

D- No. of days of storage.

SED-Standard Error of Difference between two means, CD- Critical Difference

3.5. Total phenols

200 Total phenolic content gradually decreased with enhancement of ripening. Mature green and
 201 red ripe tomato had 43.77 and 28.2 **mg (100g)⁻¹** of total phenols (Table 5). A decrease in later stages
 202 of ripening was reported earlier (Raffo et al. 2002). Total phenols decreased significantly during
 203 storage of tomato fruits at room temperature and in refrigerator. Total phenol content of mature green
 204 tomato decreased from 43.77 **mg (100g)⁻¹** to 17.3 and 30.13 **mg (100g)⁻¹** when stored at 28 and 6°C
 205 respectively. In red ripe fruits, total phenol decreased from 28.2 **mg (100g)⁻¹** to 16.47 and 24.33 **mg**
 206 **(100g)⁻¹** when stored at 28 and 6°C respectively. The decrease in phenols might be due to
 207 senescence and breakdown of cell structure during storage, as was reported by Ali et al. (2010).

208 **Table 5. Effect of harvesting stage and storage temperature on total phenol content**
 209 **[mg (100g)⁻¹] in tomato fruits**

Treatment	Days					Mean	
	0	3	6	9	12		
M1T1	43.77	40.5	30.63	21.37	17.3	30.71	
M1T2	43.77	48.43	40.6	35.6	30.13	39.71	
M2T1	32.1	28.2	21.93	18.37	17.07	23.53	
M2T2	32.1	30.23	28.13	26.6	25.03	28.42	
M3T1	28.2	23.63	18.57	17.83	16.47	20.94	
M3T2	28.2	27.83	26.53	25.23	24.33	26.42	
MEAN	34.69	33.14	27.73	24.17	21.72	28.29	
Factor	M	T	D	MT	TD	MD	MTD
SED	0.376	0.307	0.486	0.532	0.687	0.842	1.190
CD (0.05)	0.753	0.615	0.972	1.065	1.375	1.684	NS

210 M- Maturity stage, M1- Mature green, M2- Yellow green, M3- Red ripe.

211 T- Storage temperature, T1- 28°C, T2- 6°C.

212 D- No. of days of storage.

213 SED- Standard Error of Difference between two means, CD- Critical Difference

214 **4. CONCLUSION**

215 The experiments conducted have shown that the nutritional composition including total
216 sugars, ascorbic acid, lycopene and phenolics of the tomato fruit is strongly influenced during different
217 ripening stages. Ascorbic acid, sugars and lycopene contents showed their highest levels at the red
218 ripe stage while titrable acidity and phenolics were highest at mature green stage. Hence, tomato fruit
219 harvested at red ripe stage had highest nutritional quality. Low temperature is preferred for storing
220 tomato fruits since, storage at 6 °C caused less change in nutritional components.

221 **LITERATURE CITED**

- 222 Agarwal S, Rao AV. Tomato lycopene and its role in human health and chronic diseases. Canadian
223 Medical Association Journal. 2000;163(6):739-44.
- 224 Ali A, Maqbool M, Ramachandran S, Alderson PG. Gum arabic as a novel edible coating for
225 enhancing shelf-life and improving postharvest quality of tomato (*Solanum lycopersicum* L.)
226 fruit. Postharvest Biology and Technology. 2010;58(1):42-7.
- 227 Aljouni S, Kremer S, Masih L. Lycopene content in hydroponic and non-hydroponic tomatoes during
228 post harvest storage. Food Australia. 2001;53(5):195-6.
- 229 Anthon GE, LeStrange M, Barrett DM. Changes in pH, acids, sugars and other quality parameters
230 during extended vine holding of ripe processing tomatoes. Journal of the Science of Food and
231 Agriculture. 2011 May 1;91(7):1175-81.
- 232 Ghosh A, Dey K, Bhowmick N, Medda PS and Dutta P. Effect of guar gum as an edible coating to
233 improve shelf life and quality of tomato (*Solanum lycopersicum* L.) fruits during storage. The
234 Ecoscan. 2014; special issue (VI): 201-207.
- 235 Goren A, Alkalai-Tuvia S, Perzelan Y, Aharon Z, Ilic Z, Fallik E. Harvested tomato quality and
236 nutritional levels as affected by high temperatures in Mediterranean wholesale markets, and
237 home or refrigerated temperatures. Advances in Horticultural Science. 2010 Jan 1:200-6.
- 238 Grierson D, Kader AA. Fruit ripening and quality. In: Atherton J, Rudich J, editors. The tomato crop: A
239 scientific basis for improvement. Chapman and Hall. London; 1986.
- 240 Hedge JE, Hofreiter BT. Methods of estimating starch and carbohydrates. In: Whistler RL, Miller JNB.
241 Carbohydrate Chemistry. Academic Press, New York; 1962.

242 Helyes L, Lugasi A. Formation of certain compounds having technological and nutritional importance
 243 in tomato fruits during maturation. *Acta Alimentaria*. 2006;35(2):183-93.

244 Islam MK, Khan MZ, Sarkar MA, Absar N, Sarkar SK. Changes in acidity, TSS, and sugar content at
 245 different storage periods of the postharvest mango (*Mangifera indica* L.) Influenced by
 246 bavistin. *International Journal of Food Science*. 2013;Dec 29.

247 Kader AA. Fruit maturity ripening and quality relationship. In: Management of fruit ripening, post-
 248 harvest horticultural series 8, Department of Pomology, University of California, Davis;1994.

249 Kandasamy P. Quality of tomato stored under controlled atmosphere condition build by diffusion
 250 channel system. *International Journal of Food and Nutritional Science*. 2016; 5(4).

251 Kaur C, Kapoor HC. Antioxidants in fruits and vegetables—the millennium's health. *International*
 252 *Journal of Food Science & Technology*. 2001;36(7):703-25.

253 Malik CP, Singh M.B. Plant enzymology and Histoenzymology. Kalyani publishers, New Delhi;1980.

254 Mathooko FM. A comparative study of the response of tomato fruit to low temperature storage and
 255 modified atmosphere packaging. *African Journal of Food, Agriculture, Nutrition and*
 256 *Development*. 2003;2:34-41.

257 Missio JC, Renau RM, Artigas FC, Cornejo JC. Sugar-and-acid profile of Penjar tomatoes and its
 258 evolution during storage. *Scientia Agricola*. 2015;72(4):314-21.

259 Mohammed M, Wilson LA, Gomes PL. Postharvest sensory and physiochemical attributes of
 260 processing and non-processing tomato cultivar. *Journal of Food Quality*. 1999;22:167–182.

261 Moneruzzaman KM, Hossain AB, Sani W, Saifuddin M, Alenazi M. Effect of harvesting and storage
 262 conditions on the post harvest quality of tomato (*Lycopersicon esculentum* Mill) cv. Roma VF.
 263 *Australian Journal of Crop Science*. 2009;3(2):113.

264 Nour V, Trandafir I, Ionica ME. Evolution of antioxidant activity and bioactive compounds in tomato
 265 (*Lycopersicon esculentum* Mill.) fruits during growth and ripening. *Journal of Applied Botany*
 266 *and Food Quality*. 2014;23;87.

267 Parashar S, Sharma H, Garg M. Antimicrobial and antioxidant activities of fruits and vegetable peels:
 268 a review. *Journal of Pharmacognosy and Phytochemistry*. 2014;3(1):160-164.

269 Pedapati A, Reddy, RVSK, Babu JD, Kumar SS, Sunil N. Combining ability analysis for yield and
 270 physiological drought related traits in tomato (*Solanum lycopersicum* L.) under moistures
 271 stress. *The Bioscan*. 2013;8: 1537-1544.

272 Raffo A, Leonardi C, Fogliano V, Ambrosino P, Salucci M, Gennaro L, Bugianesi R, Giuffrida F,
 273 Quaglia G. Nutritional value of cherry tomatoes (*Lycopersicon esculentum* cv. Naomi F1)
 274 harvested at different ripening stages. Journal of Agricultural and Food Chemistry.
 275 2002;50(22):6550-6.

276 Rai GK, Kumar RA, Singh AK, Rai PK, Rai M, Chaturvedi AK, Rai AB. Changes in antioxidant and
 277 phytochemical properties of tomato (*Lycopersicon esculentum* Mill.) under ambient condition.
 278 Pak. J. Bot. 2012;44(2):667-70.

279 Rangana S. Manual of analysis of fruits and vegetable products, Tata McGraw Hill, New Delhi,
 280 India:1977.

281 Reid MS. Product maturity and maturity indices. In: Kader AA, Kasmire RF, Mitchel FG, Reid MS,
 282 Sommer NF, Thompson JF. editors. Post harvest technology of horticulture crops. Agricultural
 283 and Natural Resources. USA: Publication Division of Agriculture and Natural Resources,
 284 Univ. of California, Oakland. USA: 1985.

285 Sammi S, Masud T. Effect of different packaging systems on storage life and quality of tomato
 286 (*Lycopersicon esculentum* var. Rio Grande) during different ripening stages. Internet Journal
 287 of Food Safety. 2007;9:37-44.

288 Sood M, Kaul RK, Singh A. Effect of post harvest treatments on changes in sugar and lycopene
 289 content of tomato (*Lycopersicon esculentum*). World Journal of Agricultural Sciences.
 290 2011;7(5):613-6.

291 Tigist M, Workneh TS, Woldetsadik K. Effects of variety on the quality of tomato stored under ambient
 292 conditions. Journal of Food Science and Technology, 2013;50(3):477–486.

293 Tsaniklidis G, Delis C, Nikoloudakis N, Katinakis P, Aivalakis G. Low temperature storage affects the
 294 ascorbic acid metabolism of cherry tomato fruits. Plant Physiology and Biochemistry.
 295 2014;84:149-57.

296 Willis MS, Wians FH. The role of nutrition in preventing prostate cancer: a review of the proposed
 297 mechanism of action of various dietary substances. Clinica Chimica Acta. 2003;330(1):57-83.

298 Yanuriati A, Savage GP, Rowe RN. The effects of ethanol treatment on the metabolism, shelf life and
 299 quality of stored tomatoes at different maturities and temperatures. Journal of the Science of
 300 Food and Agriculture. 1999;79(7):995-1002.