

Original Research Article

Cardioprotective effect of enalapril on isoproterenol-induced myocardial infarction in rats

Abstract

Myocardial infarction is the irreversible death of heart muscle secondary to prolonged lack of oxygen supply. The present study was designed to evaluate the protective effect of enalapril in isoproterenol-induced myocardial infarction in rats using changes in haemodynamic, biochemical, histopathological and immunohistochemistry parameters. Twenty one male Wistar rats divided into three groups were used where the control (group A) were administered normal saline for 7 days, group B animals received normal saline for 7 days and thereafter isoproterenol (ISO) at 85 mg/kg on day 8 and 9. Group C animals were pretreated with enalapril (10mg/kg) for 7 days and thereafter received ISO on day 8 and 9. On day 10, the blood pressure changes of the animals were measured and thereafter sacrificed by cervical dislocation. The heart of each rat was removed, homogenized and used to assay for some oxidative stress markers and some antioxidant parameters. In this study, ISO caused myocardial infarction as seen by significant decrease in systolic, diastolic and mean arterial pressure but were corrected by enalapril. Enalapril caused significant increase in the levels of SOD, GPx, GST and GSH but significant decrease in MDA content and H₂O₂ generation. But reverse was the case for group B animals. Immunohistochemistry showed that ISO caused higher expressions of cardiac C-reactive protein (CRP) and cardiac troponins 1 (CTn1) and decrease in IL-10 β but vice-versa for enalapril. No histopathological changes were recorded for enalapril. The study thus showed that enalapril significantly exhibit cardioprotective effects.

Key words: Enalapril, myocardial infarction, cardioprotection, immunohistochemistry, antioxidant

Introduction

Human health is being seriously threatened by cardiovascular diseases (CVD) which is regarded as the main cause of death throughout the world [1, 2]. Both the underdeveloped and the developed countries have not been able to control this disease. At present CVD is the highest killer disease in US [3]. Myocardial infarction (MI) is a common presentation of ischemic heart

disease (IHD) and remains the major cause of death in the developed world. Though rapid advancements have been made in the treatment of coronary artery diseases (CAD), MI is still a major pathological issue worldwide [4]. Increased myocardial metabolic demand and decreased supply of oxygen as well as nutrients via the coronary circulation to the myocardium bring about myocardial infarction hence leading to cell injury. This pathological heart condition is one of the most lethal manifestations of cardiovascular diseases [5, 6]. Acute myocardial infarction or heart attack occurs when blood stops flowing to part of the heart leading to injury to the heart muscle due to the fact the heart is not receiving enough oxygen. The reason for this lack of oxygen supply is usually because one of the coronary arteries that supplies blood to the heart develops a blockage as a result of an unstable build up of white blood cells, cholesterol and fat. Fatty acid is the major source of fuel for energy, though glucose could also be used [7]. However in an ischemic heart as a result of less availability of oxygen, glucose becomes the major source of energy, therefore glycolysis switches from aerobic to anaerobic conditions. There is therefore a resultant shifting of metabolic utilization of substrates toward glucose from fatty acids [8]. The normal heart utilizes fatty acids because this provides the highest energy yield per molecule of substrate metabolized but glucose becomes an important preferential substrate for metabolism and ATP generation under specific pathological conditions because it can provide greater efficiency in producing high energy products per oxygen consumed compared to fatty acids [9]. Isoproterenol [1-(3, 4-dihydroxyphenyl)-2-isopropylaminoethanol hydrochloride] (ISO) a synthetic catecholamine is a β -adrenergic agonist that is very important in the regulation of myocardial contractility and metabolism. It serves as a standard model for the study of potentially beneficial effects of numerous drugs on cardiac function [10, 11]. ISO induces myocardial injury in rat because of the alteration in physiological balance between production of free radicals and antioxidative defence system [12]. It thus causes acute condition of myocardial necrosis which can lead to cardiac dysfunctions, increased lipid peroxidation, altered activities of cardiac enzymes and antioxidants [13]. It has been observed that the pathophysiological and morphological changes observed in ISO-treated rats is similar to those observed in human MI [14].

Enalapril an Angiotensin-converting-enzyme inhibitor (ACE inhibitor) is a drug used primarily for the treatment of high blood pressure and congestive heart failure where it can be used alone

or in combination with other antihypertensive agents. ACE inhibitors have also been found to be useful for other cardiovascular and kidney diseases including acute myocardial infarction, diabetic nephropathy, and cardiac failure [15]. The mechanism of action of ACE inhibitors involves reduction of the activity of the renin-angiotensin-aldosterone system (RAAS) [16].

In recent times, a novel strategy has been employed in drug discovery. It is the use of known and approved drugs and compounds for newer indications. This is termed drug repurposing. In this study, Isoproterenol was used to induce acute myocardial infarction and pharmacological activities of enalapril were then explored with the view of understanding some of its cardioprotective activities in rats and then to see if it could serve as a repurposed drug for myocardial infarction.

Materials and Methods

Chemicals and reagents

Isoproterenol, enalapril, Tween 80, Biurett's reagent, hydrogen peroxide, hydrochloric acid, sulphuric acid, xylene orange, potassium dichromate, O-diasinidine, sodium potassium tartrate, copper sulphate, ethanol, sodium azide, 2-dichloro-4-nitrobenzene (CDNB) Greiss reagent, phosphoric acid, sodium hydroxide, N 1-naphthyl ethylenediamine, sulphanilamide, distilled water, , phosphate buffer saline, creatinine reagent, copper sulphate, tri chloro acetate, reduced glutathione (GSH), thiobarbituric Acid (TBA), trichloroacetic acid (TCA), ammonium ferrous sulphate, glacial acetic acid, potassium iodide, sorbitol, Ellman's reagent (DTNB), ethanol, urea reagent. All other chemicals used were of analytical grade and obtained from British Drug Houses (Poole, Dorset, UK). All other chemicals, reagents and drugs used were of analytical grade.

Experimental animals

All experiments and protocols described in present study were approved by the UI-ACUREC. Twenty one (21) male Wistar rats weighing 90-160g were obtained from the Experimental animal unit of the Faculty of Veterinary Medicine, University of Ibadan for the experiment. They were allowed free access to standard rat pellets and fresh water *ad libitum*. The rats were housed in the animal house unit of the Department of Veterinary Physiology, Biochemistry and

Pharmacology, University of Ibadan with a 12 hour light duration. Pre-conditioning of the rats was done for two weeks before commencement of the experiment.

Cardioprotective study

The animals were randomly divided into three (3) groups with seven (7) animals in each group, and the treatment was as follow: Animals in the control (group A) were administered normal saline, group B; isoproterenol at 85mg/kg, while group C animals were pretreated with enalapril 10mg/kg for 7 days and thereafter administered ISO (85mg/kg) day 8 and 9. Normal saline was used as a vehicle. Blood pressure values of all the animals were carried out on day 10. At the end of the experimental period, blood samples were collected for haematology and serum chemistry before the rats were sacrificed by cervical dislocation. The serum in plain bottles was rapidly centrifuged at 4000 revolutions per minute (rpm) for fifteen (15) minutes and processed for determination of serum myeloperoxidase, total protein, and xanthine oxidase, AST, ALT and nitric oxide. The heart of each rat was carefully removed and homogenized on ice and then used to assay for some oxidative stress markers and antioxidant parameters. Baseline cardiovascular parameters were collected prior to the commencement of the experiment. The equipment used was a non-invasive tail cuff BP monitor, the 6-channel CODA blood pressure monitor for rats and mice. Blood pressure parameters including the systolic, diastolic, and mean arterial blood pressure parameters were determined indirectly in nonanaesthetised rats, by tail plethysmography with the use of an electrospphygmanometer (CODA, Kent Scientific, USA). The average of at least nine most consistent readings, taken in the quiescent state, following acclimatization, was recorded per animal.

Preparation of tissue homogenate

The heart tissues of the rats were harvested on ice, rinsed with normal saline and homogenized in aqueous potassium buffer (0.1 M, pH 7.4) and the homogenate centrifuged at 12,000 rpm (4°C) for 15 min to obtain the supernatant fraction.

118 *Determination of Biochemical assay*

119 Biuret method as described by Gornal et al [17] was used to determine the protein concentrations
 120 of the various samples with a slight modification. To prevent precipitation of Cu^{2+} ions as
 121 cuprous oxide potassium iodide was added to the reagent. To determine the concentration of
 122 reduced glutathione the method of Beutler et al [18] was used while glutathione peroxidase
 123 (GPX) activity was measured by the method of Rotruck et al. [19] In this case, hydrogen
 124 peroxide was used as substrate to oxidize reduced glutathione to oxidized glutathione (GSSG).
 125 Estimation of Glutathione S-transferase (GST) was by the method of Habig et al [20] using 1-
 126 chloro-2, 4-dinitrobenzene as substrate. Superoxide dismutase (SOD) assay on the other hand
 127 was carried out by the method of Misra and Fridovich [21]. MDA content was measured in the
 128 heart as an index of lipid peroxidation [22]. Hydrogen peroxide generation was measured using
 129 Wolff's [23] method while the determination of Sulfhydryl (Thiol) content was by-the method of
 130 Ellman [24]. Nitric oxide was quantified as previously described [25].

131 *Histopathology*

132 Small slices of the heart were collected in 10% buffered formalin for proper fixation and after the
 133 tissues have been processed and embedded in paraffin wax, sections that were about 5-6 μm
 134 thick were made and stained with haematoxylin and eosin for histopathological examination
 135 [26].

136 *Immunohistochemistry of Cardiac troponins-1, CRP and IL-10*

137 The heart tissues obtained from buffered formalin perfused rats were paraffin embedded and then
 138 used for immunohistochemistry. Paraffin sections were melted at 60 °C in the oven but the
 139 dewaxing of the samples in xylene was followed by passage through ethanol of decreasing
 140 concentration (100-80%). Peroxidase quenching in 3% H_2O_2 /methanol was carried out with
 141 subsequent antigen retrieval performed by microwave heating in 0.01 M citrate buffer (pH 6.0)
 142 to boil. All the sections were blocked in normal goat serum (10%, HistoMark[®], KPL,
 143 Gaithersburg MD, USA) and probed with cardiac troponins 1, CRP antibody and IL-10
 144 (Abclonal[®]), 1:375 for 16 h in a refrigerator. Detection of bound antibody was carried out using
 145 biotinylated (goat anti-rabbit, 2.0 $\mu\text{g}/\text{ml}$) secondary antibody and subsequently, streptavidin

peroxidase (Horse Radish Peroxidase- streptavidin) according to manufacturer's protocol (HistoMark[®], KPL, Gaithersburg MD, USA).

Diaminobenzidine (DAB, Amresco[®], USA) was used to enhance the reaction product for 6 – 10 min and counterstained with high definition haematoxylin (Enzo[®], NY - USA), and was thereafter dehydrated in ethanol. Once the slides were covered with cover slips, they were sealed with resinous solution. The immunoreactive positive expression of CRP, Bcl-2 and NFkB intensive regions were viewed starting from low magnification on each slice then with 400 × magnifications using a photo microscope (Olympus) and a digital camera (Toupcam[®], Touptek Photonics, Zhejiang, China).

Statistical analysis

All values were expressed as mean ± standard deviation (SD). The test of significance between two groups was estimated by Student's t-test. One-way Analysis of Variance (ANOVA) with Tukey's post-hoc test using Graph pad prism 5.0 was also performed with p-values < 0.05 considered statistically significant.

Results

In this study, ISO caused significant decreases in the levels of SBP, DBP and MAP while enalapril (ENA) caused significant increase though not to the same extent as the control (Figures 1-3). The results of haematological analysis showed that ISO caused significant increases in the levels of WBC, PCV, MCV and MCH while ENA caused significant decrease in WBC and no changes relative to ISO (Table 1). ISO also caused significant increases in the levels of AST and ALT while ENA caused significant decreases in the levels of these enzymes. On the other hand, while ISO caused significant decrease in the level of NO, ENA caused significant increase (Table 2). ISO caused significant increases in the levels of oxidative markers such as MDA, H₂O₂ and MPO while ENA caused significant decreases in the levels of these markers in a similar fashion to the control (Figures 4-6). On the other hand while ISO caused significant decrease in the levels of protein thiols and non protein thiols, ENA caused a significant increase in the levels of these molecules (Figures 7 and 8). The result also showed that ISO caused significant decrease in the levels of anti-oxidant markers such as SOD, GPx, GST and GSH but

reverse is the case for ENA (Figures 9-12). Histopathological examinations showed that while there is severe infiltration of inflammatory cells into the cardiac tissue, there was no visible lesion seen in the ENA and control groups (Figure 13). The immunohistochemical analysis showed that there was high expressions of cardiac troponin and and CRP in ISO group but lower expression of these proteins in ENA and control group (Figures 14 and 15). In the case of IL-10, there was low expression of this protein in ISO group but higher expression in ENA and control group (Figure 16).




Figure 1: Effect of enalapril on SBP in isoproterenol induced myocardial infarction using rats as a model. The superscript a showed that ISO caused significant decrease when compared to control while superscript b showed significant decrease when compared with ENA.




Figure 2: Effect of enalapril on DBP in isoproterenol induced myocardial infarction using rats as a model. The superscript a showed that ISO caused significant decrease in the level of this parameter compared to control while b showed that ENA caused significant increase relative to control and ISO groups.




Figure 3: Effect of enalapril MAP in isoproterenol induced myocardial infarction using rats as a model. The superscripts showed that ISO caused significant decrease relative to ENA and control groups.

Table 1: Effects of enalapril on RBC, WBC, HB, PCV, MCV, MCH and MCHC in isoproterenol induced myocardial infarction using rats as a model (n = 7)

Parameters	Control	ISO	Enalapril
RBC ($\times 10^{12}/L$)	4.75 $\pm$ 0.90	4.96 $\pm$ 0.43	5.03 $\pm$ 0.69
WBC ($10^3/\mu L$)	5.47 $\pm$ 0.38	6.71 $\pm$ 1.13 ^a	4.68 $\pm$ 1.68 ^b
HB (g/dl)	13.33 $\pm$ 1.40	15.15 $\pm$ 1.84	14.95 $\pm$ 1.62
PCV (%)	45.75 $\pm$ 4.65	54.25 $\pm$ 4.25 ^a	50.25 $\pm$ 3.10
MCV (fl)	83.88 $\pm$ 9.03	127.33 $\pm$ 30.12 ^a	98.87 $\pm$ 22.76
MCH (pg)	26.41 $\pm$ 3.48	38.64 $\pm$ 8.08 ^a	26.05 $\pm$ 2.25
MCHC (%)	29.97 $\pm$ 2.05	27.41 $\pm$ 2.38	30.79 $\pm$ 2.37

Values are mean $\pm$ SD, n =5, ^a - $\alpha < 0.05$ compared with control, ^b - $\alpha < 0.05$ compared with ISO.

The superscript (a) showed that ISO caused significant decrease in the level of this parameter compared to control while (b) showed that ENA caused significant increase relative to control and ISO groups.

Table 2: Effects of enalapril on ALT, AST and NO in isoproterenol induced myocardial infarction using rats as a model

Parameters	Control	ISO	Enalapril
ALT	14.51±0.02	14.67±0.05 ^a	14.41±0.05 ^{ab}
AST	19.91±0.01	19.97±0.02 ^a	19.87±0.02 ^{ab}
NO	4.11±0.68	1.72±0.47 ^a	2.67±0.71 ^{ab}

Values are mean ± SD, n =5, ^a - $\alpha < 0.05$ compared with control, ^{ab} - $\alpha < 0.05$ compared with ISO.

The superscript (a) showed that ISO caused significant decrease in the level of this parameter compared to control while (b) showed that ENA caused significant increase relative to control and ISO groups.




Figure 4: Effect of enalapril on lipid peroxidation in isoproterenol induced myocardial infarction using rats as a model. (n=5). Values are presented as mean $\pm$ standard deviation. Grp A (Control), Grp B (ISO treated only), Grp C (ENA+ 85 mg/kg ISO). The superscript (a) showed that ISO caused significant decrease in the level of this parameter compared to control while (b) showed that ENA caused significant increase relative to control and ISO groups.




Figure 5: Effect of enalapril on hydrogen peroxide generation in isoproterenol-induced myocardial infarction using rats as a model (n=5). Values are presented as mean $\pm$ standard deviation. Superscripts (a) indicates significant difference ($p < .05$) when compared with control (Grp A), whereas superscripts (b) indicates significant difference ($p < .05$) when compared with ISO treated only (Grp B). Grp A (Control), Grp B (ISO treated only), Grp C (ENA+ 85 mg/kg ISO).




Figure 6: Effect of enalapril on myeloperoxidase in isoproterenol induced myocardial infarction using rats as a model (n=5). The superscript “a” showed that ISO caused significant increase in the level of this parameter when compared to the control and ENA groups.




Figure 7: Effect of enalapril on protein thiol in isoproterenol induced myocardial infarction using rats as a model (n=5). Values are presented as mean $\pm$ standard deviation. Superscripts (a) indicates significant difference ($p < .05$) when compared with control (Grp A), whereas superscripts (b) indicates significant difference ($p < .05$) when compared with ISO treated only (Grp B). Grp A (Control), Grp B (ISO treated only), Grp C (ENA+ 85 mg/kg ISO), Grp D (ISO+100 mg/kg AP), Grp E (ISO+200 mg/kg AP) & Grp F (ISO+400 mg/kg AP).




Figure 8: Effect of enalapril on non protein thiol in isoproterenol induced myocardial infarction using rats as a model (n=5). Values are presented as mean $\pm$ standard deviation. Superscripts (a) indicates significant difference ($p < .05$) when compared with control (Grp A), whereas superscripts (b) indicates significant difference ($p < .05$) when compared with ISO treated only (Grp B). Grp A (Control), Grp B (ISO treated only), Grp C (ENA+ 85 mg/kg ISO).




Figure 9: Effect of enalapril on superoxide dismutase enzyme in isoproterenol induced myocardial infarction (n=5). Values are presented as mean $\pm$ standard deviation. Superscripts (a) indicates significant difference ($p < .05$) when compared with control (Grp A), whereas superscripts (b) indicates significant difference ($p < .05$) when compared with ISO treated only (Grp B). Grp A (Control), Grp B (ISO treated only), Grp C (ENA+ 85 mg/kg ISO).




Figure 10: Effect of enalapril on glutathione peroxidase enzyme in isoproterenol induced myocardial infarction using rats as a model (n=5). Values are presented as mean $\pm$ standard deviation. Superscripts (a) indicates significant difference ($p < .05$) when compared with control (Grp A), whereas superscripts (b) indicates significant difference ($p < .05$) when compared with ISO treated only (Grp B). Grp A (Control), Grp B (ISO treated only), Grp C (ENA+ 85 mg/kg ISO).




Figure 11: Effect of enalapril on glutathione-s- transferase enzyme in isoproterenol induced myocardial infarction using rats as a model (n=5). Values are presented as mean $\pm$ standard deviation. Superscripts (a) indicates significant difference ($p < .05$) when compared with control (Grp A), whereas superscripts (b) indicates significant difference ($p < .05$) when compared with ISO treated only (Grp B). Grp A (Control), Grp B (ISO treated only), Grp C (ENA+ 85 mg/kg ISO).





Figure 12: Effect of enalapril on reduced glutathione in isoproterenol induced myocardial infarction using rats as a model (n=5). Values are presented as mean $\pm$ standard deviation. Superscripts (a) indicates significant difference ($p < .05$) when compared with control (Grp A), whereas superscripts (b) indicates significant difference ($p < .05$) when compared with ISO treated only (Grp B). Grp A (Control), Grp B (ISO treated only), Grp C (ENA+ 85 mg/kg ISO).

378

379



380

381

382 **Figure 13:** The photomicrograph of heart from isoproterenol induced myocardial infarction
 383 using rats as a model. A (Control) shows no visible lesion. B (ISO): shows severe infiltration of
 384 inflammatory cells. C (enalapril) shows no visible lesion. Mag. ×400

385

386

387

388




Figure 14: Immunohistochemistry of cardiac troponin in heart of isoproterenol induced myocardial infarction rats. A (Control): show positive and low expression of CTnI, B (ISO): shows higher expression of CTnI than control, C (enalapril) shows lower expression of CTnI than B (ISO). The slides were counterstained with high definition haematoxylin. Mag. x100




Figure 15: Immunohistochemistry of c- reacting protein in heart of isoproterenol induced myocardial infarction rats. A (Control): show positive and low expression of CRP, B (ISO): shows higher expression of CRP than control, C (enalapril) shows lower expression of CRP than B (ISO). The slides were counterstained with high definition haematoxylin. Mag. x100




Figure 16: Immunohistochemistry of interleukin-10 in heart of isoproterenol induced myocardial infarction rats. A (Control): show positive and higher expression of IL-10, B (ISO): shows lower expression of IL-10 than control, C (enalapril) shows higher expression of IL-10 than B (ISO). The slides were counterstained with high definition haematoxylin. Mag. x100

Discussion

Myocardial infarction (MI), one of the main causes of death from cardiovascular disease is defined as an acute condition of necrosis of the myocardium and it occurs as a result of imbalance between coronary blood supply and myocardial demand [27]. MI is known to cause local inflammation and apoptosis and this can result in cardiomyocyte damage [28]. ISO induces cardiac necrosis by several mechanisms, including increased oxygen consumption, poor oxygen utilization, increased calcium overload and accumulation, altered myocardial cell metabolism, increased myocardial cAMP levels, deranged electrolyte milieu, altered membrane permeability, intracellular acidosis, and increased levels of lipid peroxides¹¹. The pathophysiological changes that occurred in heart following isoproterenol administration in rats are comparable to those taking place in human myocardial infarction [29].

Angiotensin converting enzyme inhibitors are known to prevent both the generation of the potent vasoconstrictor angiotensin II and degradation of the powerful vasodilator bradykinin, which promotes endothelial cell release of NO [30]. In this study, rats treated with ISO had significant decreases in blood pressure parameters (SBD, DBP and MAP) when compared with the controls. This was however prevented in the ENA-treated group. There have been earlier reports of hypotension in subjects with acute myocardial infarction [31, 32]. From this study, it was interesting to observe that ENA, a known antihypertensive drug, was able to preserve the blood pressure measurements of ISO-treated rats comparable to the controls. This might have been a consequence of its ability to prevent myocardial infarction. Studies have actually shown that ACEIs have been used in the management of myocardial infarction [33, 34, 35]. Isoproterenol, a β -adrenergic agonist is known to produce stress in the myocardium due to the generation of free radicals by its auto-oxidation and some of the mechanisms proposed to explain its damage to cardiac myocytes include coronary hypotension, calcium overload, hypoxia, energy depletion and excessive production of free radicals as a result of catecholamine autoxidation [36, 37, 38]. The significant decrease in the levels of systolic, diastolic and mean arterial pressure may lead to coronary hypotension as seen in this study. In a study by Owens and O'Brien [39], it was concluded that in patients suffering with ischaemic heart disease and hypotension, symptomatic and silent ischaemia occurred in a temporally causal relation with hypotension, particularly for diastolic pressures. It thus suggests that patients with coronary disease may be susceptible to

456 ischaemic events that could be incurred as a result of low blood pressure. The enalapril used in
457 this study was able to restore the haemodynamic changes caused by isoproterenol indicating its
458 cardioprotective property.

459 In this study, the results of haematological analysis showed that ISO caused significant increase
460 in the levels of WBC, PCV, MCV and MCH while ENA caused significant decrease in WBC
461 and no changes in the erythrocyte indices relative to control. The increase in the level of WBC
462 could be explained in terms of necrosis caused by ISO leading to white blood cell mobilization
463 [11]. The significant reduction in the level of this parameter by enalapril could also be seen as its
464 ability to counteract the toxic effect of isoproterenol.

465 The toxicant also caused significant increase in the levels of AST and ALT while ENA caused
466 significant decrease in the levels of these enzymes. In heart failure, the heart has an impaired
467 ability to deliver blood to the body. Many reasons may be adduced to this, for instance a heart
468 attack can damage part of the muscular wall and because of that, the delivery of blood to the
469 organs such as liver and kidney decreases. In an attempt to raise the blood pressure the kidneys
470 may react by conserving fluid and electrolytes and thus puts more strain on the heart, worsening
471 the condition. The fluid in the body builds up even more. The liver can become dysfunctional,
472 and liver enzymes can be released into the blood. The liver enzymes can be released for a
473 number of reasons. First, the liver may be receiving an inadequate supply of blood, thus
474 damaging the cells and secondly, because the heart is not effectively pumping blood, blood and
475 fluids can back up into the liver, further damaging the cells [40]. It thus means that the increases
476 noted for the liver enzymes in this study implied that isoproterenol could impair liver functions
477 and this was counteracted by enalapril indicating that enalapril has beneficial effect beyond
478 being an ACE inhibitor.

479 It was also observed that ISO caused significant decrease in the level of NO while ENA caused
480 significant increase. Nitric oxide (NO) is known to play important functional roles in a variety of
481 physiological systems. For instance within the vasculature, NO induces vasodilation, inhibits
482 platelet aggregation, prevents neutrophil/platelet adhesion to endothelial cells, inhibits smooth
483 muscle cell proliferation and migration, regulates programmed cell death (apoptosis) and
484 maintains endothelial cell barrier function [41]. Nitric oxide (NO) is known to be deficient in
485 chronic progressive renal disease (CRD) and in end-stage renal disease (ESRD) [42, 43] and this

could result from arginine deficiency [44] which may be caused by a loss of functional renal mass, increased endogenous NO synthase (NOS) inhibitors that accumulate in renal failure [44], and/or other causes, such as increased oxidant stress [45]. Low NO production may also contribute to and/or exacerbate the progression of CRD by both hemodynamic and renal growth-promoting actions [46]. It should also be noted that NO blockade can lead to increased blood pressure and attenuated or delayed the hypotensive effect of all ACE inhibitors [47]. ACE inhibitors such as enalapril also augment the hemodynamic vasodilator action of bradykinin [48]. The increased level of NO in this study due to enalapril may further affirm its antihypertensive property and hence cardioprotective effect.

ISO caused significant increase in the levels of oxidative stress markers such as MDA, H_2O_2 and MPO while ENA caused significant decrease in the levels of these markers in a similar fashion to the control. Oxidative stress constitutes an alteration produced by disequilibrium between generation of free radicals (FR) and the antioxidant system, which can lead to a damage state, in particular of the biomolecules [49]. FR generation is related to the development and evolution of diverse illnesses such as atherosclerotic disease, high blood pressure, renal disorders, and obesity [50]. FR plays a primordial role in the development of long-term complications of these illnesses [51, 52, 53]. FR generates the lipid peroxidation process in an organism. Malondialdehyde (MDA) is one of the final products of polyunsaturated fatty acids (PUFA) peroxidation in the cells. An increase in free radicals causes overproduction of MDA content. Malondialdehyde level is commonly used as a marker of oxidative stress [54]. Myeloperoxidase (MPO), is abundant in the granules of inflammatory cells and it is an important enzyme in the generation of reactive oxygen species (ROS) by conversion of hydrogen peroxide to species including $\bullet OH$, ONOO $^-$, hypochlorous acid (HOCl) and NO $_2$ [55, 56, 57]. Hydrogen peroxide (H_2O_2), a ROS has been suggested as a mediator of vascular structural and functional alterations observed in hypertension [58, 59, 60]. Vasoconstrictor responses to H_2O_2 in rat aorta [61] and superior mesenteric artery [62] are also known to increase in hypertension. The H_2O_2 -induced contraction was found to be mediated by augmented thromboxane (TXA $_2$) release [62]. The reduction of these oxidative markers by enalapril is a pointer to its ability to scavenge the radicals generated by the toxicant and it thus showed that enalapril has anti-oxidant activity. In fact, De Cavanagh et al [63] reported that enalapril inhibits free radical formation and attenuates oxidative stress and

also prevents damage to the liver and kidney. This was further confirmed by the ability of this ACE inhibitor to increase the levels of antioxidant enzymes such as SOD, GPx, GST and GSH evaluated in this study. This view is clearly supported by a study carried out by Chandra et al [64], where it was concluded that enalapril has anti-oxidative property and this may have been responsible for its cardioprotective property. As a matter of fact, ENA caused a significant increase in the levels of protein thiols and non protein thiols further confirming its anti-oxidant property. It will be recalled that thiol compounds occupy a pivotal role in cellular metabolism especially as it relates to their essential function in the maintenance of cellular redox balance and their role in controlling oxidative stress, gene expression [65, 66] and redox signalling [67]. Cells have evolved several antioxidant strategies aimed at the detoxification of ROS with glutathione redox cycle as one of the major protective systems against oxidant damage. This cycle composed of the enzymes glutathione peroxidase (GPx) and glutathione reductase (GSSG-Rd) and the cosubstrates glutathione and NADPH [68]. Glutathione is the most abundant nonprotein intracellular thiol, and has a multiple role as an antioxidant agent. It functions as a scavenger of ROS, including hydroxyl radicals, singlet oxygen, nitric oxide, and peroxynitrite. In addition, GSH is a cosubstrate for the detoxification of peroxides by GPx and of toxic metabolites by glutathione-S-transferases [69]. Though the mechanism(s) underlying the enhancement of glutathione and glutathione-related enzymes by ACEI remains unknown, however, tissue glutathione levels and GSSG-Rd and GPx activities have been shown to increase in response to experimentally induced oxidative stress [70]. Studies have shown that 11-wk enalapril or captopril treatments increased antioxidant enzymes and nonenzymatic antioxidant defenses in several mouse tissues [71, 72]. ACEI decrease angiotensin II formation as well as endogenous bradykinin degradation. As a result, long-term ACE inhibition promotes the accumulation of the latter substance [73]. Bradykinin is a potent vasodilator known to stimulate the release of nitric oxide [74]. All these showed that enalapril may have shown its cardioprotective property through its anti-oxidant effect.

In this study, histopathological examinations showed that while there was severe infiltration of inflammatory cells into the cardiac tissue of the ISO group, there was no visible lesion seen in the ENA and control groups (Figure 13). This increase in the inflammatory cells may have been responsible for the increase in the levels of WBC noted in this study (Table 1). It should be noted

that the isoproterenol-induced myocardial alterations are similar in certain respects to those occurring in human beings following a myocardial infarction [75]. It is thought that the β -adrenergic cardiostimulatory activity exerted by ISO increases cardiac oxidative metabolism to a level that exceeds the amount of oxygen available to the myocytes through the unobstructed coronary circulation. The area of the heart most susceptible to hypoxia caused by tachycardia appears to be the left ventricular subendocardium [76, 77]. Myocyte damage observed following exposure to ISO includes both apoptosis and necrosis [78]. In the study on the isoproterenol-induced myocardial damage, it was discovered that the cardiac lesions varied with treatment duration and doses and that numerous macrophages were observed in the necrotic areas. It was inferred that the coexistence of interstitial oedema, inflammatory infiltration, myocardial basement damage, and myocardial degeneration was interpreted as indicating potential reversible lesions. It was inferred that these changes are not necessarily the most important factors involved in the pathogenesis of cell death but with higher doses and longer duration, the coexistence of apoptosis, necrosis with cell membrane rupture, and fibroblast proliferation was interpreted as indicating the presence of irreversible cell damage [79]. In a study on the effect of the antihypertensive drug enalapril on oxidative stress markers and antioxidant enzymes in kidney of Spontaneously Hypertensive Rat (SHR), the histopathology results of this study confirmed the effect of L-NAME in producing kidney damage because clear pathological changes were seen in the glomerulus, tubules, and blood vessels at 28 weeks [62]. In that study, it was observed that enalapril treatment managed to prevent this damage hence confirming its renoprotective effect through blood pressure lowering as mentioned by some researchers [80]. In our study, enalapril did not show any visible cardiac tissue damage possibly through its ability to prevent cell infiltration thus preventing apoptosis and necrosis.

The immunohistochemical analysis showed that there was high expressions of cardiac troponin and CRP in ISO group but lower expression of these proteins in ENA and control groups (Figures 14 and 15). In the case of IL-10, there was low expression of this protein in ISO group but higher expression in ENA and control groups (Figure 16). Cardiac troponins are regulatory proteins within the myocardium that are released into the circulation when damage to the myocyte has occurred. Therefore, serum troponin is an exquisitely sensitive marker of myocardial injury and is necessary for establishing the diagnosis of MI. Troponin (I or T) has

576 demonstrated nearly absolute myocardial tissue specificity as well as high clinical sensitivity for
 577 myocardial ischemia [81, 82]. Troponin is the preferred biomarker for the detection of
 578 myocardial necrosis and is a Class I indication for the diagnosis of MI [82, 83, 84]. Cardiac
 579 troponin T (cTnT) and I (cTnI) have been shown to be specific and sensitive biomarkers of drug-
 580 induced myocardial cell injury in animals and humans [85, 86]. In rats, a number of studies have
 581 described a relationship between the serum levels of cTnT or cTnI and the severity of
 582 isoproterenol-induced cardiotoxicity [87, 88, 89]. This study has shown that ISO caused
 583 myocardial injury with upregulation of this biomarker. On the other hand, the down regulation of
 584 cardiac troponin by ENA also showed that this drug has ability to protect against myocardial
 585 injury in rats. It was shown that an increase in troponin I soon after high-dose chemotherapy
 586 (HDC) is a strong predictor of poor cardiological outcome in cancer patients. For instance, in a
 587 study conducted by Cardinale et al [90], it was concluded that in high-risk, HDC-treated patients,
 588 defined by an increased troponin I value, early treatment with enalapril seems to prevent the
 589 development of late cardiotoxicity. C-reactive protein (CRP) has the capacity to precipitate the
 590 somatic C-polysaccharide of *Streptococcus pneumonia*. It was the first acute-phase protein to be
 591 described and is an exquisitely sensitive systemic marker of inflammation and tissue damage
 592 [91]. It is a known fact that tissue necrosis is a potent acute-phase stimulus. In myocardial
 593 infarction, there is a major CRP response with the magnitude of this response indicating the
 594 extent of myocardial necrosis [92]. In all acute myocardial infarcts, CRP is co-deposited with
 595 activated complement [93, 94], and research findings have shown that the CRP response did not
 596 only reflects tissue damage in this context but may actually contribute significantly to the
 597 severity of ischemic myocardial injury [95]. It is very clear that CRP plays a role in the
 598 pathogenesis of cardiovascular disease and as a marker and predictor of cardiovascular disease,
 599 CRP possesses numerous cardiovascular effects including clotting, generation of oxygen
 600 radicals, increase in the expression of adhesion molecules and plasminogen activator inhibitor-1,
 601 plaque destabilization and these could result in cardiovascular disease. Prasad [96] in a review
 602 describes the effects of various cardiovascular drugs on the levels of CRP in health and disease
 603 where it showed that cyclooxygenase inhibitors such as aspirin, rofecoxib, celecoxib; platelet
 604 aggregation inhibitors such as clopidogrel, abciximab; lipid lowering agents including statins,
 605 ezetimibe, fenofibrate, niacin, diets; beta-adrenoreceptor antagonists and antioxidants (vitamin

E), as well as angiotensin converting enzyme (ACE) inhibitors (ramipril, captopril, fosinopril), reduce serum levels of CRP but enalapril and trandolapril have not been shown to have the same effect. The lowering of the level of CRP in this study by ENA is a pointer to its ability to halt cardiovascular disease hence cardioprotective effect through its anti-oxidant and anti-inflammatory properties. Immunohistochemistry in this study further showed that ENA caused increased level of IL-10. IL-10 is a Th₂-type cytokine that is produced by a wide range of immunological cell types, including monocytes/macrophages, and it is a potent inhibitor of the proinflammatory cytokines and chemokines [97]. Immunosuppressive effects of IL-10 involve both inhibition of cytokine synthesis (e.g., TNF- α , IL-6) and their biological activities on target cells [98]. It has also been reported that cotreatment with IL-10 prevents muscle insulin resistance following an acute lipid infusion [99]. Studies have shown that endogenous IL-10 limits angiotensin II (ANG II)-mediated oxidative stress, inflammation and vascular dysfunction both *in vivo* and *in vitro*, indicating a protective action of IL-10 in vascular diseases such as arterial hypertension [100]. As a matter of fact, IL-10 attenuates the increases in vascular superoxide and endothelial dysfunction during diabetes and atherosclerosis [101, 102]. In the same way, it could be suggested that IL-10 might be a mediator of cardiac protection against arterial hypertension. In a study, enalapril was able to produce an increase in plasma levels of IL-10 in patients with coronary artery disease and arterial hypertension [103]. It thus shows that the cardioprotective effect of enalapril may also be linked to its anti-inflammatory property as shown by the up regulation of IL-10.

In conclusion, this study has shown that enalapril, an ACE inhibitor has cardioprotective properties which it exhibited through its anti-oxidant, anti-inflammatory and anti-apoptotic effects. Its antihypertensive property is also exhibited through its nitric oxide increasing ability leading to vasodilation and hence decreases in peripheral resistance. Other studies have also shown its nephroprotective [63, 64] as well as antidiabetic properties [63]. This study thus showed that much is still needed to be explored on this very important drug, enalapril.

634 Conflicts of interest

635 We have no conflict of interest to declare

636 References

- 637 1. Murray CJ and Lopez AD. Global Burden of Disease and Injury Series, Global Health
638 Statistics. Boston: Harvard School of Public Health 1996; *Vols. I and II.* .
- 639 2. Gatica D, Chiong M, Lavandero S and Klionsky DJ. Molecular mechanisms of
640 autophagy in the cardiovascular system. *Circulation Research* 2015; 116(3), 456-467.
- 641 3. Hina SK, Rehman ZH, Dogar N, Jahan M, Hameed ZI, Khan K, Ahmad K, Mukhtar and
642 Valeem EE. Cardioprotective effect of gemmotherapeutically treated *Withania somnifera*
643 against chemically induced myocardial injury. *Pak J Bot* 2010; 42: 1487-1499.
644
- 645 4. Boudina S, Laclau MN and Tariosse L. “Alteration of mitochondrial function in a model
646 of chronic ischemia in vivo in rat heart,” *American Journal of Physiology—Heart and*
647 *Circulatory Physiology* 2002, vol. 282, no. 3, pp. H821–H831.
648
- 649 5. Mohanty DS, Arya A, Dinda KK, Talwar S, Joshi and Gupta SK. “Mechanisms of
650 cardioprotective effect of *Withania somnifera* in experimentally induced myocardial
651 infarction,” *Basic and Clinical Pharmacology & Toxicology* 2004; vol. 94, no. 4, pp.
652 184–190.
653
- 654 6. Sabeena Farvin KH, Anandan R, Kumar SHS, Shiny KS, Sankar TV and Thankappan
655 TK. “Effect of squalene on tissue defense system in isoproterenol-induced myocardial
656 infarction in rats,” *Pharmacological Research* 2004, vol. 50, no. 3, pp. 231–236.
657
- 658 7. Abel ED. Glucose transport in the heart. *Front Biosci* 2004; 9: 201–215.
659
- 660 8. Stanley WC, Recchia FA and Lopaschuk GD. Myocardial substrate metabolism in the
661 normal and failing heart. *Physiol Rev* 2005; 85: 1093–1129.
662
- 663 9. Nagoshi T, Yoshimura M, Giuseppe M, Rosano C, Gary D, Lopaschuk C and Mochizuki
664 S. Optimization of cardiac metabolism in heart failure. *Current Pharmaceutical Design*
665 2011; 17: 3846-3853.
666
- 667 10. Wexler BC. Myocardial infarction in young versus old male rats: pathophysiological
668 changes. *Am Heart J* 1987; 96: 70–80

669

670 11. Khalil MI, Ahmmed I, Ahmed R, Tanvir EM, Afroz R, Paul S, Gan SH and Alam N.
671 Amelioration of isoproterenol-induced oxidative damage in rat myocardium by *Withania*
672 *somnifera* leaf extract. *BioMed Research International Article* 2015. ID 624159, 10 pages
673

674 12. Rathore N, John S, Kale M and Bhatnagar D. Lipid peroxidation and antioxidant
675 enzymes in isoproterenol induced oxidative stress in rat tissues. *Pharmacol Res* 1998;38,
676 297-303

677 13. Banerjee SK, Sood S, Dinda AK, Das TK and Maulik SK. Chronic oral administration of
678 raw garlic protects against isoproterenol-induced myocardial necrosis in rat. *Comp*
679 *Biochem Physiol* 2003 Part C 136, 377-386.
680

681 14. Nirmala C and Puvanakrishnan R. Protective role of curcumin against isoproterenol
682 induced myocardial infarction in rats. *Mol Cell Biochem* 1996; 159, 85-93.

683 15. Jackson, Edwin K. "Chapter 30. Renin and Angiotensin". In Brunton, Laurence L.; Lazo,
684 John S.; Parker, Keith. Goodman & Gilman's The Pharmacological Basis of Therapeutics
685 (11th ed.) 2006. New York: McGraw-Hill

686 16. Wang W, McKinnie SM, Farhan M, Paul M, McDonald T, McLean B, Llorens-Cortes C,
687 Hazra S, Murray AG, Vederas JC and Oudit GY. "Angiotensin Converting Enzyme 2
688 Metabolizes and Partially Inactivates Pyrapelin-13 and Apelin-17: Physiological Effects
689 in the Cardiovascular System". *Hypertension* 2016; 68(2): 365-377.

690 17. Gornal AG, Bardawill JC, David MM. Determination of serum proteins by means of
691 Biuret reaction. *J Biol Chem* 1949; 177:751e66.
692

693 18. Buetler E, Duron O, Kelly BM. Improved method for the determination of blood
694 glutathione. *J Lab Clin Med* 1963; 61:882e8.
695

696 19. Rotruck JT, Pope AL, Ganther HE, Swanson AB, Hafeman DG and Hoekstra WG.
697 Seleniu biochemical role as a component of glutathione peroxidase. *Sci* 1973;
698 179:588e90.
699

700 20. Habig WH, Pabst MJ and Jacoby WB. Glutathione-S-transferase activity: the enzymic
701 step in mercapturic acid formation. *J Biol Chem* 1974; 249:130e9.
702

703 21. Misra HP and Fridovich I. The role of superoxide anion in the autoxidation of
704 epinephrine and a simple assay for superoxide dismutase. *J Biol Chem.* 1972; 25:
705 247(10): 3170–3175.
706

707 22. Varshney R and Kale RK. Effect of calmodulin antagonists on radiation induced lipid
708 peroxidation in microsomes. *Intern J Biol* 1990; 158:733e41.

23. Wolff SF. Ferrous ion oxidation in the presence of ferric ion indicator xylenol orange for measurement of hydrogen peroxides. *Methods Enzymol* 1994; 233: 182e9.
24. Ellman GL. Tissue sulfhydryl groups. *Arch. Biochem. Biophys* 1959; 82 (1): 70-77.
25. Olaleye SB, Adaramoye OA, Erigbali PP. Lead exposure increases oxidative stress in the gastric mucosa of HCl/ethanol-exposed rats. *World J Gastroent.* 2007; 13: 5121-5126.
26. Drury R, Wallington E and Cancerson R. Carleton's histological technique". 4th ed. 1976. Oxford University Press, London.
27. Bono DP and Boon NA. Diseases of cardiovascular system. In Davidson's Principles and Practice of Medicine. Edited by Edwards CRW, Boucheir IA. Hong Kong: Churchill Livingstone 1992; pp 249–340.
28. Nian M, Lee P, Khaper N and Liu P. Inflammatory cytokines and postmyocardial infarction remodeling. *Circ Res* 2004; 94: 1543-1553.
29. Wang J, Bo H, Meng X, Wu Y, Bao Y and Li Y. A simple and fast experimental model of myocardial infarction in the mouse. *Tex Heart Inst J* 2006; 33: 290-293.
30. Kerth PA and Vanhoutte PM. Effects of perindoprilat on endothelium-dependent relaxations and contractions in isolated blood vessels. *Am J Hypertens* 1991; 4:226S-234S.
31. Wagner JA, Hu K, Bauersachs J, Karcher J, Wiesler M, Goparaju SK, Kunos G and Ertl G. Endogenous cannabinoids mediate hypotension after experimental myocardial infarction. *J Am Col Cardiol* 2001; 38 (7): 2048-2054.
32. Ohman EM, Nanas J, Stomel RJ, Leeser MA, Nielsen DW, O'Dea D, Rogers FJ, Harber D, Hudson MP, Fraulo E and Shaw LK. Thrombolysis and counterpulsation to improve survival in myocardial infarction complicated by hypotension and suspected cardiogenic shock or heart failure: results of the TACTICS Trial. *J Thrombos Thrombol* 2005; 19(1): 33-39.
33. Hennekens CH, Albert CM, Godfried SL, Gaziano JM, Buring JE. Adjunctive drug therapy of acute myocardial infarction: evidence from clinical trials. *N Engl J Med* 1996; 335: 1660–1667

34. ACEIMICG. Indications for ACE inhibitors in the early treatment of acute myocardial infarction: systematic overview of individual data from 100,000 patients in randomized trials. ACE Inhibitor Myocardial Infarction Collaborative Group. Jun 1198; 97(22): 2202-2212
35. Lubarsky L and Coplan NL. Angiotensin-Converting Enzyme Inhibitors in Acute Myocardial Infarction: A Clinical Approach. *Preventive cardiology* 2007; 10(3): 156-159.
36. Rona G, Chappel CI, Balazs T and Gaudry R. An infarct-like myocardial lesion and other toxic manifestations produced by isoproterenol in the rat. *Arch Pathol* 1959; 76: 443-445.
37. Adameova A, Abdellatif Y and Dhalla NS. Role of excessive amounts of circulating catecholamines and glucocorticoids in stress-induced heart disease. *Can J Physiol Pharmacol* 2009; 87: 493-514.
38. Upaganlawar A and Balaraman R. Cardioprotective effects of *Lagenaria siceraria* fruit juice on isoproterenol-induced myocardial infarction in Wistar rats: A biochemical and histoarchitecture study. *J Young Pharmacists* 2011; 3: 297-303.
39. Owens P and O'Brien E. Hypotension in patients with coronary disease: can profound hypotensive events cause myocardial ischaemic events? *Heart* 1999; 82: 477-481.
40. Macfarlane PS, Reid R and Callander R. Pathology Illustrated. Int Student 5th Ed. Churchill Livingstone, London 2000; pp 179-188.
41. Rosselli M, Keller PJ and Dubey RK. Role of nitric oxide in the biology, physiology and pathophysiology of reproduction. *Human Reproduction Update* 1998; 4 (1): 3-24.
42. Vallance P, Leone A, Calver A, Collier J and Moncada S. Accumulation of an endogenous inhibitor of nitric oxide synthesis in chronic renal failure. *Lancet* 1992; 339: 572-575.
43. Reyes AA, Karl IE and Klahr S. Role of arginine in health and in renal disease. *Am J Physiol* 1994; 267: F331-F346.

- 782 44. Morris SM Jr. Regulation of enzymes of urea and arginine synthesis. *Annu Rev Nutr*
783 1992; 12: 81–101.
784
- 785 45. Vaziri ND, Ovelisi F and Ding Y. Role of increased oxygen free radical activity in the
786 pathogenesis of uremic hypertension. *Kidney Int* 1998; 53: 1748–1754.
787
- 788 46. Zatz R and Baylis C. Chronic nitric oxide inhibition model six years on. *Rev Hypertens*
789 1998; 32: 958–964.
790
- 791 47. Cachofeiro V, Sakakibara T and Nasjletti A. Kinins, nitric oxide, and the hypotensive
792 effect of captopril and ramiprilat in hypertension. *Hypertension* 1992; 19:138-145
793
- 794 48. Bonner G, Preis S, Schunk U, Toussaint C and Kaufmann W. Hemodynamic effects of
795 bradykinin on systemic and pulmonary circulation in healthy and hypertensive humans. *J*
796 *Cardiovasc Pharmacol* 1990; 15(suppl 6):S46-S56.
797
- 798 49. Touyz RM: Oxidative stress and vascular damage in hypertension.” *Current hypertension*
799 *reports* 2000; 2 (1): 98–105.
800
- 801 50. Wilcox CS. Reactive oxygen species: roles in blood pressure and kidney function,”
802 *Current Hypertension Reports* 2002; 4 (2): 160–166.
803
- 804 51. Sabban EL and Kvetnansky R. Stress-triggered activation of gene expression in
805 catecholaminergic systems: dynamics of transcriptional events. *Trends Neurosci* 2001;
806 24(2): 91–98.
807
- 808 52. Esch T, Sefano GB and Fricchione GL, Benson H. Stress in cardiovascular diseases.
809 *Med. Sci. Monit* 2002; 8: RA93–RA101.
810
- 811 53. Agrawal A and Sharma B. Pesticides induced oxidative stress in mammalian systems. *Int*
812 *J Biol Med Res* 2010; 1(3): 90-104.
813
- 814 54. Maddock C and Pariente CM. How does stress affect you? An overview of stress,
815 immunity, depression, and disease. *Epidemiol. Psychiatr. Soc.* 2001; 10(3): 153-62.
816

55. Furtmuller, P.G., Arnhold, J., Jantschko, W., Pichler, H and Obinger, C. Redox properties of the couples compound I/compound II and compound II/native enzyme of human myeloperoxidase. *Biochem Biophys Res Commun* 2003; 301: 551–557.
56. Arnhold J, Monzani E, Furtmüller PG., Zederbauer M., Casella L and Obinger C. Kinetics and thermodynamics of halide and nitrite oxidation by mammalian heme peroxidases. *Eur J Inorg Chem* 2006; 19: 3801–3811.
57. Zederbauer M, Furtmüller PG, Brogioni, S, Jakopitsch, C, Smulevich, G, and Obinger, C. Heme to protein linkages in mammalian peroxidases: impact on spectroscopic, redox, and catalytic properties. *Nat Prod Rep* 2007; 24: 571–584.
58. Lacy F, Kailasam MT, O'Connor DT, Schmid-Schoenbein GW and Parmer RJ. Plasma hydrogen peroxide production in human essential hypertension: role of heredity, gender, and ethnicity. *Hypertension* 2000; 36: 878–884.
59. Paravicini TM and Touyz RM. Redox signalling in hypertension. *Cardiovasc Res* 2006; 71:247–258.
60. Alvarez Y, Pe´rez-Giro´n JV, Hernanz R, Briones AM, Garc´ıa-Redondo A, Beltra´n A, Alonso MJ, and Salaices M. Losartan reduces the increased participation of cyclooxygenase-2-derived products in vascular responses of hypertensive rats. *J Pharmacol Exp Ther* 2007; 321:381–388.
61. Rodr´ıguez-Mart´ınez MA, Garc´ıa-Cohen EC, Baena AB, Gonz´alez R, Sala´ıces M and Mar´ın J. Contractile responses elicited by hydrogen peroxide in aorta from normotensive and hypertensive rats: endothelial modulation and mechanism involved. *Br J Pharmacol* 1998; 125:1329–1335.
62. Gao YJ and Lee RM. Hydrogen peroxide induces a greater contraction in mesenteric arteries of spontaneously hypertensive rats through thromboxane A (2) production. *Br J Pharmacol* 2001; 134:1639–1646.
63. De Cavanagh EMV, Inserra F, Toblli J, Stella I, Fraga CG and Ferder L. Enalapril attenuates oxidative stress in diabetic rats. *Hypertension* 2001; 38: 1130–1136.
64. Chandran G, Sirajudeen KNS, Yusoff NSN, Swamy M and Samarendra MS. Effect of the antihypertensive drug enalapril on oxidative stress markers and antioxidant enzymes

- 855 in kidney of spontaneously hypertensive rat. *Oxidative Medicine and Cellular Longevity*.
856 2014; Article ID 608512, 10 pages.
857
- 858 65. Packer L. “*Biothiols*” *Methods in Enzymology*; Academic Press Inc.: London, England,
859 1995; volumes 251, Part A & 252 Part B.
860
- 861 66. Arrigo AP. Gene expression and the thiol redox state. *Free Rad. Bio. Med.* 1999, 27,
862 936–944.
863
- 864 67. Bindoli A, Fukato JM and Forman HJ. Thiol Chemistry in peroxidase catalysis and redox
865 signalling. *Antioxid. Redox Sign.* 2008, 10, 1549–1564.
866
- 867 68. Reed DJ. Oxidative stress and mitochondrial permeability transition. in *Biothiols in*
868 *Health and Disease*, eds Packer L., Cadenas E. (Dekker, New York) 1995; pp 231–263.
869
- 870 69. Halliwell B and Gutteridge JM. Protection against oxidants in biological systems: the
871 superoxide theory of oxygen toxicity. *Free Radicals in Biology and Medicine* (Clarendon,
872 Oxford, UK), 2nd ed 1989; pp 87–187.
873
- 874 70. Forman HJ, Liu R.-M, Shi MM, Packer L and Cadenas E. Glutathione synthesis in
875 oxidative stress. *Biothiols in Health and Disease* (Dekker, New York) eds 1995, pp 189–
876 212.
877
- 878 71. Cavanagh EMV, de Inserra F, Ferder L, Romano L, Ercole L and Fraga CG. Superoxide
879 dismutase and glutathione peroxidase activities are increased by enalapril and captopril in
880 mouse liver. *FEBS Lett* 1995; 361:22–24.
881
- 882 72. Cavanagh EMV, de Fraga CG, Ferder L and Inserra F. (1997) Enalapril and captopril
883 enhance antioxidant defenses in mouse tissues. *Am. J. Physiol. Regulatory Integrative*
884 *Comp. Physiol* 1997; 272:R514–R518.
885

- 886 73. Bonner G, Schunk U and Kaufman W. Direct hypotensive action of intravascular
887 bradykinin in man. *Cardiology* 1985; 72, Suppl.1:190–193.
- 888
- 889 74. Gohlke P and Unger T. Chronic low-dose treatment with perindopril improves cardiac
890 function in stroke-prone spontaneously hypertensive rats by potentiation of endogenous
891 bradykinin. *Am. J. Cardiol* 1995; 76:41E–45E.
- 892
- 893 75. Wexler BC and Greenberg BP. Protective effect of clofibrate on isoproterenol-induced
894 myocardial infarction in arterio-sclerotic and nonarterio-sclerotic rats. *Atherosclerosis*
895 1978; 29, 373–75.
- 896
- 897 76. Balazs T, Hanig JP, and Herman EH. Toxic responses of the cardiovascular system. In
898 Casarett and Doull's Toxicology: the Basic Science of Poisons (C. D. Klaassen, M. O.
899 Amdur, and J. Doull, eds), Macmillan Publishing Company, New York, USA. third
900 edition 2006; pp 387–411.
- 901
- 902 77. Van Vleet JF, Ferrans, JV, and Herman E. Cardiovascular and skeletal muscle system. In
903 Handbook of Toxicologic Pathology (W. M. Haschek, C. G. Rousseaux, and M. A.
904 Wallig, eds), Academic Press, San Diego, CA, USA 2002; Vol. 2, pp 363–455..
- 905
- 906 78. Goldspink DF, Burniston JG, Ellison GM, Clark WA and Tan L.-B. Catecholamine-
907 induced apoptosis and necrosis in cardiac and skeletal myocytes of the rat in vivo: the
908 same or separate death pathways? *Exp Physiol* 2004; 89, 407–16.
- 909
- 910 79. Zhang J, Knapton A, Lipshultz SE, Weaver JL, and Herman EH. Isoproterenol-induced
911 cardiotoxicity in Sprague-Dawley rats: correlation of reversible and irreversible
912 myocardial injury with release of cardiac troponin T and roles of iNOS in myocardial
913 injury. *Toxicologic Pathology* 2008; 36:277-288.
- 914
- 915 80. Rugale C, Cordaillat M, Mimran A and Jover B. Prevention and reversal by enalapril of
916 target organ damage in angiotensin II hypertension. *Journal of the Renin-Angiotensin-*
917 *Aldosterone System* 2005; 6 (3): 154–160.
- 918
- 919 81. Gerhardt W, Nordin G and Ljungdahl L. Can troponin T replace CK MBmass as “gold
920 standard” for acute myocardial infarction (“AMI”)? *Scand J Clin Lab Invest Suppl.* 1999;
921 230: 83–89.

- 922
- 923 82. Morrow DA, Cannon CP, Jesse RL, Newby K, Ravkilde J, Storrow AB, Wu AHB and
 924 Christenson RH. National Academy of Clinical Biochemistry Laboratory Medicine
 925 Practice Guidelines: clinical characteristics and utilization of biochemical markers in
 926 acute coronary syndromes. *Clin Chem.* 2007; 53(4):552–574.
 927
- 928 83. Jaffe AS, Ravkilde J, Roberts R, Naslund U, Apple FS, Galvani M and Katus H. It's time
 929 for a change to a troponin standard. *Circulation* 2002; 102: 1216-1220.
 930
- 931 84. Thygesen K, Alpert JS and White HD. Universal definition of myocardial infarction. *J*
 932 *Am Coll Cardiol.* 2007; 50 (22): 2173–2195.
 933
- 934 85. Bertsch T, Bleuel H, Aufenanger J and Rebel W. Comparison of cardiac troponin T and
 935 cardiac troponin I concentrations in peripheral blood during orcipenaline induced
 936 tachycardia in rats. *Exp Toxic Pathol* 1997; 49: 467–68.
 937
- 938 86. Wallace KB, Hausner E, Herman E, Holt GD, MacGregor JT, Metz,AL et al. Serum
 939 troponins as biomarkers of drug-induced cardiac toxicity. *Toxicol Pathol* 2004; 32: 106–
 940 21.
 941
- 942 87. Bleuel, H., Deschl, U., Bolz, G., and Rebel, W. (1995). Diagnostic efficiency of troponin
 943 T measurements in rats with experimental myocardial cell damage. *Exp Toxic Pathol* 47,
 944 121–27.
 945
- 946 88. Bertinchant JP, Robert E, Polge A, Marty-Double C, Fabbro-Peray P, Poirey S, Aya G,
 947 Juan JM, Ledermann B, de la Coussaye JE and Dauzat M. Comparison of the diagnostic
 948 value of cardiac troponin I and T determination for detecting early myocardial damage
 949 and the relationship with histological findings after isoprenaline-induced cardiac injury in
 950 rats. *Clin Chim Acta* 2000; 298, 13–28.
- 951 89. Herman E, Zhang J, Knapton A, Lipshultz, SE, Rifai N and Sistare F. Serum cardiac
 952 troponin T as a biomarker for acute myocardial injury induced by low doses of
 953 isoproterenol in rats. *Cardiovasc Toxicol* 2006; 6: 211–22.
 954
- 955 90. Cardinale D, Colombo A, Sandri MT, Lamantia G, Colombo N, Civelli M, Martinelli G,
 956 Veglia F, Fiorentini C and Cipolla CM. Prevention of High-Dose Chemotherapy–induced
 957 cardiotoxicity in high-risk patients by angiotensin-converting enzyme inhibition.
 958 *Circulation* 2006; 114: 2474-2481.
 959

91. Pepys MB and Baltz ML. Acute phase proteins with special reference to C-reactive protein and related proteins (pentaxins) and serum amyloid A protein. *Adv. Immunol.* 1983; 34: 141–212.
92. de Beer FC, Hind CR, Fox KM, Allan RM, Maseri A, and Pepys MB. Measurement of serum C-reactive protein concentration in myocardial ischaemia and infarction. *Br. Heart J.* 1982; 47: 239–243.
93. Kushner I, Rakita L and Kaplan MH. Studies of acute phase protein. II. Localization of Cx-reactive protein in heart in induced myocardial infarction in rabbits. *J. Clin. Invest.* 1963; 42: 286–292.
94. Lagrand WK, Niessen HW, Wolbink GJ, Jaspars LH, Visser CA, Verheugt FW, Meijer CJ and Hack CE. C-reactive protein colocalizes with complement in human hearts during acute myocardial infarction. *Circulation.* 1997;95: 97–103.
95. Griselli M, Herbert J, Hutchinson WL, Taylor KM, Sohail M, Krausz T and Pepys MB. C-reactive protein and complement are important mediators of tissue damage in acute myocardial infarction. *J. Exp. Med.* 1999; 190: 1733–1739.
96. Prasad K. C-reactive protein (CRP)-lowering agents. *Cardiovasc Drug Rev* 2006; 24(1): 33-50.
97. Akdis CA and Blaser K (2001): Mechanisms of interleukin-10-mediated immune suppression. *Immunology* 2001; 103:131–136.
98. Pestka S, Krause CD, Sarkar D, Walter MR, Shi Y, Fisher PB (2004): Interleukin-10 and related cytokines and receptors. *Annu Rev Immunol* 2004; 22: 929–979
99. Kim HJ, Higashimori T, Park SY, Choi H, Dong J, Kim YJ, Noh HL, Cho YR, Cline G, Kim YB, Kim JK. Differential effects of interleukin-6 and -10 on skeletal muscle and liver insulin action in vivo. *Diabetes* 2004; 53:1060–1067.

- 992 100. Didion SP, Kinzenbaw DA, Schrader LI, Chu Y, Faraci FM: Endogenous
993 interleukin-10 inhibits angiotensin II-induced vascular dysfunction. *Hypertension* 2009;
994 54: 619-624.

995

996 101. Mallat Z, Besnard S, Duriez M, Deleuze V, Emmanuel F, Bureau MF, Soubrier F,
997 Esposito B, Duez H, Fievet C, Staels B, Duverger N, Scherman D and Tedgui A.
998 Protective role of interleukin-10 in atherosclerosis. *Circ Res* 1999; 85: e17-e24.

999

1000 102. Gunnett CA, Heistad DD and Faraci FM. Interleukin-10 protects nitric oxide-
1001 dependent relaxation during diabetes: role of superoxide. *Diabetes* 2002; 51: 1931-1937.
1002

1003 103. Schieffer B, Bunte C, Witte J, Hoeper K, Boger RH, Schwedhelm E and Drexler
1004 H. Comparative effects of AT1-antagonism and angiotensin-converting enzyme
1005 inhibition on markers of inflammation and platelet aggregation in patients with coronary
1006 artery disease. *J Am Coll Cardiol* 2004; 44: 362-368.

1007