

IDENTIFICATION AND ISOLATION OF FUNGI IN ABATTOIR AND POULTRY AMENDED PLOTS IN ILORIN, SOUTHERN GUINEA SAVANNA.

ABSTRACT

Traditional methods for soil remediation are often expensive and energy consuming and this has given rise to a new ecologically safer method known as mycoremediation. A field experiment was carried out at the University of Ilorin Dam site to isolate and identify fungi present in organic amended plots which are capable of remediating heavy metal polluted soil of Asa River. Randomized Complete Block design in split plot arrangement was adopted using two treatments: Poultry Dropping (PD) and Abattoir Effluent (AE) at five levels i.e control, 1.3 t/ha, 2.6 t/ha, 1.3t/ha+NPK120KgN, 2.6t/ha+NPK120KgN for abattoir effluent, and poultry droppings at control, 10t/ha, 15t/ha, 10t/ha+NPK120KgN and 15t/ha+NPK120KgN having three (3) replicates. Soil samples collected before and after planting were analysed for heavy metals (Mn, Fe, Pb, Zn, Cu, Co, Ni, Cr, Cd) using Atomic Adsorption Spectrophotometer. Result obtained showed that organic waste are effective in bioremediation of Asa River sediment with abattoir effluent having a reduction of 99.04% and poultry dropping 98.72 % of heavy metal concentration in the soil which varied in the order of: Mn>Fe>Pb>Zn>Cu>Co>Ni>Cr>Cd. Result obtained also showed that eleven metal resistant Fungi were identified from abattoir effluent and poultry dropping amended plots i.e *Aspergillus niger*, *Trichoderma viride*, *Fusarium solani*, *Penicillium notatum*, *Aspergillus flavus*, *Trichoderma harzianum*, *Trichophyton verrucosum*, *Fusarium oxysporum*, *Stachybotrys chartarum*, *Aspergillus ustus* and *Microsporum nanum*. *Aspergillus niger* was observed to have the highest population (19% in poultry droppings and 19.6% in abattoir effluent).

31

INTRODUCTION

32 Industrialization, global increase in population and indiscriminate disposal of waste has
33 become a serious concern due to their release of heavy metals which has negative effect on
34 the environment. Several toxic metals from industrial wastewater and other human activities
35 are directly or indirectly released into the environment. Unlike organic contaminants, these
36 pollutants from heavy metals are not biodegradable and able to travel up the food chain via
37 bioaccumulation (45).

38 Several conventional methods have been used for the removal and treatment of heavy metal
39 in contaminated sites which includes electrochemical treatment, ion exchange, precipitation,
40 reverse osmosis, evaporation and sorption (23). However, these methods are uneconomical
41 due to high reagents and energy requirements, or ineffective at removing all metal
42 contamination, and they can generate large quantities of toxic sludge (20).

43 More importantly, bioremediation offers a safer and more economical option for treating
44 heavy metals in contaminated sites (21). The use of microorganisms and organic waste has
45 several advantages, not least the highly efficient removal of heavy metals and cost
46 effectiveness (15).

47 Some fungi are known to be particularly tolerant of heavy metals (18, 9), and many species
48 can adapt and grow under conditions of extreme pH, temperature, nutrient availability and
49 high metal concentrations (3). Fungi tolerate and detoxify metals through various
50 mechanisms involving valence transformation, extra- and intracellular precipitation or active
51 uptake (31, 39). Scientists are currently exploring the use of microbes and associated biota
52 residing in ecosystems for the bioremediation of pollutants through degradation or
53 accumulation (26), and numerous strains isolated from contaminated sites possess such
54 abilities.

55 Fungal biomass proves to be advantageous in having a high percentage of cell wall materials
56 which offers excellent metal binding properties (29, 28). Fungal cell walls can act as a cation
57 exchanger due to their negative charge originating from the presence of different functional
58 groups, e.g. carboxylic, phosphate, amine or sulfhydryl, in different wall components
59 (hemicelluloses, pectin, lignin, etc.). Cell walls of fungi are rich in polysaccharides and
60 glycoproteins such as glucans, chitin, mannans and phospho-mannans. Metals and their
61 compounds can interact with fungi in various ways depending on the type of metal, organism
62 and environment (6). Moreover the use of fungal biomasses as biosorption materials is very
63 convenient because of their inexpensive production methods based on simple fermentation

64 techniques (30). Fungi can also serve as an economical and constant supply of biomass for
65 biosorption of heavy metals and they can also be grown using inexpensive media and
66 unsophisticated fermentation techniques.

67 Many fungal species such as *Rhizopus arrhizus*, *Penicillium spinulum* and *Aspergillus niger*
68 have been extensively studied for heavy metals biosorption and the process mechanism seems
69 to be dependent upon species (19, 24).

70 Therefore, this research was aimed at using organic waste to bioremediate Asa River
71 Sediment and also to identify the fungi biomass present in the amended plot.

72

73

MATERIALS AND METHODS

74 Soil sediments were collected from Asa River and deposited at the surface of the soil at a rate
75 of 50 kg per plot (given a total of 41, 667 t/ha) to a depth of 30 cm on which planting was
76 done. The land was arranged in plots (3m x 4m) with four ridges per plot.

77 Abattoir effluent of 1.3 t/ha (1.6litres/plot) and 2.6 t/ha (3.2litres/plot) and poultry dropping
78 of 10t/ha and 15t/ha was incorporated into the soil at two weeks prior to planting to ensure
79 decomposition and mineralization, while NPK fertilizer (20:10:10) was applied 3 weeks after
80 planting at a rate of 120 kg per hectare.

81 Randomized Complete Block design was adopted, using two (2) treatments: Poultry
82 Dropping (PD) and Abattoir Effluent (AE) at two levels with or without NPK, having three
83 (3) replicate. The treatment combination followed a randomized arrangement.

84 Randomized Complete Block design in split-split plot arrangement was adopted, abattoir
85 effluent (1.3 t/ha and 2.6 t/ha) at two levels with or without NPK, having three (3) replicate.
86 The treatment combination followed a randomized arrangement. Two seed of Swan 1 maize
87 were planted per hole at a spacing of 75 x 25cm on a 3m x 4m plot having a total of 40holes
88 per plot. Supplementary weeding was done at 3 weeks after planting and six weeks after
89 planting prior to harvesting. Water was supplied through irrigation at an interval of 3days.

Chemical Analysis of Sediment

91 The pH was determined by the method outlined by (10) using an electronically Jenway 3015
92 pH meter at ratio of 1:2.5 soil/water. Exchangeable acidity of the soil was determined by
93 titration method using 1N KCl extract as described by (34). Exchangeable cations were
94 extracted with an excess of 1M NH₄OAc (ammonium acetate) where sodium and potassium
95 were determined using flame photometry, and atomic absorption spectrophotometer was used
96 to determine the calcium and magnesium (4). Organic carbon in sediment was determined

using wet oxidation method of (40), Available phosphorus was determined using Bray 1 (13) method. Total Nitrogen was determined using micro-Kjelhdal distillation method by (7).

Heavy metals analysis of soil

Heavy metals in soil (5g) was extracted with 1% EDTA as described by (8). Solutions were filtered through Whatman No. 1 filter paper and volume was made to 50ml by adding distilled water. Soil samples were digested in triplicates and analyzed for Fe, Mn, Zn, Co, Cd, Cr, Cu, Ni and Pb. For the determination of heavy metals the atomic absorption spectrophotometer was powered on and warmed up for 30 minutes. After the heating of cathode lamp, the air acetylene flame was ignited and instrument was calibrated or standardized with different working standards.

Sterilization of Apparatus

Petri plates, media bottles, distilled water, McCartney bottles and syringes were sterilized in autoclave. For sterilization purpose all apparatus were autoclaved for 40 minutes at 121°C. After autoclaving all sterilized material dried in oven at 95°C.

Media Preparation and Fungal Isolation

Potato Dextrose Agar (PDA) media was used for the isolation of fungi. Fungi were isolated on Potato Dextrose Agar (PDA) by soil dilution method (33).

Preparation of plates

Fungi were isolated on Potato Dextrose Agar (PDA) by soil dilution method (33). Poured the media in Petri-dishes and allowed to solidify for 24 hour. To suppress the bacterial growth, 30 mg/lit of streptomycin was added in the medium (33). Once the agar was solidified, and then put plates in an inverted position for 24 hours at room temperature (33).

Identification of fungi

The fungal cultures were identified on the basis of macroscopic (colonial morphology, color, texture, shape, diameter and appearance of colony) and microscopic (septation in mycelium, presence of specific reproductive structures, shape and structure of conidia, and presence of sterile mycelium) characteristics (41). Pure cultures of fungi isolates were identified with the help of literature (17,11).

RESULTS AND DISCUSSIONS

Table 1: Soil chemical analysis prior to planting

Chemical Properties	Soil	Abattoir Effluent	Poultry dropping
Nitrogen%	0.07	1.06	0.71
Available P	0.046	24.52	11.92
Exchangeable Ca (Cmol/kg)	2.79	11.58	33.87
Exchangeable Mg (Cmol/kg)	0.14	2.61	17.85
Exchangeable K (Cmol/kg)	0.02	4.55	16.93
Exchangeable Na (Cmol/kg)	0.11	0.11	0.14
Organic carbon%	0.5		
Organic Matter %	0.87		
pH in H ₂ O	9.3		
pH in KCl	8.9		
Acidity	0.4		
Sand	75.96		
Silt	11.28		
Clay	12.76		
Textural Class	Sand		
Cation exchange capacity	3.06		
Chromium mg/kg	25.3		
Cadmium mg/kg	0.5		
Nickel mg/kg	2.5		
Lead mg/kg	9.1		
Manganese mg/kg	350.5		
Iron mg/kg	25,250		
Copper mg/kg	9.65		
Zinc mg/kg	54.7		
Cobalt mg/kg	1.9		

The table 1 below shows the initial soil chemical properties of Asa River sediment, in which the sediments were observed to be sandy, organic matter content of the sediment was ranked low at (0.0.87%), while the nitrogen content of the sediment also ranked as low (0.07). The pH of the sediment showed 8.9 in KCl which means that the sediment was alkaline in nature and available phosphorus was also seen to be extremely low (0.046). Calcium was observed to be low, magnesium and potassium were also observed to be extremely very low and sodium also observed to have low value (0.11). Cation exchange capacity of the sediment was observed to fall below the range considered to be low. The chemical analysis of the sediment showed that the soil is low in terms of fertility. The heavy metal analysis of the sediment revealed that all metals fall below maximum permissible limit as set by USEPA except manganese (350.5mg/kg). Table 1 also revealed that abattoir effluent has a high nitrogen content of (1.06) compared to poultry dropping of 0.71%. Table 1 further revealed that abattoir effluent had a higher phosphorus content (24.52%) than poultry dropping (11.92%), potassium, calcium magnesium and sodium were observed to be higher in poultry dropping than abattoir effluent.

146

Table 2: Effect of organic waste on heavy metal availability in soil after cropping

Treat./Levels	Co (mg/kg)	Mn (mg/kg)	Cr (mg/kg)	Cd (mg/kg)	Pb (mg/kg)	Zn (mg/kg)	Fe (mg/kg)	Cu (mg/kg)	Ni (mg/kg)
Control	0.30	58.1	0.04	0.00	1.8	0.45	4.23	0.47	0.10
AE 1.6t/ha	0.18	46.9	0.05	0.01	1.53	0.57	3.00	0.48	0.067
AE 3.2 t/ha	0.24	44.0	0.04	0.01	1.47	0.59	2.37	0.51	0.03
AE 1.6t/ha+NPK120KgN	0.34	57.9	0.04	0.003	1.5	0.51	4.33	0.45	0.10
AE 3.2 t/ha +NPK120KgN	0.31	52.2	0.12	0.003	1.7	0.52	4.20	0.38	0.00
Control	0.30	58.1	0.04	0.00	1.8	0.45	4.23	0.47	0.10
PD10 t/ha	0.24	55.6	0.04	0.01	1.5	0.82	2.60	0.55	0.067
PD15 t/ha	0.31	47.0	0.07	0.003	1.77	0.86	3.73	0.62	0.067
PD10 t/ha+NPK120KgN	0.33	48.0	0.05	0.003	1.37	0.81	2.93	0.50	0.067
PD15 t/ha+NPK120KgN	0.24	40.5	0.04	0.00	1.37	0.78	4.13	0.42	0.067
SED	0.003	5.32	0.027	0.003	0.246	0.099	0.507	0.07	0.037
LSD(0.05)	ns	Ns	Ns	ns	Ns	ns	*	ns	ns

147

148 Table 2 shows that there are no significant differences at <0.05 between the treatments at various levels of application except for Fe with the
149 highest reduction in concentration observed at sole application rate of 3.3t/ha. Although they were no significant differences between the
150 treatment applied yet it was observed that the organic waste were very effective in reducing the concentration of heavy metals that were
151 available in the soil. This result is supported by the studies of (42, 43) who were both of the opinion that organic waste possesses large surface
152 area, which implies a high capacity for complex heavy metals on their surface. Impact of organic amendments in bioremediation is effective and
153 their effectiveness is related to the quantity applied (44).

154

Occurrence of fungi isolates in poultry droppings

Figure 1 shows the percentage of occurrence of the fungi in poultry dropping amended plots. Ten (10) fungal species were isolated. It shows that *Aspergillus niger* had the highest of occurrence (19.1%), followed by *Aspergillus flavus* (14.6%), *Trichoderma viride* (14%), *Trichoderma harzianum* (12.3%), *Fusarium solani* (12.1%), *Penicillium notatum* (10.3), *Trichoderma verrusocum* (6.4%), *Fusarium oxysporum* (5.3%), *Microsperum nanum* (3.8%) and *Aspergillus ustus* (2.1%).




Figure 1: Percentage occurrence of fungi in poultry droppings

Occurrence of fungi isolates in abattoir effluent

Fig. 2 shows the percentage of occurrence of fungi in abattoir effluent. Eleven (11) fungal species were isolated. *Aspergillus niger* had the highest percentage of occurrence (19.6%), followed by *Trichoderma viride* (15.1%), *Fusarium solani* (11.2%), *Penicillium notatum* (10.1%), *Aspergillus flavus* (9.7), *Trichoderma harzianus* (7.8%), *Trichophyton verrusocum* (6.5%), *Fusarium oxysporum* (6.2%), *Stachybotrys chartarum* (5.4%), *Aspergillus ustus* (4.6%) while *Microsporium nanum* had the lowest occurrence (3.8%).




Figure 2: Percentage occurrence of fungi in abattoir effluent

Result above shows that different fungi of Ascomycotina were isolated and identified in the amended plots and these fungi showed excellent bioremediation capability for different heavy metals such as Co, Cd, Cu, Ni, Pb and Zn. These agrees with the findings of (12) who evaluated the efficiency of different fungi and found out that *Aspergillus niger* and *A. flavus* has high capacity in remediating Pb, Zn, Cr and Cd from refinery effluent. Furthermore, (5) in their study of Mycoremediation: utilization of fungi for reclamation of heavy metals at their optimum remediation conditions found out that *Aspergillus niger*, *A. flavus*, *A. fumigates* and *A. ustus* has great potential in reducing heavy metals concentration in contaminated environment. This finding is similar to that of (1,32). Two *Fusarium sp.* were identified in which *Fusarium solani* was known to degrade Cr, Zn and Nickel (36), while *F. oxysporum* was also identified to be effective in reducing the concentration of Cd, Cu, Pb, and Zn. This is similar to the findings of (14) who discovered the *F. oxysporum* helps to biosorb Zn and Cu and *F. solani* removes Cu (2). Similarly, two species of *Trichoderma* were identified; *T. harzanium* and *T. viride*. *T. viride* was identified to have a high accumulating potential for Ni, Cd and Cr (27), and (22) found out that *T. viride* has high uptake potential for chromium. While *T. harzanium* has been observed to have a high reducing capacity for Siddique *et al.*, 2011, 35). Moreso, *P. notatum* was found to have capacity for reducing high chromium concentrations (37). Result further revealed that *Stachybotrys chartarum*, plays an important role in reducing copper (Cu) in contaminated soil. In a study conducted by (16) to evaluate the tolerance capacity of *Stachybotrys chartarum* in a copper based preservative, they found that the fungi has high tolerance level for copper. Furthermore, *Microsporum nanum* was found to play an important role Cd, Cr, Pb and this was supported by the findings of (12) who opined that *M. nanum* reduced the concentration of heavy metal from a refinery effluent.

It was also observed from Figure 1 and 2 that *Aspergillus niger* is the dominant isolated specie from the two amended plots. *Aspergillus niger* has been known to be able to tolerate the heavy metals in organic wastes and the result obtained in this study confirm the earlier assertion by different authors who asserted that *Aspergillus sp.* is the dominant species in wastewater (25).

CONCLUSION

From the result of this study, it was deduced that organic waste are effective material for bioremediation, and their effectiveness was enhanced as a result of the activities of certain fungi which performs important functions coupled with their inherent potential to tolerate and reduce heavy metals. Fungi isolated and identified in abattoir and poultry amended plots includes *Aspergillus niger*, *Trichoderma viride*, *Fusarium solani*, *Penicillium notatum*, *Aspergillus flavus*, *Trichoderma harzianum*, *Trichophyton verrucosum*, *Fusarium oxysporum*, *Stachybotrys chartarum*, *Aspergillus ustus* and *Microsporum nanum*. *Aspergillus niger* (19% in poultry droppings and 19.6% in abattoir effluent) was observed to have the highest population in all the amended plots.

REFERENCES

1. Ahmad, I., Ansari, I. and Aqil, F. (2005a). Biosorption of Ni, Cr and Cd by metal tolerant *Aspergillus niger* and *Penicillium sp.* using single and multi-metal solution. Indian J Exp. Biol. 44: 73-76.
2. Amatussalam, A., Abubacker, M.N. and Rajendran, R.B. 2011. In situ Carica papaya stem matrix and *Fusarium oxysporum* (NCBT-156) mediated bioremediation of chromium. Indian J Exp. Biol. 49: 925-31.
3. , P., Isar, J., Saran, S., and Saxena, R.K. (2006). Bioaccumulation of copper by *Trichoderma viride*. Bioresour. Technol. 97:1018-1021.
4. Anderson, J.M. and Ingram, J.S.I. (1993). Tropical Soil Biology and Fertility: A Handbook of Methods. Second edition. CAB International, Wallingford, UK. p.37
5. Archana and Jaitly, A. K. (2015). Mycoremediation: utilization of fungi for reclamation of heavy metals at their optimum remediation conditions. Journal of Biology and Life Sciences. 3(1): pp. 77-106.
6. Aung, M.M. and ing, Y.P., (2005). Bioleaching of spent fluid catalytic cracking catalyst using *Aspergillus niger*, Journal of Biotechnology, 116, 159-170.
7. AOAC International (formerly the Association of Official Analytical Chemists), 1999. *Official Methods of Analysis*. Arlington, VA: AOAC International.
8. Association of Official Analytical Chemist, (1980). *Official Methods of Analysis*. 13th edition. AOAC, Washington, DC.
9. Baldrian, P., (2003). Interactions of heavy metals with white-rot fungi. Enzyme Microb. Technol. 32:78-83.
10. Bates R.G, (1954). *Electromeric pH determination*. John Willey and sons Inc., New York, USA., Pp 87-92.
11. Barnett H.L., Hunter, B.B. (1999). Illustrated genera of imperfect fungi, fourth edition. Prentice Hall Inc.
12. Bello, O.A., Abubakar, B.Y. and Abdullahi I.O. (2016). Efficiency of *Aspergillus niger*, *Aspergillus flavus* and *Microsporum nanum* to remov heavy metal from refinery effluent. Journal of Advances in Biology and Biotechnology, 6(3): pp. 1-6.
13. Bray, R.H. and Kurtz, L.T. (1945), Determination of total, organic, and available forms of phosphorus in soils. *Soil Science*, 59: 39-45.

14. Chandrakar, V., Verma, P. and Jamaluddin (2012). Removal of cu and zn by fungi in municipal sewage water. *Int. J. Adv. Biotechnol. Res.* 2: 787-790.
15. Concas, A., and Cao G., (2004) Technologies for the Remediation of Contaminated Sites (Italian Language), CUEC, 1-85.
16. Danguolė, B. and Loreta L. (2007). Fungal tolerance towards copper-based wood preservatives. *BIOLOGIJA*. 2007. Vol. 53. No. 4. P. 54–61.
17. Domsch KH, GamsW, Anderson TH. 1980. *Compendium of soil fungi*. London, England: Academic Press.
18. Gavrilesca M (2004). Removal of heavy metals from the environment by biosorption. *Eng. Life Sci.* 4(3):219-232.
19. Hafez, N., Abdel-Razek, A.S., Hafez, M.B., (1997) Accumulation of some heavy metals on *Aspergillus flavus*. *J Chem Technol Biotechnol* 68:19–22.
20. Hemambika, B., Johncy R.M., and Rajesh V.K., (2011). Biosorption of heavy metals by immobilized and dead fungal cells: A comparative assessment. *Journal of Ecology and the Natural Environment* Vol. 3(5), pp. 168-175.
21. Iskandar, N.L., Zainudin NAIM, Tan, S.G. (2011). Tolerance and biosorption of copper (Cu) and lead (Pb) by filamentous fungi isolated from a freshwater ecosystem. *Journal of Environmental Science* 23(5):824-830.
22. Joshi, P.K., Swarup, A., Maheshwari, S., Kumar, R. and Singh, N. 2011. Bioremediation of heavy metals in liquid media through fungi isolated from contaminated sources. *Indian J. microbial.* 51: 482-487.
23. Kadirvelu K, Senthilkumar P, Thamaraiselvi K, Subburam V (2002). Activated carbon prepared from biomass as adsorbent: Elimination of Ni(II) from aqueous solution. *Bioresour. Technol.*, 81: 87-90.
24. Kapoor, A., and Viraraghavan, T., (1997) Heavy metal biosorption sites in *Aspergillus niger*. *Biores Technol* 61:221–227
25. Sharma Kavita (2010). Isolation of soil mycoflora of katar near Gangtok, India. *Journal of Phytology* 2010, 2(5): 30–32.
26. , S., Q. Cao, Y.M. Zheng, Y.Z. Huang and Y.G. Zhu. (2008): Health risks of heavy metals in contaminated soils and food crops irrigated with wastewater in Beijing, China. *Environ.Pollut.*, 152: 686-692.
27. Levinskaite, L. 2001. Simultaneous effect of Ni, Cd and Cr on soil micromycetes. *Biologija*. 13-15.
28. Luef, E., Prey, T. and Kubicek, C.P., (1991). Biosorption of Zinc by fungal mycelia waste. *Appl.Microbiol. Biotechnol.*, 34:688-692.
29. Mann, H., (1990). Biosorption of heavy metals by bacterial biomass. In: Volesky B (ed) *Biosorption of heavy metals*. CRC Press, Boca Raton, FL, pp 93-137.
30. Maurya, N.S., Mittal, A.K. Cornel, P., Rother, E. (2006). Biosorption of dyes using dead macro fungi: Effect of ye structure, ionic strength and pH, *Bioresour. Technol.* 97: 512-521.
31. Mala, J.G.S, Nair, B.U., Puanakrishnan, R. (2006). Accumulation and biosorption of chromium by *Aspergillus niger* MTCC 2594. *J. Gen. Appl. Microbiol.* 52:179-186.
32. Ramasamy, R.K., Congeevaram, S. and Thamaraiselvi, K. 2011. Evaluation of isolated fungal strain from e-waste recycling facility for effective sorption of toxic heavy metal pb (II) ions and fungal protein molecular characterization- a mycoremediation approach. *Asian J. Exp. Biol. Sci.* 2: 342-247.
33. Razak AA, Bachman G, Farrag R. 1999. Activities of microflora in soils of upper and lower Egypt. *African Journal of Mycology and Biotechnology* 7, 1-19.

34. Rhoades, J.D. (1982). Cation exchange capacity. In: A.L. Page (ed.) *Methods of soil analysis*. Part 2: Chemical and microbiological properties (2nd ed.) Agronomy 9:149-157.
35. Sarkar, S., Satheshkumar, A., Jayanthi, R. and Premkumar, R. 2010. Biosorption of Nickel by live biomass of *Trichoderma harzianum*. *Res. J. Agr. Sci.* 1: 69-74.
36. Sen, M. (2013). Biosorption of zinc and nickel from wastewaters using nonliving cells of *Fusarium solani*. *Int. J. Chem. Tech. Appl.* 2: 63-70.
37. Seshikala, D. and Charya, M.A.S. 2012. Effect of pH on Chromium biosorption. *Int. J. Pharma. Bio.Sci.* 2: 298-302.
38. Siddiquee, S., Salleh, N.A., Sujjat, A., Saili, N.S. and Naher, L. 2013. Tolerance and biosorption capacity of Zn^{2+} , Pb^{2+} , Ni^{3+} and Cu^{2+} by filamentous fungi (*Trichoderma harzianum*, *T. aureoviride* and *T. virens*). *Adv. Biosci. Biotechnol.* 4: 570-583.
39. Turnau, K., Orlowska, E., Ryszka, P., Zubek, S., Anielska, T., Gawronski, S., Jurkiewicz, A., (2006). Role of mycorrhizal fungi in phytoremediation and toxicity monitoring of heavy metal rich industrial wastes in southern Poland. *Soil Water Pollut. Monit. Prot. Remed.* 3(23):533-551.
40. Walkley, A., and T.A. Black. (1934). An examination of the Degtjarett method for determining soil organic matter and a proposed modification of the chromic acid titration method. *Journal of Soil Science*, 37:29-38
41. Zafar, S., Aqil, F. and Ahmad, I. (2007). Metal tolerance and biosorption potential of filamentous fungi isolated from metal contaminated agriculture soil. *Bioresource Technology* 98, 2557-2561.
42. Uchimiya, M., Lima, I. M., Klasson, K. T., and Wartelle, L. H., 2010: Contaminant immobilization and nutrient release by biochar soil amendment: *Roles of natural organic matter*, *Chemosphere*, 80, 935–940.
43. Park, J. H., Choppala, G. H., Bolan, N. S., Chung, J. W., and Chuasavathi, T. (2011): Biochar reduces the bioavailability and phytotoxicity of heavy metals, *Plant Soil*, 348, 439–451.
44. Walker, D. J., R. Clemente, A. Roig and M. P. Bernal, 2003. The effects of soil amendments on heavy metal bioavailability in two contaminated Mediterranean soils. *Environ. Pollut.*, 122: 303–312.
45. Rovina K, Azad SA, Naher L, Suryani S, Siddiquee, S . (2015). Heavy Metal Contaminants Removal from Wastewater Using the Potential Filamentous Fungi Biomass. *Journal of Microbial & Biochemical Technology Journal of Microbial & Bioch. Technol* 7: 384-393.