

# Comparative Evaluation of bioactive compounds in some raw and boiled egg varieties: Eggs, potential nutraceuticals?

## ABSTRACT

**Aims:** To comparatively evaluate the bioactive compounds (egg white proteins) of chicken (exotic and local), turkey, quail and guinea fowl eggs. This study also aimed at ascertaining the claims on egg (particularly the raw form) being a functional food.

**Study design:** Experimental.

**Place and Duration of Study:** Department of Biochemistry, University of Calabar, Calabar, and Department of Pharmacology, University of Nigeria, Nsukka, February to July 2017.

**Methodology:** Five species of bird eggs studied in the raw and boiled forms are: exotic chicken, local chicken, turkey, quail and guinea fowl eggs. Freshly-laid eggs were purchased from poultries, cleaned and then some were boiled before homogenizing and analyses. A combination of methods involving egg white preparation, separation of egg white proteins using ion-exchange chromatography, purification using tangential flow filtration and quantification using the colorimetric Bradford assay, was used. Results of the quantitative estimation of avidin, lysozyme, ovalbumin, ovotransferrin and flavoprotein concentrations were statistically compared using analysis of variance (ANOVA).

**Results:** It was generally observed that the concentrations of the bioactive compounds (except ovalbumin) were significantly ( $P < 0.05$ ) higher among the raw eggs than the boiled ones. Raw turkey egg had the highest concentration of avidin ( $15.83 \pm 0.15 \mu\text{g/g}$ ) and this was significantly different ( $P < 0.05$ ) from the others, while quail had the lowest concentration ( $8.47 \pm 0.20 \mu\text{g/g}$ ) even among the boiled samples. Ovalbumin, a storage protein, was found to be the most abundant of the egg white proteins (average of 50-55%).

**Conclusion:** Some egg varieties such as quail egg, have relatively safer content of avidin; others such as raw turkey egg with quite higher avidin concentrations, should be consumed in moderation in order to reduce the risk of biotin deficiency. The presence of these bioactive compounds in significant quantities also show that eggs may serve as functional foods.

**Keywords:** bioactive compounds, eggs, nutraceuticals, proteins

## 1. INTRODUCTION

Nutrition Food is any material of plant or animal origin that consists of essential body nutrients such as carbohydrates, fats, proteins, minerals or vitamins, and is ingested or assimilated by an organism to produce energy, sustain growth and maintain [1]. The nutrient composition of various foods depend on several factors which include species, breeds, cultivars, ecological factors, post-harvest handling, preservation and storage techniques. Some foods are considered healthy depending on their nutrient content while others are considered unhealthy [2].

The importance of good and adequate nutrition for maintenance of health can never be overemphasized. Poverty, diseases, harmful economic systems, food insecurity, inadequate food and agricultural policies, poor nutrition education, adverse climatic changes, conflicts/wars are some of the causes of hunger and malnutrition [3]. With the increase in malnutrition and poverty globally, it has become necessary to develop strategies and remedies to the growing number of nutrition-related diseases that populations are being faced with. There are food based approaches and agriculture based approaches employed in tackling the problem of malnutrition. Food-based approaches, particularly utilizing animal source foods (such as meat and eggs) offer potentially sustainable solutions to multiple deficiencies [4].

Research has shown that food biodiversity can provide some sustainable solution for combating food insecurity and malnutrition [4]. Consumption of different breeds/varieties may have significant impact on nutrition and health outcomes. Some specific data are required to promote the use of biodiversity for food and nutrition. These information which include food composition and consumption from various varieties/breeds and their dietary contribution to nutrition and human health, have been increasing as well as awareness. Different varieties/breeds vary significantly in their nutrient compositions [5]. Macronutrients from different varieties of the same species could vary by 10-fold, and micronutrients by up to 1000-fold [6] by virtue of the genetic resource itself.

Bird eggs which are of different varieties, many of which are consumed by man for food, are made up of the yolk, albumen and shell. The albumen (egg white) consist primarily of water (87%) and protein (13%) with no cholesterol and little, if any fat [7]. According to Kovacs (2005), ovalbumin (54%), ovotransferrin (12%), ovomucoid (11%), lysozyme (3.5%) and ovomucin (3.5%) are the major egg white proteins while avidin (0.05%), cystatin (0.05%), ovoflavoprotein (0.8%), ovomacroglobulin (0.5%), glycoprotein (1%) and ovoidinhibitor (1.5%) are the minor ones. Each of these proteins are recognized for their functional importance and they possess many functional properties such as gelation, emulsification and coagulation [8].

According to Li-Chan *et al.* (1995) [9], many egg white proteins have also been found to possess various antimicrobial and antioxidant properties. Technologies have now been developed for separating egg white proteins commercially, hence there are possibilities of using these egg white proteins for their antimicrobial and antioxidant properties [10]. Ovalbumin is the major egg white protein synthesized in the hen's oviduct and accounts for 54% of the total egg white proteins [11]. Conalbumin was renamed as ovotransferrin after findings that it can bind iron. One molecule of ovotransferrin can bind 2 molecules of iron, and it transports iron in the body [8]. Lysozyme is another important egg white protein found in nature as a monomer, but is occasionally present as a dimer with more thermal stability. It is considered as a strong basic protein present [12]. Ovomucin, one of the major egg white proteins, has more coiled regions at its extremities, like the structure of human mucin; there are the soluble and insoluble forms. Previous studies have shown that at least 3 types of carbohydrate chains are found in ovomucin, which are composed of galactose, galactosamine, sialic acid, and sulfate with a molecular ratio of 1:1:1:1. On average, 33% of ovomucin is carbohydrates [13]. Ovomuroid is one of the most highly glycosylated proteins found in egg white [14]. A research [15] reported that ovomucoid can be used to control *Streptomyces erythraeus*. Ovomucin is also considered as a trypsin inhibitor, which is a negative property of the protein; it has the capability to control microorganisms and so can be used as an antimicrobial agent in foods.

Separation of egg white proteins have been done for many years but new, simple, economical, and sequential methods with better yield and purity are emerging. Among the egg white proteins, lysozyme is currently used as antimicrobial agent in the food industry, and others proteins such as ovalbumin has a strong potential as a drug carrier, ovotransferrin as an antimicrobial agent or iron carrier, ovomucin and ovomucoid as antimicrobial and immunomodulating agents. Peptides derived from ovotransferrin, ovalbumin, ovomucoid, and ovomucin showed cytotoxic, anticancer, immunomodulating, ACE-inhibitory, antimicrobial, and antioxidant activities, and have high potentials to be used in the pharmaceutical, nutraceutical, and food industries [8].

Numerous species of bird eggs exist which are nutritious but there seems to be scanty information on the nutritional and non-nutritional content of different egg varieties, except the popular chicken egg. According to Kiple (2007) [16], the chicken egg is the most consumed by humans but other eggs including those of quail, guinea fowl, goose, turkey and duck are also important in human nutrition; however, information on egg quality characteristics have been quite limited to chicken eggs and comparative evaluation of egg varieties are not many.

This study therefore seeks to evaluate and compare the concentrations of five bioactive compounds (egg white proteins) in some popularly consumed bird egg varieties namely exotic chicken, local chicken, turkey, quail and guinea fowl eggs, in their raw and boiled forms.

## 2. MATERIALS AND METHODS

### 2.1 Egg sample collection and preparation

84 Eggs were sourced from poultries in Nsukka, Enugu state; only freshly laid eggs were purchased for  
85 the purpose of this research. The different egg species were cleaned and prepared separately for  
86 analyses.

87 **2.1.1 Preparation of raw egg samples:** Shells of fresh eggs were cleaned, broken and the  
88 contents emptied into clean glass beakers. The raw egg content was then homogenized and placed in  
89 clean, labelled beakers and sealed with parafilm.  
90

91 **2.1.2 Preparation of hard-boiled egg samples:** The fresh eggs were boiled by adding them to tap  
92 water already boiling at 100°C. The tap water was put to cover the eggs in the pot. The eggs were left  
93 to boil for 10mins, immediately after which they were removed and allowed to cool in tap water at  
94 room temperature. Each type of eggs was boiled separately and then shelled after they had cooled.  
95 The boiled eggs were placed in clean, labelled beakers and sealed with parafilm.  
96

97 After homogenising, 20g of each of the samples (raw and boiled samples) were stored in properly  
98 labelled, air-tight sample glass bottles until ready for analyses. All the reagents used in the laboratory  
99 analyses were of standard analytical grade (AR).  
100

## 101 **2.2 Determination of bioactive compounds**

102 This was carried out according to the method of He (2011) [17] and Bradford (1976) [18]. The  
103 bioactive compounds (egg white proteins) were analysed using a combination of methods involving  
104 egg white preparation, separation of egg white proteins using ion-exchange chromatography,  
105 purification using tangential flow filtration and quantification using the colorimetric Bradford assay.

106 **2.2.1 Principle and Procedure:** Ion-exchange chromatography separates ions based on their  
107 charged groups. A change in pH changes the charge on the particular molecules and therefore alters  
108 binding. The molecules then start eluting based on the changes in their charges from adjustments. In  
109 the Bradford assay, the separated proteins form a protein-dye complex with the Bradford reagent and  
110 the concentration of the protein is colorimetrically determined using a spectrophotometer.

111 **2.2.1.1 Egg White Preparation:** Fresh eggs were broken, and egg white and yolk were separated  
112 using a kitchen egg separator. The egg white was diluted 10-fold with 10mM Tris-HCl and 10mM  
113 disodium orthophosphate. This was stirred for 30 minutes and stored overnight at 4°C. The diluted  
114 sample was then centrifuged at 10000g for 30min to remove all insoluble debris, and the supernatant  
115 was collected. Supernatant was prepared for ion exchange chromatography.

116 **2.2.1.2 Ion Exchange Chromatography** (Liu *et al.*, 2012) [19]: The separation of egg white proteins  
117 was carried out using tandem ion-exchange chromatography. This method incorporates both HiLoad  
118 26/10 High Performance Q and SP Sepharose Fast Flow columns for the separation of both anionic  
119 and cationic species, on an ÄKTA explorer FPLC System (GE Healthcare). Columns were washed  
120 with elution buffer (10mM Tris-HCl, 10mM DSOP, 1M NaCl, pH 7). Further equilibration with 4 column  
121 volumes of equilibration buffer (10mM Tris-HCl, 10mM DSOP, pH 7). 100mL of diluted egg white  
122 (1:10 equilibration buffer) was loaded at a flow rate of 8.0mL/min initially onto the Q Sepharose  
123 column to bind anionic compounds. The unbound cationic components were then trapped onto the SP  
124 Sepharose column. Compounds that did not bind either Q or SP Sepharose were collected as the  
125 flow through. Egg white proteins were first eluted from the Q Sepharose column with elution buffer  
126 with isocratic steps at 5% for 30 minutes, 10% for 30 minutes, 20% for 20 minutes, and 100% for 20  
127 minutes. Afterwards, the in-line valve was switched to isolate the SP Sepharose column. Elution was  
128 then achieved with elution buffer on the HiLoad SP Sepharose column using isocratic steps at 5% for  
129 20 minutes and 100% for 20 minutes.

130 **2.2.1.3 Purification of fractions:** Tangential Flow Filtration (TFF) and Stirred Cell Filtration were used.  
131 Tandem ion-exchange chromatography separated diluted egg white into nine fractions. Fractions  
132 were concentrated on two 1 kDa nominal molecular weight limit (NMWL) via TFF using a ProFlux M12  
133 (Millipore, Bedford, MA, USA) at a trans membrane pressure of 40 psi. Ultrafiltration and buffer  
134 exchange (diafiltration) were used to ensure conductivity of the sample was approximately  $\leq 300 \mu\text{S}$ .

Owing to the large hold-up volumes associated with TFF, an Amicon stirred cell with a 1 kDa NMWL regenerated cellulose ultracel membrane (millipore) was used to further concentrate protein fractions and filter the samples in PBS. Nitrogen gas was applied at a maximum pressure of 75 psi while the sample was stirred at 25 rpm to avoid “gelling” and accumulation of egg white proteins on the filter membrane. Concentration of each protein fraction was quantified using the colorimetric Bradford assay. The purified sample fractions were stored at  $-40^{\circ}\text{C}$ .

### 3. RESULTS AND DISCUSSION

#### 3.1 Bioactive compounds

Table 1 shows the results of the quantitative estimation of bioactive compounds (egg white proteins). Comparing the raw and boiled samples, it was generally observed that the bioactive compounds concentrations were significantly higher ( $P < 0.05$ ) in the raw egg samples than the boiled samples; for instance, the concentration of avidin in raw quail egg was  $8.47 \pm 0.20 \mu\text{g/g}$  but it was  $4.60 \pm 0.10 \mu\text{g/g}$  in the boiled sample. This was the trend except for ovalbumin which had slightly higher concentrations in the boiled egg samples. Among the raw samples, turkey egg had the highest concentration of avidin ( $15.83 \pm 0.15 \mu\text{g/g}$ ) and this was significantly ( $P < 0.05$ ) different from the others, while quail had significantly ( $P < 0.05$ ) lower concentration ( $8.47 \pm 0.20 \mu\text{g/g}$ ) even in the boiled samples. Ovalbumin was of greater percentage composition with values ranging between 47 and 53 g/100g egg white in both raw and boiled samples. The avidin content of quail egg reduced by half after boiling. It was also observed that for lysozyme, all the raw and boiled concentrations varied significantly ( $P < 0.05$ ) except for guinea fowl egg where boiling did not significantly affect the lysozyme content. Ovotransferrin concentrations of raw exotic chicken, quail and guinea fowl eggs, were statistically similar ( $P > 0.05$ ) and higher than that of local chicken and turkey egg which were also statistically similar ( $P > 0.05$ ).

The concentrations of bioactive compounds found to be higher in the raw eggs than in the boiled ones, may help to explain why raw eggs are prescribed for use as functional foods for ‘treating’ conditions such as hypertension, infections and hyperlipidemia because these bioactive compounds have been reported to have antihypertensive, hypolipidemic, anticancer and antibacterial properties [20]. In another research, Verrinder-Gibbins (1997) [21] indicates that there are potential approaches for increasing egg white proteins through genetic engineering. This will afford possibilities for enhancing functional properties and increasing their use for industrial and pharmaceutical applications. The high content of avidin in raw turkey egg causes it not to be recommended for individuals who need increased levels of biotin (vitamin  $\text{B}_7$ ). This is because avidin binds biotin and reduces its bioavailability to the body. In an article [22], it was reported that biotin improves glycemic control in diabetic patients and hence, biotin supplementation is one of the recommended treatments for diabetes since the vitamin helps in lowering blood sugar and also functions in energy production. Quail egg had the least avidin concentrations (both in the raw and boiled samples) and may be a healthier choice for diabetic patients who choose drink raw eggs for perceived health benefits. Ovalbumin had the highest concentration and this agreed with the findings of Stevens (1996) [23] and Zabik (1992) [24] who reported ovalbumin as the major egg white protein. During storage, ovalbumin is changed to s-ovalbumin, an extra heat-stable form [25]. This may be the reason for the observed non-decrease of its concentration in the boiled sample. Ovotransferrin makes up about 13% of the egg white proteins and its ability to bind to iron is related to its antimicrobial activity [24]. According to Akkouche *et al.* (2012) [26], when egg white is heated, its globular proteins are prone to changes in structure and conformation. Depending on the temperature and duration of heating, these changes can range from denaturation, to gelation or coagulation. This may help explain the decrease in concentration of some of the bioactive compounds in the boiled samples. Flavoprotein is reported to have the highest selenium content (a potent antioxidant) of 1800 ng/g, among the egg white proteins [27]. This may be responsible for some of the speculated anticancer activities of raw eggs. Of all the species, raw turkey eggs were found to be richest in flavoprotein.

189 **Table 1. Bioactive compounds (egg white proteins) in raw and boiled egg varieties.**

EGG SPECIE	Avidin (µg/g EW)	Lysozyme (g/100g EW)	Ovalbumin (g/100g EW)	Ovotransferrin (g/100g EW)	Flavoprotein (g/100g EW)
<b>Raw samples</b>					
Exotic chicken	9.37 ± 0.18 <sup>b</sup>	4.47 ± 0.07 <sup>b</sup>	49.90 ± 0.64 <sup>b</sup>	1.50 ± 0.12 <sup>a</sup>	0.25 ± 0.00 <sup>c</sup>
Local chicken	9.77 ± 0.07 <sup>b</sup>	4.23 ± 0.09 <sup>b</sup>	52.37 ± 0.18 <sup>c</sup>	1.70 ± 0.06 <sup>b</sup>	0.26 ± 0.00 <sup>c</sup>
Turkey	15.83 ± 0.15 <sup>c</sup>	5.30 ± 0.12 <sup>c</sup>	52.17 ± 0.74 <sup>c</sup>	1.17 ± 0.03 <sup>b</sup>	0.31 ± 0.01 <sup>d</sup>
Quail	8.47 ± 0.20 <sup>a</sup>	3.63 ± 0.09 <sup>a</sup>	47.63 ± 0.32 <sup>a</sup>	1.73 ± 0.03 <sup>b</sup>	0.16 ± 0.01 <sup>a</sup>
Guinea fowl	9.40 ± 0.17 <sup>b</sup>	4.50 ± 0.12 <sup>b</sup>	51.20 ± 0.35 <sup>c</sup>	1.40 ± 0.10 <sup>a</sup>	0.22 ± 0.00 <sup>b</sup>
<b>Boiled samples</b>					
Exotic chicken	7.17 ± 0.07 <sup>c</sup>	2.37 ± 0.15 <sup>a</sup>	53.37 ± 1.05 <sup>b</sup>	1.27 ± 0.07 <sup>c</sup>	0.21 ± 0.01 <sup>b</sup>
Local chicken	6.30 ± 0.12 <sup>b</sup>	3.30 ± 0.12 <sup>b</sup>	52.67 ± 0.81 <sup>b</sup>	1.20 ± 0.06 <sup>b</sup>	0.21 ± 0.01 <sup>b</sup>
Turkey	11.40 ± 0.17 <sup>d</sup>	4.17 ± 0.03 <sup>c</sup>	53.47 ± 0.19 <sup>b</sup>	1.17 ± 0.03 <sup>b</sup>	0.25 ± 0.01 <sup>b</sup>
Quail	4.60 ± 0.10 <sup>a</sup>	3.23 ± 0.07 <sup>b</sup>	42.53 ± 1.22 <sup>a</sup>	1.27 ± 0.03 <sup>c</sup>	0.83 ± 0.03 <sup>c</sup>
Guinea fowl	6.77 ± 0.29 <sup>b</sup>	4.17 ± 0.07 <sup>c</sup>	53.60 ± 0.21 <sup>b</sup>	1.07 ± 0.03 <sup>a</sup>	0.13 ± 0.03 <sup>a</sup>

190 Values are expressed as mean ± SEM, n = 3. Values with different superscripts in the same column are significantly ( $P < 0.05$ )  
 191 different from each other. EW means egg white.  
 192

#### 193 4. CONCLUSION

194  
 195 Comparative evaluation of the bioactive compounds of some bird eggs has shown that some egg  
 196 varieties have relatively higher content of some egg white proteins than others. These bioactive  
 197 compounds have also been reported to have antimicrobial and antioxidant activities, thus causing the  
 198 eggs to have the ability to serve as functional foods. In order to reduce the risk of biotin deficiency,  
 199 consumption of varieties such as quail egg (with lower avidin levels) should be more recommended  
 200 especially for diabetics who need more biotin due to its hypoglycaemic activity.

#### 201 REFERENCES

- 202  
 203 1. [www.merriam-websterunabridged.com](http://www.merriam-websterunabridged.com)  
 204

- 205 2. Ezzati, M., Hoorn, S. V. & Rodgers, A. (2002). Comparative Risk Assessment  
206 collaborating Group Estimates of global and regional potential health gains from reducing  
207 multiple major health risk factors. *The Lancet*, 362 (9380): 271-80.
- 208 3. FAO, IFAD & WFP (2014). The state of food insecurity in the world. Food and Agriculture  
209 Organisation of the United Nations, Rome.
- 210 4. FAO (2013). Combating Micronutrient Deficiencies: Food-based Approaches. The Food and  
211 Agriculture Organization of the United Nations. Eds B. Thompson and L. Amoroso).
- 212 5. Burlingame, B., Charrondiere, R. & Mouille, B. (2009). Food composition is fundamental to the  
213 cross-cutting initiative on biodiversity for food and nutrition, *Journal of Food Composition and*  
214 *Analysis*, 22: 361–365.
- 215 6. Stadlmayr, B., Nilsson, E., Mouille, B., Medhammar, E., Burlingame, B., Charrondiere, U. R. (2011).  
216 Nutrition indicator for biodiversity on food composition – A report on the progress of data availability.  
217 *Journal of Food Composition and Analysis*, 24(4-5): 692-698.
- 218 7. Pamplona-Roger, D. (2004). Encyclopedia of Foods and Their healing Power (Volume 1).  
219 *Education and Health Library*. Review and Herald Publishing.
- 220 8. Abeyrathne, E. D., Lee, H. Y. & Ahn, D. U. (2013). Egg white proteins and their potential use in  
221 food processing or as nutraceutical and pharmaceutical agents- A review. *Poultry Science*, 92: 3292-  
222 3299.
- 223 9. Li-Chan, E. C. Y., Powrie, W. D. & Nakai, S. (1995). The chemistry of eggs and egg products. In  
224 Egg Science and Technology. Edited by W. J. Stadelman and O. J. Cotterill. The Haworth Press Inc.  
225 New York.
- 226 10. Abdel-Aal, E. M., Akhtar, H., Zaheer, K. & Ali, R. (2013). Dietary sources of lutein and zeaxanthin  
227 carotenoids and their Role in Eye health. *Nutrients*, 5(4): 1169-1185.
- 228 11. Stadelman, W. J. & Cotterill, O. J. (2001). Egg Science and Technology. 4th Edition. Avi Publ.  
229 Co., Westport, CT.
- 230 12. Huopalahti, R., Fandino, R. L., Anton, M. & Schade, R. (2007). Bioactive Egg Compounds.  
231 Springer, New York, NY.
- 232 13. Mine, Y. (2008). Egg Bioscience and Biotechnology. John Wiley & Sons Inc., Hoboken, New  
233 Jersey.
- 234 14. Kovacs-Nolan, J. K. N., Zhang, J. W., Hayakawa, S. & Mine, Y. (2000). Immunochemical and  
235 structural analysis of pepsin-digested egg white ovomucoid. *Journal of Agriculture and Food*  
236 *Chemistry*, 48: 6261–6266.
- 237 15. Nagata, K. & Yoshida, N. (1984). Interaction between Trypsin-like enzyme from *Streptomyces*  
238 *erythraeus* and chicken ovomucoid. *Journal of Biochemistry*, 96: 1041–1049.
- 239 16. Kiple, K. F. (2007). A Moveable Feast: Ten Millenia of Food Globalization.  
240
- 241 17. He, F. (2011). Bradford Protein Assay. *Bio-protocol*, Bio101: e45. DOI: 10.21769/BioProtoc.45.
- 242 18. Bradford, M. M. (1976). A rapid and sensitive method for the quantitation of microgram quantities  
243 of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry*, 72(1-2): 248-254.
- 244 19. Liu, Y. J., Du, Q., Yang, B. C., Zhang, F. F., Chu, C. H., Liang, X. M. (2012). Silica based click  
245 amino acid stationary phase for ion Chromatography and Hydrophilic Interaction Liquid  
246 Chromatography. *Analyst*, 137: 1624-1628.  
247



- 248 20. Froning, G. W. (1998). Recent Advances in Egg Products Research and Development: A paper  
249 presented at the University of California Egg Processing workshop. Riverside and Modesto.
- 250 21. Verrinder-Gibbins, A. M. (1997). Genetically-Engineered Poultry. International Egg  
251 Commission Annual Production and Marketing Conference. Toronto, Ontario.
- 252 22. Gaby, A. R. (2009). Chromium and Biotin for Type 2 Diabetes – Literature Review and  
253 Commentary. *Townsend Letter*, May edition.
- 254 23. Stevens, L. (1996). Egg proteins: what are their functions? *Science Progress*, 79: 65-87.
- 255 24. Zabik, M. (1992). Eggs and Products. In: BOWERS *Journal of Food Theory and Applications*. 2<sup>nd</sup>  
256 edition. New York: Macmillan Publishing Co. pp. 359-424.
- 257 25. Donovan, J. W. & Mapes, C. J. (1976). A Differential Scanning Calorimetric study of conversion of  
258 ovalbumin to s-ovalbumin. *Journal of the Science of Food and Agriculture*, 27: 197-204.
- 259 26. Akkouche, Z., Aissat, L. & Madani, K. (2012). Effect of Heat on egg White Proteins. A Paper  
260 presented at International Conference on Applied Life Sciences (ICALS). © INTECH pp. 407-413.
- 261 27. Jacobs, K., Shen, L., Benemariya, H. & Deelstra, H. (1993). Selenium distributions in egg  
262 white proteins. *Z Lebensm Unters Forch*, 196(3): 236-238.