

Original Research Article

Effect of traditional and improved cassava processing on Cassava Derivative Products in Central African Republic

ABSTRACT

Cassava (*Manihot esculenta* Crantz) is a starchy staple food that previous researches have showed to contain cyanogenic compounds, precursors of hydrocyanic acid, undoubtedly toxic for humans. With the aim to determine food security in cassava, the aim of this study is to compare the effect of traditional and improved cassava processing in Cassava Derived products. Data were analysed by unpaired student's t-test at a confidence level of 95% ($P < 0.05$). The concentration of cyanogenic compounds in cossette of Cassava from improved processing (shucking + fermentation + drying) obtained after retting of cassava root without skin is significantly lower than the levels in traditional processing.

Keywords: Cassava, Derivative Products, Technological Treatment

Introduction

Cultivated cassava is usually *Manihot esculenta* Crantz. Since its domestication thousands of years ago in the Amazon region, cassava is now spread around the world and is widely cultivated for consumption in the tropics and sub-tropic (Siritunga, 2002; Tewe and Lutaladio, 2004; Burns *and al.*, 2010; FAOSTAT, 2012). Cassava is a valuable source of food for developing countries and Central African Republic (CAR) is no an exception (Conaway *and al.*, 2012). Cassava is a nutritionally strategic famine crop could support food security in areas of low rainfall. Mature roots are able to survive for a long time without water and still retain nutritional value. Roots are a valuable source of calories, whereas cassava leaves are a valuable source of protein, minerals, and vitamins (Aboua and Kamenan, 1996; Sahoré Drogba and NemlinGnopo, Wobeto *and al.*, 2006; Chijindu and Boateng, 2008).

Cassava is grown in CAR as a subsistence crop. Its utilization as food varies from region to region. It is a source of food security, not only because it can be grown on less productive land, but because it is a source of income for urban and rural populations (Kosh-Komba and *al.*, 2016). People in many parts of CAR rely largely on starchy foods for their carbohydrates intake. The population consume it in various forms: tuber without any preparation, chicwangué, porridge soup, roasted, braised (retted or unretted) in the form of cooked pasta prepared from flower made out of pieces of cassava. More than 80% of Central African populations' energetic needs come from cassava. However, the status of cassava as a food security crop to most subsistence farmers is threatened by pests, diseases and weeds and inadequate processing techniques to eliminate the cyanogenic glycoside in cassava (Pohlandt *and al.*, 1983; Ernesto *and al.*, 2002; Diallo *and al.*, 2014).

Usually, cassava is well processed before being consumed. Inadequate processing however may result in appreciable amounts of cyanogenic glycosides remaining and this may pose a public health risk (WHO, 2004; ATSDR, 2006; FSANZ, 2008; Mburu *and al.*, 2012; Abraham *and al.* 2016).

The aim of this study is to assess cyanide content in various processed cassava to determine how different processing techniques can help reduce cyanide content and to provide information to relevant stakeholders on cyanide contents in cassava derived products.

2. Materials and Methods

2.1. Choice of Surveyed Sites

In response to the invitation to participate in the research project, samples were received from Yaloké, Sibut and Pissa (figure 1). The sites were chosen to represent a range of environments and management practices in cassava-based cropping systems in the mid-altitude zone of CAR. The climate in all sites is sub-humid with a bimodal rainfall distribution. This allows for the production of most annual crops during both the long (March–August) and the short rains (September–October). Altitude ranges between 1200 and 1500 masl. Cassava is planted in the first 2 months of the short or long rains and remains in the field for about a year. Agricultural systems are diverse with farmers growing 4–6 main crops on average (Boulvert, 1986).




Figure 1. Location of sites for data collection in Central African Republic

2.2 Sample Collection

A questionnaire was prepared and used as a tool for the collection of information in the different sites. The questionnaire focused on the different processing to eliminate the cyanogenic glycoside in cassava. 250- 350 households per site were randomly selected. Structured interviews, in combination with a visit of different processing techniques (traditional and improved processing) to eliminate the cyanogenic glycoside in cassava were documented. Cassava Derivative Products (Locally called cossette of cassava and bread of war) were obtained from farms market vendors from Yaloké, Sibut and Pissa.

2.3 Humidity level

5 g of each sample is putting in an oven at 85 °C until their mass is stabilized. The mass obtained after this operation, subtracted from the initial mass (5g), give the water content (AOAC 1975) according to the form $\% H = [(M1-M2)/M1] \times 100$. M1 = initial mass (5g), M2 = stabilized mass and % H= Humidity level.

2.4 Glucide dosage

The reducing sugars are assayed according to the MILLER method (1959). At elevated temperature and in basic medium, 3,5-dichlorosilicic acid is reduced to 3-amino-5-nitrosalicylic acid in the presence of reducing sugars. The product of the reaction is yellow-orange, and has a maximum absorption at 575 nm. This method makes it possible to determine concentrations of reducing sugars ranging from 0 to 1 g/l. To 1 ml of the suitably diluted sample was added 1 ml of the Dinitrosilicic acid reagent. The mixture is boiled for 5 minutes and cooled rapidly by immersing it in an ice bath. After the addition of 8 ml of distilled water, the contents of the tubes vortex-homogenized and the optical density read at 575

nm. A glucose standard range from 0 to 1 g / l carried out in parallel for each series of measurements.

2.5 Lipids dosage

The lipids were determined on soxhlet (AOAC 1975) by extraction with hexane for 6 hours

2.6 Cyanogenic glycoside dosage

2.6.1 Extraction of cyanogenic compounds

Samples of 10 g from Cassava Derivative Products (cossette of cassava) were homogenized in 30 ml of acid extraction medium. The amount of sample varied, but the ratio of sample to extraction medium was always approximately 1:3. The homogenized samples were left to stand for 10 min and then centrifuged at 10,000 g for 10 min (Barthet and Bacala, 2010). The supernatant was stored at 4°C until assayed for cyanogenic compounds.

2.6.2 Cyanogenic glycoside determination in cassava derivative products

The cyanide concentration in cossette of cassava was determined using ninhydrin based spectrometer of trace cyanide at 485 nm maximum wavelength (Ikédiobi *and al.*, 1980; Surleva *and al.*, 2013; Lambri *and al.*, 2013). A calibration graph was first constructed using standard solutions of CN⁻ at concentrations of 0.02, 0.04, 0.08, 0.1 and 0.2 µg/mL (which is within the linear range) and was prepared by adding appropriate volumes of cyanide solutions at concentration of 20 µg CN⁻/mL to 1 mL of 2% Na₂CO₃. Ninhydrin solution (0.5 mL) containing 5 mg/mL in 2% NaOH was added to each standard cyanide solution. The mixture was homogenized and incubated for 15 minutes for color development. Similarly, the blank was prepared in the same way as above, except that instead of 1 mL 2% Na₂CO₃ containing CN⁻, 1 mL of 2% Na₂CO₃ without CN⁻ was added. UV-Visible absorption of the reaction product (Cyanide-ninhydrin adduct) of the different concentrations of cyanide was measured using UV/Vis Spectrophotometer at 485 nm. Total cyanide in the samples was determined by adding 0.1 g of the ground sample in a standard volumetric flask (5 mL) and made up to mark with 0.1% NaHCO₃. The samples were sonicated for 20 minutes in a water bath and the mixture centrifuged at 10,000 rpm for 10 minutes. The supernatant was pipetted with automatic pipette, two aliquots (2 mL each) and added to 0.5 mL ninhydrin in NaOH, allowed for fifteen minutes for color development and absorbance measured at 485 nm.

2.7 Data analysis

Data were analysed by unpaired student's t-test at a confidence level of 95% (P<0.05) to find the significant difference between various parameters. XLSTAT statistical software (version 2008) was used for data analysis. Excel (version 2003) was used for showing the correlation between the traditional cassava processing and improved cassava processing.

3. Results

Cassava derivate products were obtained using traditional and improved processes to closely reflect the way that people consume these products. In this study, the selected products were Cossette of cassava and bread of war.

3.1. Traditional cassava processing to obtain cossette of cassava and bread of war

Traditional processing techniques for cassava are described in Figures 2, 4 and 6. Most of the prospective households do the retting of unpeeled cassava roots in the backwaters with a retting period ranging from 3 to 5 days according to the seasons. The drying time varies from 10 to 20 hours depending on the dry and rainy seasons.




Figure 2. Traditional cassava processing to obtain cossette of cassava and bread of war

3.2. Improved cassava processing to obtain cossette of cassava and bread of war

Improved cassava processing techniques are described in **Figures 3, 5 and 7**. The retting process in Plastic Drums ranging in length from 24 to 36 hours is done with peeled roots. Drying time varies from 1 to 2 hours depending on dry and rainy seasons.




Figure 3. Improved cassava processing to obtain cossette of cassava and bread of war



Figure 4 : Traditional retting



Figure 5 : Improved retting

199

200



Figure 6 : Traditional drying



Figure 7 : Improved drying



Figure 8 : Cassava Derivative Products
Locally called Cossette of cassava



Figure 9 : Cassava Derivative Products
Locally called bread of war

201

202 3.3. Result of carbohydrate determination

203 The Glucide profile of Cassava in traditional and improved processing is depicted in Table 1.

204 The difference observed is significant ($P < 0,0001$). This proves that there is variability between the
205 measured parameters according to the two cassava processing.

206

207

208 **Table 1: Glucide contain in Cossette of Cassava from traditional and improved processing**

Difference	91,869
t (observed value)	385,378
t (critical value)	-1,990
Degree of freedom	79
p-value	< 0,0001
alpha	0,05

209

210

211

The correlation test also shows that the two measured parameters through the two techniques are not related (figure 10). Thus, there is a weak negative correlation ($R^2 = -0,068$).




Figure 10: Glucose contain in Cossette of Cassava from traditional and improved processing

3.4. Result of Lipids determination

The lipids profile of Cassava in traditional and improved processing is depicted in Table 2. The difference observed is significant ($P < 0,0001$) this proves that there is variability between the measured parameters according to the two cassava processing.

Table 2: Lipid contain in Cossette of Cassava from traditional and improved processing

Difference	0,147
t (observed value)	20,575
t (critical value)	-1,990
Degree of freedom	79
p-value	< 0,0001
alpha	0,05

The correlation test also shows that the two measured parameters through the two techniques are not related (figure 11). Thus, there is a weak negative correlation ($R^2 = -0,856$).




Figure 11: Lipid contain in Cossette of Cassava from traditional and improved processing

3.5. Result of Cyanogenic glycoside

The Cyanogenic glycoside profile of Cassava in traditional and improved processing is depicted in Table 3.

The difference observed is significant ($P < 0,0001$). This proves that there is variability between the measured parameters according to the two cassava processing.

Table 3: Cyanogenic glycoside contain in Cossette of Cassava from traditional and improved processing

Difference	12,184
t (observed value)	102,567
t (critical value)	-1,990
Degree of freedom	79
p-value	< 0,0001
alpha	0,05

The correlation test also shows that the two measured parameters through the two techniques are not related (figure 12). Thus, there is a weak negative correlation ($R^2 = -2,090$).




Figure 12: Cyanure contain in Cossette of Cassava from traditional and improved processing

3.4. Humidity level in cossette of cassava and bread of war

The Humidity profile of Cassava derived products is depicted in Table 4.

The difference observed is significant ($P < 0,0001$). This proves that there is variability between the measured parameters according to the two cassava processing.

Table 4: Humidity level in cossette of cassava and bread of war

Difference	253 49,128
t (observed value)	254 63,956
t (critical value)	255 -12,960
Degree of freedom	79
p-value	< 0,0001
alpha	0,05

The correlation test also shows that the two measured parameters through the two techniques are not related (figure 13). Thus, there is a weak negative correlation ($R^2 = -0,70$).




Figure 13: Humidity level in cossette of cassava and bread of war

4. Discussion

Cassava roots are rich in energy and contain mainly starch and soluble carbohydrates (Abraham and al., 2016). Although they are low in protein, it is a staple food for about 200 to 300 million people worldwide (Chijindu and Boateng, 2008). Cassava is consumed in various forms. Even though cassava is an important food, it contains toxic and antinutritional substances that interfere with digestion and uptake of nutrients (Wobeto and others 2007). Cyanide is the most toxic factor restricting the consumption of cassava roots and leaves. Cassava contains cyanogenic glucosides that are toxic for humans and can lead to serious health disorders.

In this study, the concentration of cyanogenic compounds is higher in traditional cassava processing. On the other hand, the concentration of cyanogenic compounds in improved cassava processing is significantly lower than the levels in traditional processing ($P < 0.05$) (figure 12). World Health Organization recommends 10 mg/kg body weight as the maximum safe cyanide level (Okafor and al., 2002; WHO, 2004) for fresh cassava intake. The concentration of cyanogenic compounds in cossette of Cassava from improved processing (shucking + fermentation + drying) obtained after retting of cassava root without skin is significantly lower than the levels in traditional processing. The presence of skin in traditional processing could be in the origin of higher of cyanogenic compounds. Cyanide content ranges from 53 to 1300 mg cyanide (HCN) equivalents/kg DW (Siritunga and Sayre 2003; Wobeto and others 2007) in leaves, and from 10 to 500 mg HCN equivalents/kg DW (Arguedas and Cooke 1982; Dufour 1988; Siritunga and Sayre 2003, Cardoso and al., 2005) in root parenchyma.

Cassava contains cyanogenic glycosides, mostly linamarin (2- β -D-glucopyranosyloxy-2-methylpropanenitrile) and lutostralin (2- β -D-glucopyranosyloxy-2-methylbutyronitrile) (ATSDR, 2010; Jorgensen *and al.*, 2011). These glycosides are supposed not to be toxic, but their decomposition into hydrogen cyanide (HCN) by intestinal flora enzymes would take place if humans ingested them (Sahoré and NemlinGnopo, 2010). The produced HCN is toxic for humans and it was reported to be responsible for certain pathological disorders such as Konzo diseases, thyroid goiter and tropical ataxic neuropathy (Ernesto *and al.*, 2002). The (per)acute oral toxicity of cyanide ions and HCN in mammals is well known. The small molecule is readily absorbed and distributed via systemic circulation. Above a certain tissue level, cyanide inhibits the cytochrome c oxidase (complex IV, the terminal enzyme of the mitochondrial electron transport chain by competitively binding to the oxygen-reducing cofactor of the protein (Piero and al., 2015). This crucial effect causes decreased utilization of oxygen and increased anaerobic metabolism, leading to excess of lactic acid and metabolic acidosis, and finally to cell death through energy deprivation. Due to its high dependence on oxidative metabolism, the central nervous system is particularly vulnerable to cyanide intoxication. With high oral doses, symptoms occur within a few minutes and may include nausea, vomiting, giddiness, headache, palpitations, hyperpnoea then dyspnoea, bradycardia, unconsciousness, and violent convulsions, followed by death (JECFA 2012; WHO 2004). In terms of the minimal lethal dose of cyanide in humans, the number of about 0.5 mg/kg body weight is commonly cited. It dates back to a paper by Gettler and Baine (1938) who applied a formula for back-calculation of the dose derived from experimental data in dogs to tissue levels found in humans after lethal intoxications. Often, a range of 0.5–3.5 mg/kg body weight is cited in the literature for the acute lethal dose of cyanide in humans which first was published by Halstrøm and Møller (1945). However, in some cases, much higher doses have been survived (e.g. 2.5 g potassium cyanide equivalent to 1000 mg cyanide: Tassan et al. 1990).

The carbohydrate content varies between 90% and 95% in the sample cossettes. Studies by other authors have revealed the content in some cassava varieties of 94% (Sahoré Drogba and NemlinGnopo, 2010).

Fermentation of cassava pulp or dough for 4-5 days results in a decrease in total cyanogens of 52-63% (Kendirim *and al.*, 1995; Obilie *and al.*, 2004).

5. Conclusion

The concentration of cyanogenic compounds in cossette of Cassava from improved processing (shucking + fermentation + drying) obtained after retting of cassava root without skin is significantly lower than the levels in traditional processing. This achievement is a strategic step in cassava detoxification.

References

- Diallo Y., Gueye M.T., Ndiaye C., Sakho M., Kane A., Barthelemy J.P. and Lognay G. A New Method for the Determination of Cyanide Ions and Their Quantification in Some Senegalese Cassava Varieties. *American Journal of Analytical Chemistry*. 2014; 5: 181-187.
- Pohlandt C., Jones E. A. and Lee A. F, "A Critical Evaluation of Methods Applicable to the Determination of Cyanides," *Journal of the South African Institute of Mining and Metallurgy*, Vol. 83, 1983, pp. 11-19.
- Ernesto M., Cardoso A. P., Nicala D., Mirione E., Massaza F., Cliff J., Haque M. R. and Bradbury J. H. 2002. "Persistent Konzo and Cyanide Toxicity from Cassava in Northern Mozambique," *Acta Tropica*, Vol. 82, No. 3, pp. 357-362. [http://dx.doi.org/10.1016/S0001-706X\(02\)00042-6](http://dx.doi.org/10.1016/S0001-706X(02)00042-6).
- Sahoré Drogba A. and NemlinGnopo J. 2010. Effect of Technological Treatments on Cassava (*ManihotEsculentaCrantz*) Composition.*Food and Nutrition Sciences*, 2010, 1, 19-23.
- FAOSTAT. (2012). Fiche produits manioc. Food and Agriculture Organization of the United Nations, Rome, Italy. 6P.
- Ikédiobi,C.O., Onyia,G.O.C., Eluwa C.E.,(1980) A rapid and inexpensive assay for total Cyanid in Cassava (*Mnihot esculenta Crantz*) and cassava Products, *Agr.biol.chem.*, 44,2803-2809.
- Barthet VJ and Bacala R (2010) Development of optimized extraction methodology for cyanogenic glycosides from flaxseed (*Linum usitatissimum*). *J AOAC Int* 93:478–484.

- Jorgensen K, Morant AV, Morant M, Jensen NB, Olsen CE, Kannangara R, Motawia MS, Møller BL, Bak S (2011) Biosynthesis of the cyanogenic glycosides linamarin and lotaustralin in cassava: isolation, biochemical characterization, and expression pattern of CYP71E7, the oxime-metabolizing cytochrome P450 enzyme. *Plant Physiol* 155:282–292.
- JECFA, Joint FAO/WHO Expert Committee on Food Additives (2012) Cyanogenic glycosides. In: WHO food additives series no. 65, safety evaluation of certain food additives and contaminants.
- WHO, World Health Organization (2004) Hydrogen cyanide and cyanides: human health aspects. Concise international chemical assessment document 61, Geneva. <http://www.who.int/ipcs/publications/cicad/en/cicad61.pdf>. (Accessed 08 Feb 2015).
- Abraham K. · Buhrke T. · and Lampen A. Bioavailability of cyanide after consumption of a single meal of foods containing high levels of cyanogenic glycosides: a crossover study in humans. *Arch Toxicol* (2016) 90:559–574. DOI 10.1007/s00204-015-1479-8.
- Kosh-Komba E., Aba-Toumno L., Zinga I., Yandia P., Atato A., Kadekoy-Tigague D., Wabolou F., Kongbo Dembo E., Batawila K., Akpagana K. The Social and Economical Factors Influence the Use of Cassava for Improving Livelihood in Central African Republic. *Resources and Environment* 2016, 6(2): 41-45 DOI: 10.5923/j.re.20160602.03.
- ATSDR (2006). Toxicological Profiles for Cyanide. Agency for Toxic Substances and Disease Registry, Atlanta, GA: U.S department of Health and Human Services, Public Health Services. <http://www.atsdr.cdc.gov/ToxProfiles/tp.asp?id=72&tid=19>.
- Piero NM, Joan MN, Richard OO, Jalemba MA, Omwoyo OR, et al. (2015). Determination of Cyanogenic Compounds Content in Transgenic Acyanogenic Kenyan Cassava (*Manihot esculenta* Crantz) Genotypes: Linking Molecular Analysis to Biochemical Analysis. *J Anal Bioanal Tech* 6: 264 doi:10.4172/2155-9872.1000264.
- Egan SV, Yeoh HH, Bradbury JH (1998) Simple picrate paper kit for determination of the cyanogenic potential of cassava flour. *J Sci Food Agric* 76: 39-48.
- Cumbana A, Mirione E, Cliff J, Bradbury JH (2007) Reduction of cyanide content of cassava flour in Mozambique by the wetting Method. *Food Chem* 101: 894-897.
- Mburu FW, Swaleh S and Njue W (2012) Potential toxic levels of cyanide in cassava (*Manihot esculenta* Crantz) grown in Kenya. *African Journal of Food Science* 6: 416-420.
- Food and Agriculture Organization/World Health Organization Food standards programme (1991) Codex Alimentarius Commission X11. Supplement 4. FAO/ UN, Rome, Italy.
- Cumbana A, Mirione E, Cliff J, Bradbury JH (2007). Reduction of cyanide content of cassava flour in Mozambique by the wetting Method. *Food Chem* 101: 894-897.
- Mburu FW, Swaleh S, Njue W (2012) Potential toxic levels of cyanide in cassava (*Manihot esculenta* Crantz) grown in Kenya. *African Journal of Food Science* 6: 416-420.
- Siritunga SD (2002). Generation of acyanogenic cassava (*Manihot esculenta* Crantz): transgenic approaches. The Ohio State University, USA.
- Chijindu, E.N. and Boateng, B. (2008) Effect of Nutritional Content of Processed Cassava Chips on Development of *Prostephanus truncates* (Horn). *World Journal of Agricultural Sciences*, 4, 404-408.
- Cardoso, A.P., Mirione, E., Ernesto, M. and Massaza, F. (2005) Processing of Cassava Roots to Remove Cyanogens. *Journal of Food Composition and Analysis*, 18, 451-460. <http://dx.doi.org/10.1016/j.jfca.2004.04.002>.
- Lambri, M., Fumi, M.D., Roda, A. and De Faveri, D.M. (2013) Improved Processing Methods to Reduce the Total Cyanide Content of Cassava Roots from Burundi. *African Journal of Biotechnology*, 12, 1285-1291.
- Surleva, A., Zaharia, M., Ion, L., Gradinaru, R.V., Drochioiu, G. and Mangalagiu, I. (2013) Ninhydrin-Based Spectrophotometric Assays of Trace Cyanide. *Acta Chemica IASI*, 21, 57-70. <http://dx.doi.org/10.2478/achi-2013-0006>.
- Tchacondo, T., Karou, S.D., Osseyi, E., Agban, A., Bawa, L.M., Atcha, A.R., Soumana, K., Assignon, K., Kpemoua, A. and De Souza, C. (2011) Effect of Harvest and Cook Processing on Cyanides Content of Cassava Cultivars and Cassava-Based Dough Consumed in Lomé, Togo. *Advanced Journal of Food Science and Technology*, 3, 398-402.
- Wobeto C, Corrêa AD, de Abreu CMP, dos Santos CD, de Abreu JR. 2006. Nutrients in the cassava (*Manihot esculenta* Crantz) leaf meal at three ages of the plant. *Ciência e tecnologia de alimentos*, Campinas 26:865–9. Available from: <http://www.scielo.br/pdf/cta/v26n4/23.pdf>.
- Tewe OO and Lutaladio N. 2004. Cassava for livestock feed in sub-Saharan Africa. Rome, Italy: FAO.

394 Burns, A., Gleadow, R., Cliff, J., Zacarias, A., Cavagnaro, T. 2010. Cassava: The drought, war and
395 Famine crop in a changing world. *Sustainability* 2, 3572-3607.

396 Cardoso, A.P., Mirione, E., Ernesto, M., Massaza, F., Cliff, J., Haque, M.R., Bradbury, J.H. 2005.
397 Processing of cassava roots to remove cyanogens. *J. Food. Comp. Anal.* 18, 451-460.

398 Boulvert Y., (1986). Carte Phytogéographique de la République Centrafricaine (feuille Ouest- feuille
399 Est) à 1:1000000. ORSTOM, 31 p.

400 Conaway J.L., Ouedraogo A.K. et Coneff J. (2012). *Activité de zonage plus de moyens d'existence de*
401 *la République centrafricaine*. USAID (United States Agency International Development),
402 Bangui, Centrafrique 41p.

403 SahoréDrogba A. and NemlinGnopo J. 2010. Effect of Technological Treatments on Cassava
404 (*ManihotEsculentaCrantz*) Composition. *Food and Nutrition Sciences*, 2010, 1, 19-23.

405 Okafor PN, Okorowkwo CO, Maduagur EN. 2002. Occupational dietary exposure of humans to
406 cyanide poisoning from large scale Cassava processing and ingestion of Cassava foods. *Food*
407 *Chem Toxicol.* 40:001–5.

408 Agbor-Egbe T, Lape Mbome I. (2006) The effects of processing techniques in reducing cyanogen
409 levels during the production of some Cameroonian cassava foods. *Journal of Food*
410 *Composition and Analysis*; 19: 354-363.

411 Kendirim OC, Chukwu OA, Achinewhu SC. (1995) Effect of traditional processing of cassava on the
412 cyanide content of gari and cassava flour. *Plant Foods for Human Nutrition*; 48(4): 335-339.

413 Obilie EM, Tano-Debrah K, Amoa-Awua WK. (2004) Souring and breakdown of cyanogenic glucosides
414 during the processing of cassava into akyeke. *International Journal of Food Microbiology*; 93:
415 115-121.