

Determination of PSP toxins in Moroccan shellfish by MBA, HPLC and RBA methods: Links to causative phytoplankton *Alexandrium minutum*

ABSTRACT

Paralytic shellfish toxins (PST) or paralytic shellfish poisoning (PSP) toxins are secondary metabolites of toxic species of phytoplankton. The consumption of shellfish accumulating these toxins can cause neurological symptoms and even death. Within the framework of the surveillance program of seafood safety along the Moroccan littoral environment established by National Institute of Fisheries Research (INRH), a study of PST was conducted from 2004 to 2016 in south Moroccan's shellfish, mussels from south Agadir region and solen from Dakhla bay. The surveillance was carried out bi-monthly or weekly using the AOAC official method of analysis (AOAC 959.08) mouse bioassay (MBA). In parallel, monitoring of toxic phytoplankton in water was conducted. In the aim to determine the shellfish toxin profile, ion pair high-performance liquid chromatography with post column derivatization and fluorescence detection (HPLC-FD) was performed. The Receptor Binding Assay (RBA) also was used for determination of total toxicity of PSP toxins in Agadir's mussels. In both regions, the analysis of sea water revealed the presence of the toxic algae *Alexandrium* spp during toxics events. Along the coast of Agadir, PSP toxins in shellfish were associated with the presence of *Alexandrium cf. minutum* in seawater. These toxic events were widely distributed in time and space and mainly detected during the summer and fall seasons. In some samples concentrations exceeded the sanitary threshold (ST) of 800 $\mu\text{g eq STX /kg}$. HPLC analysis revealed that the toxin profile was dominated by Saxitoxin and Gonyautoxins. The comparison between different methods showed a strong uphill (positive) linear relationship, with a coefficient correlation of $r=0.79$ between MBA and HPLC and $r = 0.809$ between MBA and RBA.

Keywords

Paralytic shellfish poisoning, *Alexandrium minutum*, Morocco, HPLC, RBA, MBA.

1. INTRODUCTION

PSP toxins form a family of more than 20 molecules, chemically close to each other, all derived from Saxitoxin by substitution of various radicals of this molecule, the chemical structures are illustrated in fig. 1. PSTs are accumulated by shellfish filtering on algae producing toxins, mainly by dinoflagellates belonging to the genus *Alexandrium*, which may occur in the temperate and tropical climate regions [1], [2].

PSP toxins were detected repeatedly in the bivalves of Moroccan coasts and can represent a real danger for the consumers of shellfish especially of mussels collected from natural beds along atlantic and mediterranean coasts. The first case of intoxication by shellfish was recorded in 1961. The year 1994 was marked with many episodes of human intoxication [3], [4].

From 1998 to 2016, several episodes of shellfish toxicity were reported, involving Moroccan Mediterranean coasts [5], [6], [7] and in bivalves of Moroccan Atlantic coasts [8] but no case of PSP human intoxication were reported during this period. In the aim to protect the consumer of this seafood, a phycotoxin and toxic phytoplankton monitoring system was established early nineties by the National Institute for Fisheries Research (INRH) along Moroccan coasts.

The aim of this work is to present a spatio-temporal variation of toxic phytoplankton in waters and PSP toxin analysis in south Moroccan shellfish (in mussels from the south of Agadir and in solen from- Dakhla bay). Moreover, an investigation of PSP toxin profile is presented and three detection methods were compared, the AOAC official method of analysis by mouse bioassay (MBA), HPLC-FD and the AOAC official method by receptor binding assay (RBA). The results of different methods were compared by presenting the advantages and disadvantages of each test. This is the first report concerning the use of the nuclear RBA for detection of PSP toxins in Moroccan shellfish.




Figure 1: Chemical structures of PSP toxins

2. MATERIALS and METHODS

2.1. Sampling site

Two shellfish species were taken from two natural beds of south Atlantic Moroccan coasts, *Perna perna* from Agadir coast (Dar sfint site 29°40',404 N - 09°59',115 W) and *solen marginatus* from Dakhla bay (Pk25 site 23°54',746N - 15°46', 060W). The shellfish of these areas are collected by local populations for consumption and monitored by a national program of the safety of the marine environment. Mussels were collected weekly from January 2010 to December 2016, more than 362 samples were studied. 286 samples of solen were harvested from 2004 to 2010. Three kg of shellfish were taken at low tide, in parallel, 362 sea water samples were collected weekly, from the same sites, at high tide (± 2 hours of peak) with net (20 µm) and with a bottle (1 L). All samples were kept in a cooler box during the transport to the laboratory.




Figure 2. Sampling sites

2.2. Toxin analysis

2.2.1 MBA

The MBA was conducted under monitoring program started in the early nineties along Moroccan coasts. During this study, analysis were performed from 2004 to 2016 in shellfish of south of Morocco according to the AOAC method for paralytic shellfish poisons 959.08 [9] using white male mice weighing between 18 and 22 g. The mice were injected with 1 ml of extract obtained from fresh shellfish meat (100 g $\pm$ 1) extracted with 0.1 M HCl (100 ml $\pm$ 1), pH adjusted to 3 $\pm$ 0.1. The mice (3 mice per extract) were observed for 1 h, the mouse weight and mean survival time were used to calculate the toxicities expressed in μ g STX equi./kg by using a conversion factor (0.2). This factor was calculated by standardization of this method using a certified reference standard of Saxitoxin (STX) purchased from, National Research Council Canada (IMB, NRCC, Halifax, Nova Scotia, Canada). These analyzes were carried out at the marine biotoxins laboratory of the National Institute for marine Research (INRH), Agadir, Morocco.

2.2.2. HPLC-FD

PSP Toxin profiles were performed by HPLC/FD in samples of Dakhla and Agadir collected during PSTs events of 2006.

Analysis were based on the method of Thielert et al. [10] with modification described by Yu et al [11].

All solvents used were analytical grade, 4 ml acetic acid 0.03 M was used for 2 g of sample.

- Extraction of phycotoxins: the sample and the solvent were mixed in Vortex for 1 min and then homogenized during 10 min in an ultrasonic bath and finally, samples were centrifuged for 5 min at 140000 rpm. The supernatant was filtered by a 0.45 μ m diameter filter syringe to obtain the filtrate ready for analysis by HPLC.

- Hydrolysis: For the determination of C-toxins (N-sulfocarbamates) a hydrolysis of the filtrate was performed. To 150 μ l of the filtrate, 37 μ l of 1 M HCl were added and the mixture was mixed in a vortex for 30 seconds and heated for 15 minutes at 90 $^{\circ}$ C. The mixture was vortexed a second time

Formatted: Font: Italic

for 30 seconds. Finally the Extract was neutralized with 75 µl of sodium acetate 1 N and mixed for 30 seconds- to obtain the sample is ready for HPLC analysis.

- Chromatography was conducted according to conditions as described by Diener [12]. The injection volume was 10 µl, column used was a Phenomenex type, Luna 5 µm RP-C18, 250 mm x 4.6 mm. The mobile phases were as follows: eluent A: 6 mM octanesulfonic acid, 6 mM heptanesulfonic acid, 40 mM ammonium phosphate, The pH adjusted to 7 with H₃PO₄ and 0.75% tetrahydrofuran and eluent B: 7 mM octane sulfonic acid, 7 mM heptanesulfonic acid, 48 mM ammonium phosphate, The pH is adjusted to 7 with H₃PO₄ and with 1% tetrahydrofuran and 10% acetonitrile. The flow rate was 1ml/min.

- The temperature set for the post-column derivatization was 50 °C, a solution containing 10 mM periodic acid and 550 mM NH₃ solution with a flow rate of 0.3 ml / min was used, pH was reduced to 2-3 by 0.75 M Nitric acid 0.75M with a flow rate of 0.4 ml/-min. Detection wavelengths were set at 330 nm for excitation and 390 nm for emission. STX, neosaxitoxin (NEO), and GTX1, 2, 3, 4- standards were purchased from the National Research Council Canada (NRC-PSP- 1B). The standards solutions were diluted with acetic acid 0.03 N as required.

2.2.3. Receptor Binding Assay RBA

The extracts of Agadir's samples taken during toxics events from 2010 to 2016 were analyzed following the procedure of the AOAC OMA 2011.27. Extracts obtained following the AOAC 959.08 method were filtered using a 0.45 filter paper and stored at -20 °C. These analyzes were conducted at the Environment Laboratories of the International Atomic Energy Agency in. The determination of PSP toxins by RBA was made according to the protocol of Van Dolah et al [13].

This method is based on the competition between a radioligand (STX-tritium) and the different dilutions of the samples on the same receptors. Thus, the amount of radioligand complexes decreases with increasing concentrations of toxin in the extract. The counting of the completed radioactivity is performed by liquid scintillation counter which gives a competition curve. This curve showing the percentage of the complex formed according to the toxins added, allows the determination of total amount of the toxin present in the sample.

2.2.4. Identification and phytoplankton counting

Observations were carried out using an inverted microscope, in the sedimentation tank 10 or 25 ml, according to Uthermol method [14]. The determination was made through appropriate systematic identification keys. The counting was performed with horizontal scanning. The results densities were expressed as number of cells per liter (cells * L⁻¹).

3. Results and discussion

3.1. MBA analysis and toxic species of phytoplankton investigation

The results of MBA analysis in Agadir's mussels show that five PST episodes were recorded during this study from 2010 to 2016. Toxins were detected in 6.62% of the samples (23 MBA positive on a total of 362 samples) (Fig 3). The longest period of mussels toxicity was recorded by this biological method in 2012 with 7 weeks against the shortest one was noted in 2013 (1 week). But, the highest concentration of this toxin was recorded in 2015 with 7068 µg eq STX-/kg.

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript




Figure 3: MBA positives samples in southern Agadir coasts 2010 to 2016.

The monitoring of toxic phytoplankton *Alexandrium* spp showed a significant spatial and temporal variability. Indeed, the species were present at low concentrations between February 2010 and September 2011 to increase in winter 2011. The maximum concentration reached 7.2×10^3 cells L^{-1} was recorded in the fall. *Alexandrium* remain in waters, with low densities that reach high values in late spring (May) - early summer (June) 2012 (2×10^3 cells L^{-1}), then they proliferate in July 2012 with a maximum of 4×10^3 cells L^{-1} ; then, no detection of this species until the spring 2014 with a maximum in June of 1.5×10^3 cells L^{-1} ; after this species remain present throughout the year in superficial waters, but at low concentrations ranging between 10^2 and 9×10^2 cells L^{-1} . In the spring of 2015, *Alexandrium* reached concentrations around 8×10^2 cells L^{-1} at July to mark a major efflorescence in August with 25×10^3 cells L^{-1} , bloom spread through the waters until the beginning of fall 2015 (since October). The concentrations of *Alexandrium* fell gradually to 7×10^2 and 40 cells L^{-1} , recorded respectively in spring (March) and summer (July) 2016.

The analysis of PSP toxins by MBA shows that many samples contain high levels of toxins compared to the regulatory limit (800 μg STX equi/kg), maximums noted were 845, 1145, 357, 2109, and 7068 μg STX equi./kg respectively in fall 2011, in summer 2012, in fall 2013, in the end of spring 2014 and in summer 2015 (Fig. 4). This toxicity is explained by the high concentrations of *Alexandrium* sp recorded in seawater during these episodes.

Comment [SCH1]: ??

Formatted: Not Superscript/ Subscript

Formatted: Not Superscript/ Subscript




Figure 4: Variation of *Alexandrium* spp concentrations in sea waters and PSP toxins in Agadir's mussels (Dar sfint area) (2010- 2016)

At the Dakhla bay (Pk25 area), the MBA analysis of PSP toxins in shellfish showed that one period was recorded from 2004 to 2010, where Solen was toxic for 5 weeks, toxins were found in 2.09 % of total analyzed samples (Fig 5). No case of PST was detected before winter 2006; in December 2006, the toxicity was exceeded sanitary threshold with a maximum of 1215 µg STX_{eq}/kg, in general, from October to November 2006 the concentrations of PSP toxins were near or below the sanitary threshold between 395 and 600 µg STX_{eq}/kg. From January 2007 to December 2016, PSP toxins were not recorded in Solen. Toxins were detected in fall, during this season the temperature was high in this region between 18 and 20 °C [15]. Toxic phytoplankton observation conducted from 2004 to 2006 shows that *Alexandrium* spp concentrations in seawater recorded a low spatial and temporal variability. Indeed, winter 2004 and spring 2006 showed an irregular presence of the species with low concentrations (ranging from 100 to 800 cells·L⁻¹). In May and June the concentrations exceeded the health threshold of *Alexandrium* (10³ cells·L⁻¹) with 2.5*10³ cells·L⁻¹, after the concentrations increased sharply in fall-winter 2006. The maximum value reached 49.4*10³ cells L⁻¹ registered in November 2006 (Fig. 5, 6). The detection of this toxic dinoflagellate coincided with the shellfish toxicity of this bay.




Figure 5: MBA positives samples in Dakhla bay coasts 2004 to 2006




Figure 6: Variation of *Alexandrium spp* concentrations in sea waters and PSP toxins in Solen from Dakhla Bay (Pk 25 area) (2004- 2006)


In the southern Moroccan Atlantic, PSP toxins were detected barely and with low concentrations, but in recent years these toxins became more frequent with very high concentrations. In general over the last decades the occurrence and intensity of HAB appears to be increasing on a global scale due to rising ocean temperatures and growing coastal eutrophication [16]. -The species

from the genus *Alexandrium*, detected during the episodes of shellfish toxicity from 2004 to 2016, have been known as producers for the neurotoxins of PSTs, which is one of the most dangerous marine toxins known [17]. This genus is most often associated with PSP toxins, as about 10 of the approximately 30 *Alexandrium* species are PSP producer [18]. The *Alexandrium* species was identified as *A. minutum* and confirmed in the seawater of the southern region of Agadir (Dar sfint) and pk25 (Dakhla bay) by scanning microscopymicroscopy and epifluorescence microscopy [19]. This species is primarily an inhabitant of environments with a high terrestrial influence, such as lagoons, bays and estuaries where nutrient levels are high, the water column is stratified, and disturbance low. In spite of this, *A. minutum* is able to grow in coastal areas where land-derived influences are low. An important factor in *Alexandrium*'s presence in coastal areas is the requirement for a shallow water depth, as cysts which provide the seed population needed to initiate a bloom must be exposed to light for germination. In this study, these conditions were more widely present in Dakhla bay than in Dar sfint area which could explain why *Alexandrium* never dominated algal community and were presented the very low density and increase only in the bloom period. Also, At Dar sfint waters, the net sample presented some temporary cysts of *Alexandrium* suspended in the water column. The annual and inter annual variation of *Alexandrium* bloom occurrence may be due to variations in the hydro-climatic conditions of coastal water in this region, where the water is warmer (17 to 22 °C). The warmer period in the south of Atlantic Moroccan coasts starts generally from June to September [20]. *Alexandrium minutum* have preferential for more temperate waters. Statistical studies on all bloom of *A. minutum* highlight the evidence that temperature is a major factor controlling this efflorescence.

In many countries MBA is used for many years as an official method for PSTs monitoring for protection of the public health against this type of intoxication [21]; [22]. The results obtained from both the *Alexandrium* cell counts and the MBA for PSTs analysis suggest a clear correlation between the high cell counts recorded and shellfish containing higher concentrations of PSTs. The shellfish PSP toxicity events recorded coincide with the typical "bloom season" for *Alexandrium* which generally occurred in summer and fall, but with high inter-annual variability. The same results was recorded by Guallar-Morillo [23] who reported that the period of appearance of *Alexandrium* was often between the end of spring early summer, its time niche and risk period of PSP toxins contamination.


HPLC-FD analysis

HPLC-FD analysis confirms the presence of PSP detected by MBA, the amounts detected did not exceed 511 µg STX.equi./kg. The PSP toxin profile of Dakhla's solen and Agadir's mussel samples showed the presence of Carbamate toxins (GTxs), and Saxitoxin (STX). The Main toxins that had been detected were GTX2, GTX3 and STX (figs.7, 8).



246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269

Figure.7 : PSP toxin profile of Agadir's mussels on june 2006



270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286

Figure 8: PSP by MBA and HPLC with **toxines** profile of Dakhla's solen during toxic event of end 2006

In Both Agadir coast and Dakhla bay, **carbamate** toxins were the main toxins detected with majority of GTXs and minority of STX. Similar results have been reported [5] in wedge shells taken from the north of Agadir with a profile dominated by GTXs. This toxin profile is very close to that of *Alexandrium minutum* collected from Cap Town in South Africa, this species had a toxin profile dominated by gonyautoxins [24]. Generally, the shellfish toxin profile is very close to that of ingested phytoplankton [25]. This dinoflagellate was detected in sea water during all toxic event in Agadir coast and Dakhla bay from 2004 to 2016 [19], [26], what leads us to implicate *Alexandrium minutum* in PSP toxicities of shellfish of both regions, however several studies reported that this toxic specie of dinoflagellate was implicated in PSTs events. [27], [28], [29].

3.2. Comparison between the different methods used for PSP toxin analysis

Figure 9 and figure 10 illustrate respectively the correlation between toxicities of HPLC and MBA and that between RBA and MBA. The first information obtained there is a strong positive correlation between the two different non-animal methods against the reference MBA, the comparative results are summarized in Table 1. Results yielded an r of 0.79, indicating that the MBA reports somewhat higher STX equivalents in shellfish, relative to the HPLC. Probably this can be explained by conversion of the *n*-sulfocarbamyl toxins (low relative toxicity) to the corresponding carbamate toxins (high relative toxicity) by hydrolysis of acid extraction used in MBA [30]. The fact that the MBA reports somewhat higher toxicity than the HPLC at levels near or below the regulatory limit is beneficial from a food safety standpoint. These conclusions are in agreement with that found in previous study [13]. When the data from MBA and RBA were compared, a *highgood* linear correlation was found $r = 0.809$, this positive correlation means that the concentrations detected by the two methods vary in the same way, which means that there is a great resemblance in the results of those two methods. For high concentrations (above the sanitary threshold), the RBA reported no false results, with regard to the low concentrations, some small differences between the results of the two methods were found; this is probably due to the very high sensitivity of the radioactive assay to detect very low concentrations below the limit of detection of the bioassay.




Figure 9: Correlation between MBA and HPLC-FD method for analysis of PSP in mussel (Number of compared samples = 8)




**Figure 10: Correlation between MBA and RBA method for analysis of PSP in mussel
(Number of compared samples = 12).**

**Table 1 : Summary of results in mussels and solen comparing HPLC-FD and RBA against the reference
MBA (ST : Sanitary threshold RSD : Relative Standard Deviation)**

	RBA	HPLC-FD
Number of samples	12	8
Mean ratio to (RSD%)	61.41%	104.6%
Pearson correlation coefficient (r)	0.809	0.79
Linear refression slope equation	$y=0.6404x+675.6$	$y=0.3263x-28.867$
Other method> ST ; MBA< ST	3 (25 %)	0
Other method< ST ; MBA> ST	0	1 (12.5%)
Other method and MBA both < or > ST	9 (75 %)	7 (87.5 %)

4. CONCLUSION

361 The comparison between three methods used for the analysis of PSP in shellfish from South Atlantic
 362 Moroccan coasts shows that HPLC was more sensitive and more specific than MBA and RBA. This
 363 chemical method allows for detection of small amount and identification of the different compounds of
 364 PSP toxins present in samples of this region. The higher costs of HPLC testing (expensive equipment,
 365 purchase frequently several types of standard PSP) and longer testing time mean that many
 366 laboratories use the mouse bioassay for investigation of PSP toxins in shellfish. The three MBA, HPLC
 367 and RBA methods report the total toxicity in $\mu\text{g\_STX\_equi./kg}$, but the advent of the chemical
 368 method, the data of can also be presented as a complete profile of the toxin.
 369 The RBA test has allowed detecting low concentrations, use of a minimal amount of the standard and
 370 analysis of several samples, but this method requires adequate facilities for the management of
 371 radioactive waste despite the low radioactive emission of tritium used in the marking of the STX.
 372 For the MBA, among the known drawbacks for the use of laboratory animals and the high price of
 373 standard STX frequently used to standardize this biological test, it remains effective for monitoring PSP
 374 toxins. During this PSP investigation by using bioassay, no human intoxications were reported.
 375 Concerning investigation of toxic phytoplankton, *A. minutum* was identified and confirmed as
 376 dinoflagellate responsible for the production of PSP toxins. However, the relationship between shellfish
 377 toxicity levels distribution, occurrence and abundance of vegetative cells of *Alexandrium* in southern
 378 Moroccan coasts is complex.
 379 In Morocco, in order to protect the consumer from paralytic shellfish poisoning, an alert system (official
 380 monitoring program) is established for stopping the collection and commercialization of shellfish. These
 381 activities are permitted when the toxin concentrations are below the sanitary threshold value.
 382
 383

Comment [SCh2]: ???

References

- 384 1. Mons, M.N., Van Egmond, H.P. & Speijers, G.J.A. 1998. Paralytic shellfish poisoning: A
 385 review. RIVM Report 388802 005. June 1998.
- 386 2. Quilliam, M.A. Committee on Natural Toxins and Food Allergens. Phycotoxins. General
 387 Referee Reports. JAOAC. 2001; Int. 84(1): 194-201.
- 388 3. Bourhili, E.H., Intoxication alimentaire par la moule (51 cas), Agadir, Octobre et Novembre
 389 1982. Thèse N° 33 Fac. Med. et Pharm., Rabat. 1982; 84p. French.
- 390 4. Tber A. Communication personnelle. Laboratoire d'analyses des services vétérinaires.
 391 Casablanca, Maroc. 5 p. 1983.
- 392 5. Taleb H., Vale P. et Blaghen M. Spatial and Temporal evolution of PSP toxins along the
 393 Atlantic shore of Morocco. 2003; Toxicon 41(2): 199-205.
- 394 6. Taleb H., Vale P., Jaime E., Blaghen M., Study of paralytic shellfish poisoning toxin profile in
 395 shellfish from the Mediterranean shore of morocco. Toxicon, 2001; 39 (12): 1855-1861.
- 396 7. Amanhir R, Benhadouch A, TalebH, Rijal Leblad B, Vale P, Blaghen M. International Journal
 397 of Biochemistry Research & Review, 2017; 17(2): 1-11, 2017; Article no.IJBCRR.14413.
- 398 8. Abouabdellah R, Taleb R, Bennouna A, Erler K, Chafik A, Moukrim K. Paralytic shellfish
 399 poisoning toxin profile of mussels *Perna perna* frsouthern Atlantic coasts of Morocco.
 400 Toxicon, 2008; 51(4): 780-786.
- 401 9. AOAC (Paralytic Shellfish Poison, Biological method, Final Action, Method 959.08, in Official
 402 Methods of Analysis of the Association of Official Analytical Chemists (AOAC), Natural
 403 Toxins, 1995 Chapter 49, page 46).
- 404 10. Thielert G., Kaiser I. and Luckas B., HPLC determination of PSP toxins. In: J.M. Fremy
 405 (eds.) Proceedings of symposium on marine biotoxins. Editions CNEVA, Maisons-Alfort.
 406 1991; pp. 121-125.
- 407 11. Yu R.C, Hummert C, Luckas B, Qian P.Y, Yhou M.J., Modified HPLC method for analysis of
 408 PSP toxins in algae and shellfish from China. Chromatographia, 1998; (48):671- 675
- 409 12. Diener, M, Erler K, Hiller S, Christian B and Luckas B. Determination of paralytic shellfish
 410 poisoning (PSP) toxins in dietary supplements by application of a new HPLC/FD method. Eur
 411 Food Res Technol 2006; 224, 147-151.
- 412 13. Van Dolah, F.M, Fire,S.M, Leighfield, T, A., Mikulski, C, M, Doucette, G, J, 2012.
 413 Determination of Paralytic Shellfish Toxins in Shellfish by Receptor Binding Assay:
 414 Collaborative Study Journal of AOA C International 2012; Vol. 95, No. 3,.
- 415 14. Uthermol, H., Zur Vervollkommung der quantitativen phytoplankton .Methodik. mitt. int.
 416 Ver.Theor. Angrevo. Limnol 1958; (9): 1-38.
- 417 15. Anhichem M, Dellal M, Chfiri r, Bouthir FZ, Yahyaoui A, Tazi O, Benbrahim S. Etude de la
 418 qualité des eaux et des sédiments de la baie de Dakhla (maroc) physic-chimie et

- contamination métallique. 5ème congrès franco-maghrébain de zoologie et ichthyologie, 2016, 8-10 December.
16. McCarthy M., Bane V., García-Altares M., van Pelt F. N. A. M., Furey A., O'Halloran J. Assessment of emerging biotoxins (pinnatoxin G and spirolides) at Europe's first marine reserve: Lough Hyne. *Toxicon* 2015; (108): 202–209.
17. Hashimoto, K., Noguchi, T. Recent studies on paralytic shellfish poison in Japan. *Pure. Appl. Chem.* 1989; 61, 7–18.
18. Anderson, D.M, Alpermann, T.J, Cembella, A.D.; Collos, Y.; Masseret, E.; Montresor, M. The globally distributed genus *Alexandrium*: Multifaceted roles in marine ecosystems and impacts on human health. *Harmful Algae* 2012; 14, 10–35.
19. Bennouna A. Le phytoplancton du littoral atlantique marocain Abda-Doukalla : Organisation, Dynamique et impact des taxons potentiellement toxiques, Doc National, Université Chouaib Doukkali, 2008; 50p. French.
20. Makaoui A, Idrissi M., Benazzouz A., AitLaame M, Agouzouk A., Larissi J., and Hilmi K.: Hydrological variability of the upwelling and the filament in Cape Juby (28°N) Morocco (2010). *International Journal of Advanced Research* (2015), Volume 3, Issue 10, 9 – 26.
21. Sommer, H., Meyer, K. F. Paralytic shell-fish poisoning. *Am. Med. Assoc. Arch. Pathol.* 1937; 24, 560-598.
22. Fernandez, M.L. Regulation of marine microalgae toxins: towards harmonization of methods and limits. *S. Afr. J. Mar. Sci.* 2000; (22). 339-346.
23. Guallar-Morillo. Pressions environnementales et niches écologiques d'*Alexandrium minutum* sur les côtes françaises. 2014; Journées RePHY, Nantes.
24. Pitcher GC, Cembella A.D., Joyce L.B., Larsen J., Probyn T.A., Ruiz Sebastia'n C. The dinoflagellate *Alexandrium minutum* in Cape Town harbour (South Africa): Bloom characteristics, phylogenetic analysis and toxin composition. *Harmful Algae* 2007; (6)823–836.
25. Cembella A.D, S. E Shumway and R. Larocque. Sequestering and putative biotransformation of PSP toxins by the sea scallop *Placopecten magellanicus*: seasonal and spatial scales in natural populations. *J. Exp. Mar. Biol. Ecol.* 1994; 180: 1- 22.
26. Abouabdellah R., A. Bennouna, J. El Attar, Katrin Erler, Abdelghani Chafik. The detection of PSP and DSP toxins in bivalves from the southern Moroccan coasts by HPLC and mouse bioassay. *Marine and Freshwater Toxins Analysis. Second Joint Symposium and AOAC Task Meeting* Baiona, 2011; Spain May 1-5.
27. Giacobbe MG, Maimone G First report of *Alexandrium minutum* Halim in a Mediterranean lagoon. *Cryptogam Algolog* 1994; (15):47–52
28. Forteza, V., Quetglas, G., Delgado, M., Reyero, M., Fraga, S., Franco, JM. & Cacho, E. Toxic *Alexandrium minutum* bloom in Palma de Mallorca harbour, (Balearic Islands), western mediterranean. In Reguera, B., Blanco, J., Fernandez, M. & T. Wyatt, eds. 1997. *Harmful Algae, Proceedings of the VIII International Conference on Harmful Algae*, (June 1999, Vigo, Spain), pp. 270-273. Xunta de Galicia and IOC of UNESCO.
29. Vila M, Camp J, Garcés E, Masó M, Delgado M High resolution spatio-temporal detection of HABs in confined waters of the NW Mediterranean. *J Plankton Res* 2001; 23(5): 497–514
30. Indrasena, W.M. and Gill, T.A. Thermal degradation of partially purified paralytic shellfish poison toxins at different times, temperatures and pH. *J. Food Sci.* 2000; 65(6): 948-953.