

# 1    **Nutritional and Phytochemical Composition of *Vitellaria paradoxa* (Shea fruit pulp)**

2  
3

## 4    **Abstract**

5    The nutritional and phytochemical compositions of the **Shea fruit** (*Vitellaria paradoxa*) were  
6    investigated using standard methods. The phytochemical screening **reveal** the presence of  
7    tannins, alkaloids, saponins, carbohydrates, steroids, flavonoids, anthraquinones and cardiac  
8    glycosides **while** the proximate analysis **(%)** showed the moisture (75.4), ash (11.6), crude fat  
9    (10.6), crude protein (4.4), crude fibre (9.1), carbohydrate (17.8) and energy value (385.2  
10    kcal/g) contents. **Also**, the mineral composition (%) of the dried Shea fruit pulp as evaluated  
11    in this study **consist** of K (1.97), Na (0.47), Ca (5.50), Mg (1.75), P (1.24); Zn (1.81) and Fe  
12    (3.01). The results obtained in this study further reiterate the reason why there is high  
13    consumption of Shea fruits by farmers in West and Central Africa for both its nutritional and  
14    medicinal benefits.

15

16    **Key words: Nutritional properties, Phytochemicals, Shea fruit pulp**

17

## 18    **Introduction**

19    Traditional and folk medicines derived from plants-stem, bark, root, fruit, leaves and seeds  
20    have been discovered to play significant role in the prevention and cure of diseases, sickness  
21    and disorders in humans around the globe (Gabbe *et al.*, 2006). About three quarter of the  
22    world's population relies on plants and their extracts for healthcare (Premanathan *et al.*,  
23    2006). More so, a good number of people especially those living in the rural areas **depends**  
24    **largely** on the therapeutic effects of herbs because most of these effects have stood the **taste**  
25    **of time** particularly for the treatment of allergic, metabolic, cardiovascular and other  
26    degenerative or life threatening diseases and as well as food sources (Igole *et al.*, 2005), this  
27    is because of their bioactive and nutritional compositions (Lamien *et al.*, 2007). Generally,

fruits from plants particularly those from agro forestry species have been reported as good source of dietary supplement because of their vital minerals and vitamins which provides the nutritional and energy requirements needed by the human body to grow and also curtail malnutrition among children (FAO, 2005).

Fruits from agro forestry species such as that of *V. paradoxa* tree have been recognized by the rural dwellers to supplement their daily energy need (FAO, 2005). *Vitellaria paradoxa* commonly known as Shea butter tree (*Sapotaceae*); is the only specie in the genus indigenous to Africa and found prominently growing in west and central Africa (Lovett *et al.*, 2000). The Shea tree fruit consist of a thin tart nutritious pulp that surrounds a relatively large oil-rich seed from which Shea butter is extracted. The importance of the Shea nut is considered second to the palm nut because of the benefit of butter to many industries both locally and internationally in cosmetics, pharmaceutical and food uses (Akihisa *et al.*, 2010). The sweet pulp is consumed locally when ripped because it is a rich source of sugars, calcium, iron, potassium, magnesium and phosphorus (Maramz *et al.*, 2004). The major constituent of the pulp is vitamin C which is required as an essential nutrient by both humans and animals as an anti-oxidant (Niji *et al.*, 2002). Vitamin C also facilitate lowering of hypertension, enhance human immunity and lack of it may result in scurvy (tooth bleed) diseases (Lang *et al.*, 2007). The pulp is also anti-carcinogenic hence have chemotherapeutic quality It is also used as pain reliever in bones, nerves, inflammation, dislocation and joints arthritis (Akihisa *et al.*, 2010). The oil is also used as local lamp illuminant and for soap production. The fruit contains carbohydrates such as glucose, fructose and galactose (Aremu *et al.*, 2006) and usually, the period of harvesting of the fruit coincides with the high energy requirements in farm planting and the consumption of this fruit meet this immediate energy demand by farmers after hard labour (Okullo *et al.*, 2010). The objective of this study is to evaluate the nutritional and phytochemical properties of dried Shea fruit pulp.

53 **Materials and methods**

54 **Sample collection and preparation**

55 The mature *Vitellaria paradoxa* (Shea butter) fruit were collected from Dzwayagi village,  
56 Gbako Local Government Area in Niger State in June, 2017 following which it was depulped  
57 manually. The pulp comprising of the epicarp and mesocarp were oven dried at 35°C to a  
58 constant weight for five (5) days and then pulverized using laboratory blender, sieved (mesh  
59 size 350µm); the fine powder obtained after sieving was weighed then packaged in a  
60 transparent air tight glass bottle and stored at ambient temperature (31±2°C) for analyses.

61 **Proximate analysis**

62 The proximate content of the dried pulverized Shea fruit was carried out according to  
63 standard methods described by AOAC (2012) to estimate the ash, crude protein, crude fat,  
64 crude fibre and carbohydrate contents of the sample.

65 **Mineral analysis**

66 The analysis of the mineral constituents of the sample were carried out using Atomic  
67 Absorption Spectrophotometer (Model 320N, Surgicare, England) for Mg, Zn, Ca, Fe, P  
68 while the Na and K contents were determined using Flame Photometer (Model FP 640,  
69 Mumbai, India) using the methods of AOAC (2003).

70

71

72 **Phytochemical analysis**

73 The phytochemical analysis of the methanol extract of the Shea fruit pulp (*Vitellaria*  
74 *paradoxa*) was carried out according to standard methods as described by Sofowora (1993)  
75 and Trease and Evans (2012).

76 **Statistical analysis**

77 The mean data of the proximate and mineral analyses were computed.

78 **Results and discussion**

79 The result in Table 1 showed the nutritional composition of the Shea fruit pulp with moderate  
 80 moisture content of 75.39% which contributes to it being perishable if poorly preserved but  
 81 which also contributes to its softness and edibility. The moisture content of the dried Shea  
 82 fruit pulp obtained in this study (Table 1) is in agreement with the moisture content of 72.4 -  
 83 75.3% reported by Okullo *et al.* (2010). The total carbohydrate content of 17.96% obtained in  
 84 this study is within the range of the carbohydrate value (12.4-19.4%) reported in a similar  
 85 study by Neuwinger (1994). Like in other edible fruits, the Shea fruit pulp is rich in  
 86 carbohydrate content such as galactose, fructose and glucose which serves as source of high  
 87 energy required by farmers during farm planting. Shea fruit has been reported to contain  
 88 more carbohydrates that are vital in nutrition and as good source of energy (Anwange *et al.*,  
 89 2004) and it is believed that regular eating a Shea fruit after hard labour could provide  
 90 immediate energy sources (Neuwinger, 1994).

91 Also, the result in Table 1 showed that the crude fibre content in the dried Shea fruit was 9.06  
 92 %; crude fibre helps in the maintenance of normal peristaltic movement of the digestive tract.  
 93 Thus diet containing high crude fibre may reduce disorders such as constipation, colon  
 94 diseases, diabetes, heart diseases and obesity (Omosuli *et al.*, 2009). The crude protein  
 95 (4.49%) in the Shea fruit obtained in this study is higher than that reported on Shea fruit pulp  
 96 in Uganda by Prokarite (2007). This variation may be due to differences in soil, climate and  
 97 other environmental conditions. Also, the crude fat content (10.6%) obtained in study is  
 98 higher than the 1.5-3.5% reported by Wilhelmina (2004) in his study on Shea fruit in Ghana.  
 99 This may be due to differences in post harvest handling processing. The ash content (11.55%)

of the Shea fruit pulp obtained in this study is comparable to those reported for other edible fruits across the West African region. Ash content of fruits and seeds is important because it determines the amount of minerals and trace metals that are vital in human dietary intake (Onwuka, 2005). Furthermore, the high energy value of Shea fruit recorded in this study (385.21 Kcal/g) is an indication of its high carbohydrate content and as well as the varieties of carbohydrates (fructose, galactose and glucose) it contains (Neuwinger, 2004); this therefore maybe the reason why it is highly patronize by farmers in West and Central Africa (Yeshajau and Clifton, 2004).

#### **Mineral composition in Shea fruit pulp**

The result in Table 2 showed that the Shea fruit pulp investigated in this study, contains Na, K, Mg, Fe, P, Zn, and Ca in various amounts which are important minerals that constitute the nutrients required by human and animals for growth and development (Agatemor and Ukhun, 2006). The K content in Shea fruit obtained in this study (1.97%) is comparable with the 2.0% K content reported in other wild and domesticated edible fruits. Potassium plays a vital role in protein, synthesis, body fluid balance, nerve and muscular function, glucose and glycogen absorption and regulating blood pressure (Omosuli *et al.*, 2009). The amount of Na present in the studied Shea fruit pulp is 0.47% and is still within the recommended daily allowance of 0.50 % as reported by NRC (2009). From this study, the Na/K ratio was calculated to be 0.24 % which is less than 1 %; this indicates that Shea fruit pulp is recommended for consumption and in the management and control of hypertension (NRC, 2009). The amount of Ca estimated in the Shea fruit pulp is 5.50% which indicates that Ca is the major mineral present in the Shea fruit pulp although it is lower than the 8.8 % Ca reported in a similar study (Okullo *et al.*, 2010). Calcium play the role of cofactor in many physiological and metabolized functions such as in bone and teeth formation, nervous system, hormonal secretions, enzyme activations, and blood clotting (Agatemor and Ukhun, 2006).

The result in Table 1 also showed that the Shea fruit pulp analyzed in this study contains P (1.24%), Zn (1.86%) and Fe (3.01%). Also, the Mg in the Shea fruit pulp (1.75%) as evaluated in this study is below the 6.0% recommended by NRC (2009) in similar fruit pulp. Magnesium is an important element required for release of enzymes required for synthesis and break down of carbohydrates, fats, proteins in synthesis of RNA and DNA (Grober *et al.*, 2005).

### **Phytochemical constituent of Shea fruit pulp**

The presence of alkaloids, saponins, tannins, anthraquinones, terpenes, flavonoids, carbohydrates and glycosides as phytochemical constituents in the Shea fruit pulp (Table 2) may be the basis of its medicinal and health properties and hence explains why it has been used to cure and prevent various illness and diseases (Hassan *et al.*, 2004); therefore, the valuable pharmaceutical properties of *V. paradoxa* may be due to the presence of these bioactive compounds (Table 2). Alkaloids are heterocyclic nitrogen containing compounds found to have anti-malaria, antibacterial, anti-cancer, analgesic, anti-asthma, anti-hyperglycemic and anti-arrhythmic effect (Karou *et al.*, 2007). This may be responsible for the use of the Shea pulp in treating diseases such as cancer, body pain and diabetes. Tannins are bioactive compounds with hot astringent taste that are toxic to microorganisms therefore, their presence in Shea fruit pulp may be the basis for their anti-diarrheal and anti-hemorrhage roles in human (Asquith and Butter, 1996). Furthermore, flavonoids are polyphenolic bioactive compounds which reduces the risk of cardiovascular disorder and lower hypertension; and this may be the reasons why Shea fruits is eaten by rural dwellers especially those diagnosed as being hypertensive (Edeoga *et al.*, 2005). Also, the presence of saponins could be responsible for the use of the Shea fruit ash in local soap making and to cure skin diseases. Saponins generally have soapy feel, hemolytic activity and lowering of

149 cholesterol thus the pulp is chewed raw to reduce high cholesterol in man (Ahmad and  
150 Geelen, 2013).



## 151 Conclusion

152 The findings of this research show that Shea fruit pulp has a high nutritional potential and  
153 health benefits. Therefore, large scale cultivation of *V. paradoxa* should be promoted so as to  
154 meet up with its high consumption.

155

## 156 References

- 157 Agatemor, C. and Ukhun, M. E. (2006). Nutritional potential of the nut of tropical almond  
158 (*Terminalia catappa* L.). *Pakistan Journal of Nutrition*, 5(4): 334 – 336.
- 159 Ahmad, F. and Geelen, D. (2013). Saponins and their role in biological processes in plants  
160 and animals. *Phytochemistry Review*. 12(4): 877 – 893.
- 161 Akihisa, T. (2011). Triacylglycerol and triterpene ester composition of Shea nut from seven  
162 African countries. *Journal of Oleo Sciences*, 60(8): 385 – 391.
- 163 Akihisa, T., Kojima, N., Kikuchi, T., Yasukawa, K., Tokuda, H., Masters, E. T., Manosroi,  
164 A. and Manosroi, J. (2010). “Anti-inflammatory and chemopreventive effects of  
165 triterpene cinnamates and acetates from Shea fat”. *Journal of Oleo Sciences*, 59 (6): 273  
166 – 280.
- 167 Anwange, B. A., Ajibola, V. O. and Oniye, S. J. (2004). Chemical studies of the seeds of  
168 *Moringa Obifera* (Lam) and *Detarium microcarpum* (Gull and Sperr). *Journal of*  
169 *Biological Sciences*, 8 (4): 711 – 715.
- 170 AOAC (2003). *Standard Methods of Analysis* (20<sup>th</sup> ed). Association of Analytical Chemist  
171 Washington D C, USA. pp 5 – 6.
- 172 AOAC (2012). *Standard Methods of Analysis* (26<sup>th</sup> ed). Association of Analytical Chemist  
173 Washington D C, USA. pp 10 – 13.
- 174 Aremu, M., Olonisakin, A., Bako, D. and Madu, P. (2006). Compositional study and physio-  
175 chemical characteristics of cashew nut (*Anarcadium occidentals*). *Pakistan Journal of*  
176 *Nutrition*, 5:328 – 333.
- 177 Asquith, T. N. and Butter, L. G. (1996). Introduction of condensed tannins with selected  
178 proteins. *Phytochemistry*, 25(7): 1591 – 1593.
- 179 Edeoga, H. O., Okwu, D. E. and Mbaebie, B. O. (2005). Phytochemical constituents of some  
180 Nigerian plants. *African Journal of Biotechnology*, 4(6): 685 – 688.
- 181 FAO (2005). *Non-Wood Forest Products in Africa: A Regional and National Overview*. Food  
182 and Agriculture Organization, Rome. pp 303 – 305.
- 183 Gabbe, S. and Tatke, P. (2006). Evaluation of the Immunomodulatory activity of methanol  
184 extract of *Fiasis benghalonsis* roots in rats. *Indian Journal of Pharmacology*, 38(4): 271  
185 – 274.
- 186 Grober, U., Schmidt, J. and Kiesters, K. (2005). *Nutrients*. Oxford University Press London.  
187 pp. 23 – 25.

- Hassan, M., Oyewale, A., Amupitan, J., Abdullahi, M. and Okwonko, E. (2004). Preliminary phytochemical and antimicrobial investigations of crude extracts of root bark of *Detarium microcarpum*. *Journal of Chemical Society of Nigeria*, 29: 26 – 29.
- Igoli, J., Ogaji, O. and Torayin, A. (2005). Traditional medicine practices amongst people of Igede, Nigeria. *African Journal of Trad.*, 2(2): 134 – 152.
- Karou, D. A., Savadogo, A., Canni, C., Yamego and Montesano, C. (2007). Antibacterial activity of alkaloids from *Sida acuta*. *African Journal of Biotechnology*, 5: 195 – 200.
- Lamien, N., Tiabu, M., Guiniko, S. and Oden, P. (2007). Variation in denchrometric and fruiting characteristics of *Vitellaria paradoxa*: Population and multivariate models for estimation of fruit yield. *Agroforestry Systems*, 69(1): 9 – 11.
- Lang, W. and Bulley, S. (2007). What control vitamin C levels in plants? Accessed on 2/9/2008 <<http://www.isl.vt.edu/articles/oct0703.htm>>
- Lovett, P. and Hag, N. (2000). Evidence for a tropic selection of the Shea nut tree (*Vitellaria paradoxa*). *Agro-forestry Systems*, 48: 293 – 304.
- Maranz, S., Kpikpi, W., Wasman, Z. and Champagne, B. (2004). Nutritional values and indigenous preferences for Shea fruit (*Vitellaria paradoxa*) in African Agro-forestry park lands. *Journal of Economic Botany*, 58(4): 588 – 600.
- Neuwinger, H. D. (1994). African ethno-botany. poisonous drugs, chemistry. *Pharmacology and Toxicology Journal*, 3 (4): 823 – 826.
- Niji, F. F. and Onajobi, F. (2002). The vitamin C contents of some tropical fruits challenges to organic farming and sustainable land use in the tropics and sub tropics. Book of Abstracts, University of Benin. Institute of Animal Nutrition. Volume 3. pp273.
- NRC (2009). *Recommended Dietary Allowances* (10 ed). National Research Council, National Academic Press Washington DC. pp 24 – 25.
- Okullo, J. B. L., Omujal, F., Agea, J. G., Vuzi, P. C., Namutebi, A., Okello, J. B. A. and Nyanzi, S. A. (2010). Proximate and mineral composition of Shea (*Vitellaria paradoxa*) Fruit pulp in Uganda. *African Journal of Food, Agriculture, Nutrition and Development*, 11(10): 4430 – 4443.
- Omosuli, S., Ibrahim, T., Oloye, D., Agbaje, R. and Jude-Ojei, B. (2009). Proximate and mineral composition of roasted and defatted cashew and Shea nuts. *Pakistan Journal of Nutrition*, 8(10): 1649 – 1651.
- Onwuka, G. (2005). *Food Analysis and Instrumentation, Theory and Practice*. Napthali Publishers, Nigeria. pp. 63 – 63.
- Premanathan, M., Ajendran, S., Ramanathan, T., Kathuresan, K., Nakashima, H. and Yamamoto, N. (2000). A survey of some Indian medicinal plants for anti-human immune deficiency virus (HIV) activity. *Indian Journal of Medicinal Research*, 112: 73 – 77.
- Prokarite (2007). La Base de Donnees. Vitellaria, World Agroforestry Center. <http://prokarite.org/vitellaria.dbase/fat-percentage.html>, Retrieved on 11/03/2018.
- Sofowora, A. (1993). *Medicinal Plants and Traditional Medicine in Africa* (2<sup>nd</sup> ed.). Spectrum Books Ltd., Ibadan, Nigeria. pp. 134 – 156.
- Trease, G. E. and Evans, N. C. (2012). *Pharmacognosy* (15<sup>th</sup> ed) Elsevier Publishers, London. pp. 42 – 93.
- Wilhelmina, K. (2004). Effects of production and processing factors on major fruits and vegetables anti-oxidants. *Journal of Food Science Research*, 13(1): 427-434
- Yeshajau, P. and Clifton, E. (2004). *Food Analysis, Theory and Practice* (3<sup>rd</sup> ed). GBS Publishers, New Delhi. pp 602 – 621.



**Table 1: Proximate composition of dried Shea fruit pulp**

Analysis	Concentration <sup>1</sup>
<b>Proximate content (%)</b>	
Moisture	75.39 ± 0.01
Ash	11.55 ± 0.78
Crude fat	10.60 ± 0.22
Crude protein	4.49 ± 0.28
Crude fibre	9.06 ± 0.67
Carbohydrate	17.97 ± 0.58
Energy value (Kcal/g)	385.21 ± 4.26
<b>Mineral content (%)</b>	
Potassium	1.97 ± 0.01
Sodium	0.47 ± 0.01
Calcium	5.50 ± 0.14
Magnesium	1.75 ± 0.01
Phosphorus	1.24 ± 0.11
Zinc	1.81 ± 0.12
Iron	3.01 ± 0.13

<sup>1</sup>Each data is mean ±SD of triplicate determinations

**Table 2: Qualitative phytochemical constituent of methanol extracts of dried Shea fruit pulp**

Phytochemical constituent	Inference <sup>1</sup>
Saponins	++
Flavonoids	++
Alkaloids	+++
Tannins	++
Anthraquinones	+
Carbohydrates	++
Steroids	+++
Cardiac glycosides	+

<sup>1</sup> Key ++ = high concentration; +++ = very high concentration and + = moderate concentration