

Original Research Article

Temporal and spatial changes in host defense enzymes in response to *Fusarium* wilt in Pigeonpea (*Cajuns cajan* (L.) Millspaugh)

ABSTRACT

Fusarium wilt caused by *Fusarium udum* Butler (*Fud*) is a constant threat to pigeonpea productivity in several parts of the world. Understanding the molecular basis of pigeonpea-*Fud* interaction is necessary to improve resistance to *Fud* and thereby the productivity of pigeonpea. Temporal and spatial changes were studied in defense enzymes namely catalase, peroxidase (PO), polyphenol oxidase (PPO), Phenylalanine ammonia lyase (PAL), β -1,3-glucanase and chitinase at pre-initiation (S1), disease initiation (S2) and severe diseased (S3) stage in root, stem and leaf tissues of 2 susceptible and 2 resistant cultivars of pigeonpea inoculated with *Fud*. Comparison of various treatment combinations revealed that all the defense enzymes showed temporal and spatial variation in activity/expression pattern. PO increased in all the tissues of all the varieties from S1 to S2 and declined thereafter in S3 stage and induction was higher in resistant cultivars. Catalase and PPO were induced highly in resistant cultivars inoculated with *Fud* and pattern was stage-specific and tissue-specific in nature. Catalase induction occurred at S2 in uninoculated plants where as it started at S1 in *Fud* inoculated plants. PPO level diminished from S2 to S3 stage and in susceptible varieties it reached even below the control. PAL, β -1,3-glucanase and chitinase activity were induced at S1 and S2 and values at S3 decreased progressively. Induction of chitinase was significantly high in leaves whereas β -1,3-glucanase activity was high in stem tissues. Both control and challenged plants had higher level of β -1,3-glucanase activity at later stages of disease development, but the proportionate increase was much higher in resistant cultivars. The activity/expression pattern of these defense enzymes reveals their use as established resistance markers and provides scope for manipulating their expression and development of wilt-resistant transgenic pigeonpea.

Keywords: Pigeonpea, India, *Fusarium udum*, Pathogenesis, Defence enzymes, antioxidant enzymes, PR proteins

Abbreviations:

gfw⁻¹: per gram fresh weight, gdw⁻¹: per gram dry weight, PAL: Phenylalanine ammonia lyase, PO: Peroxidase, PPO: Polyphenol oxidase, PR: Pathogenesis-related, ROS: Reactive oxygen species

1. INTRODUCTION

Pigeonpea (*Cajuns cajan* (L.) Millspaugh) is among the world's most important pulse crops. Around 76% of the total global area and around 73% of total global production of pigeonpea is in India. Globally it is cultivated on 4.75 million hectares producing 3.68 million tonnes with productivity of 774.30 kg/ha [1]. Wilt caused by *Fusarium udum* Butler is an important disease of pigeonpea in north, central and south India inflicting heavy losses in yield [2]. Plants activate a large array of defence mechanisms in response to pathogen attack and crucial factor determining the success of these mechanisms is the speed of their activation. There is a considerable interest in understanding how plants recognize pathogen attack and control expression of defence mechanisms. After pathogen recognition, highly localized biochemical events are rapidly induced to inhibit the further development of the attacking pathogen. The rapid production and accumulation of reactive oxygen species (ROS), particularly the superoxide anion (O₂⁻) and hydrogen peroxide (H₂O₂), culminates with a hypersensitive response (HR) and localized programmed cell death (PCD) to impair the pathogen establishment and development inside the host tissues and deprive the pathogen of further access to nutrients [3]. Concomitant with or following HR, several pathogen defense-related genes are translated into antioxidant enzymes and PR-proteins [4] that confer resistance to the plant in defense against pathogen attack. Enzymes of the phenyl propanoid pathway e.g. phenyl alanine ammonia lyase (PAL) are involved in the synthesis of phenolic compounds like flavonoids, salicylic acid,

coumarins etc. which are known as phytoalexins. Differential expression of PAL genes was investigated using northern blot analysis in chickpea seedlings challenged with *Fusarium oxysporum* f. sp. *ciceri* [5]. Datta and Lal [6] observed Temporal and spatial changes in phenolic compounds in response to infection by *Fusarium* wilt pathogen in chickpea and pigeonpea and suggested that interaction between *Fusarium* and host plants was found to enhance defense responses against wilt disease in resistant cultivars. Broetto et al. [7] also reported similar results in bean, *Phaseolus vulgaris* L. infected by *Fusarium oxysporum* f. sp. *phaseoli*. The host proteins important for plant defence against pathogens also include enzymes like polyphenol oxidase, peroxidase, superoxide dismutase and "pathogenesis related" (PR) proteins viz. chitinase and β -1,3-glucanase [3, 4, 8].

Higher activity of enzyme in highly resistant cultivars and lower in susceptible ones suggested the possible role of β -1,3-glucanase as a biochemical marker for screening of pearl millet cultivars for downy mildew [9]. Giri et al. [10] reported differential induction of chitinase in susceptible and resistant cultivars of chickpea and decrease in β -1,3-glucanase in the resistant cultivars in response to the pathogen infection. The observed decrease in the resistant cultivars may be associated with the reduction in pathogen invasion as a result of the defence reaction. Stevenson et al. [11] reported that root exudates of chickpea plants containing phytoalexins contribute at least partly to resistance against wilt as they show anti- *F. oxysporum* activity *in vitro*. In a recent study, Oliveira et al. [12] studied time course activities of antioxidant enzymes and PR proteins in cowpea plants infected with *Colletotrichum gloeosporioides* and observed differential induction among resistant and susceptible genotypes and confirmed participation of peroxidase, chitinase and β -1,3-glucanase as important component of host defense. Similarly, Pareek et al. [13] recorded higher induction/activation of chitinase and β -1,3-glucanase in resistant genotypes of *Vigna aconitifolia* inoculated with *Macrophomina phaseolina* and implicated these enzymes as a part of general defense response against fungal pathogens. Gharbi et al. [14] compared biochemical defense responses of susceptible and resistant cultivars of *Olea europaea* to *Verticillium* wilt and observed early activation of peroxidase and polyphenol oxidase and early and simultaneous upregulation of chitinase and β -1,3-glucanase correlated with reduced susceptibility in resistant cultivars.

Biochemical and molecular mechanism of resistance provides information for better understanding of the pathogens and their reaction to crop and therefore helps in management of disease in better way. Keeping this in view, the present investigations were undertaken to correlate and analyse the biochemical basis of disease resistance, by measuring levels of Peroxidase, catalase, PPO (as antioxidant enzymes), PAL (enzyme involved in phenyl propanoid pathway), β -1,3-glucanase and chitinase spatially and temporally in wilt resistant and wilt susceptible cultivars of pigeon pea.

2. MATERIAL AND METHODS

2.1 Plant Material

Two wilt resistant and wilt susceptible cultivars of pigeonpea (*C. cajan* (L.) Millspaugh) were used for biochemical and molecular study and comprised of Asha (ICPL 87119), Maruthi (ICP 8863) in resistant group and Bahar and Type 7 as susceptible.

2.2 Inoculation of plants with the fungus and sample collection for biochemical analyses

F. udum (*Fud*) isolates collected from wilt affected fields of pigeonpea. The pathogens were isolated from fourth-node stem sections taken from wilted pigeonpea plants according to Tullu et al. [15] and were colonized on filter paper, dried in the transfer hood, and aseptically cut into small pieces. The colonized filter paper pieces were placed in potato-dextrose broth and incubated to produce liquid cultures of the pathogen. The liquid cultures were filtered through cheese cloth to remove mycelia. The spore suspension was pelleted by centrifugation. After discarding the supernatant, the conidia were washed with sterile water to adjust the spore suspension to 1×10^6 spores ml^{-1} with a haemocytometer. These isolates were further characterized at the laboratories of Department of Life Sciences, and Department of Biochemistry, I.B.S.B.T, C.S.J.M. University, Kanpur. Single spore culture of fungus was obtained by serial dilution method. Isolated fungus was identified as *F. udum* and their pathogenicity tested on pigeonpea cultivar Bahar in pot experiments. Plastic pots of 30 cm diameter were surface-sterilized with 0.1% (w/v) mercuric chloride (HgCl_2). Pots were filled with 2 kg sterilized soil (three subsequent sterilizations at 1.1 kg/cm^2 for 1 h for 3 days). Seven days before sowing, pots were inoculated with the 14-day-old culture of the pathogen multiplied on sand maize meal water medium (90 g sand, 10 g maize meal and 20 ml distilled, sterilized water) @ 50 g kg^{-1} soil.

Seeds were surface-sterilized using 2% sodium hypochlorite for 3 min, and rinsed in sterile water. Ten seeds of each cultivar were sown in each pot for disease scoring. The root, shoot and leaf tissues were collected separately at 7, 15 and 30 days after sowing (DAS) and were frozen immediately in liquid nitrogen to store at -20°C. Biochemical basis of disease resistance was studied by changes in major biomolecules in *Fusarium* wilt resistant and susceptible genotypes.

2.3 Extraction of antioxidant enzymes

The tissue samples (500 mg fresh weight) were ground in 0.1 M phosphate buffer (pH 7.0) using a pre-chilled mortar and pestle, and homogenate was centrifuged at 15000 X g at 4°C for 30 min and aliquots were used as the source of enzyme.

2.4 Isolation and activity assay of peroxidase (EC 1.11.1.7)

Peroxidase activity was determined according to Kar and Mishra [16] with the following modifications. Assay mixture for the peroxidase comprised: 2 ml 0.1 M of phosphate buffer, pH 7.0, 1 ml 0.01 M of pyrogallol ($C_6H_6O_3$), 1 ml of 0.005 M hydrogen peroxide (H_2O_2), and 1 ml of the 2 times diluted enzyme extract. The reference cuvette contained equal volume of the inactivated enzyme (by boiling) and pyrogallol. This was incubated for 5 min. at 25°C after which the reaction was stopped by adding 0.5 ml of 5% (v/v) H_2SO_4 . The amount of purpurogallin formed was determined by taking the absorbance at 420 nm. One unit of peroxidase activity is defined as that amount of the enzyme which cause change of 0.01 absorbance unit per minute under assay condition and was expressed as U per gram fresh weight ($U\text{ gfw}^{-1}$).

2.5 Isolation and activity assay of polyphenol oxidase (EC 1.14.18.1)

Assay mixture of polyphenol oxidase comprised of same ingredients as in case of peroxidase [16] except H_2O_2 . The reaction mixture was incubated for 5 min. at 25°C and stopped by adding 0.5 ml of 5% (v/v) H_2SO_4 . The amount of purpurogallin formed was determined by taking the absorbance at 420 nm. One unit of polyphenol oxidase activity is defined as that amount of the enzyme which cause change 0.01 absorbance unit per minute under assay condition and was expressed as U per gram fresh weight ($U\text{ gfw}^{-1}$).

2.6 Isolation and activity assay of catalase (EC 1.11.1.6)

The activity of catalase was assayed using method of Braber [17] with the following modification. Five ml of the assay mixture for the catalase activity comprised: 300 μ moles of phosphate buffer, pH 6.8 100 μ moles of H_2O_2 , and 1 ml of twice diluted enzyme extract. After incubation at 25°C for 1 min, the reaction was stopped by adding 10 ml of 2% (v/v) H_2SO_4 and the residual H_2O_2 was titrated against 0.01 N $KMnO_4$ until a faint purple colour persisted for at least 15 sec. A control was run at the same time in which the enzyme activity is defined as that same time in which the enzyme activity was stopped at "zero" time. Catalase activity was expressed as μ mole of H_2O_2 used per minute under the assay condition described, considering the concentration of H_2O_2 using the extinction coefficient 0.036/ μ mol/ml.

2.7 Isolation and activity assay of phenylalanine ammonia lyase (EC 4.3.1.5)

Tissues were homogenized in 0.1 M Borate buffer (pH 8.8) at 4°C. Homogenate was centrifuged at 4°C and 38000 Xg and the supernatant used as enzyme source. The activity of L-phenylalanine ammonia lyase was determined by the method of Zucker [18] with certain modifications. Assay mixture consisted of 1.5 ml Borate buffer, 1.0 ml H_2O_2 , 1.0 ml phenylalanine solution (10 μ M) and 0.5 ml enzyme extract. After incubation of this assay mixture for 2 hrs at 38°C water bath, the change of absorbance at 290 nm compared to a reference sample without added substrate was measured with a spectrophotometer. PAL activity was expressed as $\Delta A_{290nm}\text{ ml}^{-1}\text{ h}^{-1}$.

2.8 Isolation and activity assay of chitinase (EC 3.2.1.14)

Defatted and depigmented tissue powder was stirred with extraction buffer (1:6 w/v in 0.1 M phosphate buffer, pH 6.9 containing 0.05 M NaCl) at 4°C for 12 hrs. The mixture was centrifuged at 10,000 X g for 20 min and the proteins from the supernatants were precipitated by adding ammonium sulphate [$(NH_4)_2SO_4$] to 90% saturation (60 g $(NH_4)_2SO_4$ /100 ml extract). The precipitated protein were collected by centrifugation, resuspended and dialyzed against the extraction buffer [10]. Chitinase activity was determined according to Chen et al. [19] and Tsukamoto et al. [20]. The reaction mixture contained 1.0 ml of colloidal chitin solution (7 mg), 1.0 ml of sodium acetate buffer (50 mM, pH 5.2) and 1.0 ml of suitably diluted enzyme. After incubation at 50°C for 1 hr the released reducing sugar was measured as N-acetyl glucosamine (NAG) equivalents by the method of Reissig et al. [21].

2.9 Isolation and activity assay of β -1,3-glucanase (EC 3.2.1.39)

β -1,3-glucanase activity was estimated using the procedure of Koga et al. [22]. The assay mixture contained 1.0 ml of suitably diluted enzyme and 1.0 ml of 1% laminarin solution in sodium acetate buffer (50 mM, pH 5.2). The mixture was incubated at 40°C for 30 min and the released reducing sugar was measured as glucose equivalents [23].

2.10 Statistical analyses

Data from the enzyme assays were subjected to analysis of variance (ANOVA) followed by Tukey's test and values are expressed as Mean $\pm$ SE.

3. RESULTS

3.1 Spatial and temporal activity of peroxidase

The activities of peroxidase increased in all the tissues of all the varieties from pre initiation (S1) to disease initiation stage (S2) and declined thereafter in S3 *i.e.* severe diseased stage ~~are~~ shown in Fig. 1. This change was prominent in case of all challenged as well as un-inoculated plants. In case of disease initiation stage (S2) *i.e.*, after 15 DAS, irrespective of whether the plants was challenged or un-inoculated, there was a significant increase in activity of peroxidase in all the tissues compared to S1 stage. At this stage conspicuous changes occurred at the rate of increase in activity of the enzyme between resistant and susceptible cultivar upon inoculation with respect to their corresponding control. Upon *Fud* infection, peroxidase activity increased by 6.31% and 25.7% in the roots of resistant varieties Asha and Maruthi (Fig. 1) whereas, in the susceptible cultivars Type 7 and Bahar, rates of increase were 1.6 and 3%, respectively, with respect to their controls. The rate of increase was more in the stem and leaf tissues of resistant varieties. Activity increased by ~~the~~ 12% in stem of resistant cv. Maruthi and 2% in susceptible cv. Type 7 upon inoculation. Highest level of increase (13.5%) in peroxidase activity was observed in leaves of Maruthi and the lowest increase (5%) was observed in Bahar leaves. Irrespective of inoculated or un-inoculated plants, there was a significant decrease in enzyme activity in severe disease stage. But comparative level of induction of peroxidase was higher in resistant cultivars than in susceptible cultivars.




Fig. 1. Peroxidase activity (U gfw⁻¹) in different tissues of Pigeonpea cultivars differing in susceptibility to *Fusarium* wilt

3.2 Spatial and temporal activity of polyphenol oxidase

In pigeonpea roots in control condition, prior to disease initiation, the highest activity (53.42 U gfw^{-1}) was noticed in susceptible variety Bahar and lowest (46.40 U gfw^{-1}) in resistant cultivar Maruthi and the same was true for stem tissue (Fig. 2). At disease initiation stage (S2), enzyme activity was elicited to 21.19 and 52.0% in the stem of the resistant cultivars Asha and Maruthi as compared to their un-inoculated control plants. Upon *Fud* inoculation, the activity increased by 9.80 and 7.56% in the stem tissue of susceptible cultivars Type 7 and Bahar, respectively. Similarly in root tissues at S2 stage, the rate of increase was 16.80, 23.36, 13.44 and 0.77% in Asha, Maruthi, Type 7 and Bahar, respectively, as compared to control. In pigeonpea, it was interesting to see that induction of the activity was up to S3 in resistant cultivars. In resistant varieties the enzyme accumulated in a significant higher rate as compared with the untreated plants. The level of the enzyme diminished from S2 to S3 stage and in susceptible varieties the level reached even below the control level.




Fig. 2. Polyphenol oxidase activity (U gfw^{-1}) in different tissues of Pigeonpea cultivars differing in susceptibility to *Fusarium* wilt

3.3 Spatial and temporal activity of catalase

Type 7 (control) showed highest ($212 \mu\text{mol H}_2\text{O}_2 \text{ used min}^{-1} \text{ gfw}^{-1}$) and Maruthi (control) exhibited the minimum ($179 \mu\text{mol H}_2\text{O}_2 \text{ used min}^{-1} \text{ gfw}^{-1}$) activity. Activity of catalase in leaf tissues was significantly higher compared to root and stem (Fig. 3). Catalase activity in leaves at S1 stage was as high as $222.39 \mu\text{mol H}_2\text{O}_2 \text{ used min}^{-1} \text{ gfw}^{-1}$ (WR 315, control). Activity of catalase at S2 stage in susceptible cultivar Bahar increased by 13%, whereas, in resistant cv. Asha and Maruthi, the increase was 12.7% and 25%, respectively, compared to their corresponding controls in root tissues (Fig. 3). Highest induction was attained at S3 stage in the same tissues. For example, in resistant cv. Maruthi upon infection the increase was 1.65 times with respect to its control. In case of stem tissue, the same pattern of enzyme activity was followed and increase in induction was highest in cv. Asha at S3 (16.47% over control). The same pigeonpea cultivar showed highest increase in catalase activity in leaf tissue as well. Another significant pattern observed in all the cultivars was that compared to S1 Stage the enzyme activity increased in challenged tissues at S2 stage.




Fig. 3. Catalase activity ($\mu\text{mol H}_2\text{O}_2$ used $\text{min}^{-1} \text{gfw}^{-1}$) in different tissues of Pigeonpea cultivars differing in susceptibility to *Fusarium* wilt




Fig. 4. Phenylalanine ammonia lyase activity ($\Delta A_{290\text{nm}}/\text{ml h}^{-1}$) in different tissues of Pigeonpea cultivars differing in susceptibility to *Fusarium* wilt

3.4 Spatial and temporal activity of phenylalanine ammonia lyase

It is evident that irrespective of whether the plants were inoculated or un-inoculated, there was significant increase in PAL activity from S1 to S3 stage (Fig. 4). The activity either slightly decreased or remained unchanged at S3 stage. At S1 stage, highest PAL activity was observed in susceptible pigeonpea cv. Bahar under un-inoculated condition. At S2 stage, increase in activity was mainly in the root tissues of all varieties with respect to the controls (Fig. 4). In resistant cultivars, Asha and Maruthi, activity increased by 36 and 72%. In the susceptible varieties, activity increased by 6 and 9% in root tissues as compared to the respective controls. Significant increase in activity by 12 and 9% was observed Asha and Maruthi, respectively, in the stem tissues at S2 stage. PAL activity in

susceptible cv. Bahar increased by only 3% in stem tissue at this stage. But activity of this enzyme increased by 15% in leaf tissue of susceptible cv. Bahar at S2 stage as compared to its control. At severe disease stage (S3), in case of *Fud* treated pigeonpea, highest activity of PAL in root tissue at S3 stage was observed in root tissues of Asha and lowest in Type 7. At this stage, stem and leaf tissue of resistant cultivars PAL activity showed only marginal increase. In case of susceptible varieties, the activity either remained unchanged or fell below the constitutive level of PAL activity.




Fig. 5. Chitinase activity (U gfw⁻¹) in different tissues of Pigeonpea cultivars differing in susceptibility to *Fusarium* wilt

3.5 Spatial and temporal activity of chitinase

Induction of chitinase in pigeonpea was significantly high in leaves. Likewise peroxidase and polyphenol oxidase, chitinase also showed a marked increase in activity in different tissues upon inoculation of the plants with *Fusarium*. Maximum enzymatic activity was observed in resistant cultivars at S2 stage. It was observed that from S1 to S2 stage, the enzyme activity increased progressively and thereafter showed decrease with progression of the disease and at S3 stage enzyme activity was very low (Fig. 5). In contrast, these changes were much less conspicuous in case of un-inoculated plants, where the increase or decrease was only marginal suggesting role of chitinase of plants in resistance to wilt disease. In the leaf tissues the increase was very clear in all the varieties and at S1 stage enzyme activity was up to 4.37 U gfw⁻¹ (Bahar, control) (Fig. 5). At disease initiation (S2), activity increased by 3.10 and 3.58 fold in the resistant variety Asha and Maruthi as compared to control. On the other hand, in the leaf tissue of susceptible cultivars at S2 stage, the activity increased by 2.29 fold. Similarly in the resistant varieties, enzyme activity increased 3.0 fold in root and stem tissue, whereas in susceptible variety, it increased 2.0 fold with respect to their individual control at S2. At S3 stage the activity was slightly decreasing or it was almost unchanged in all the cases (Fig. 5).




Fig. 6. Glucanase activity ($\text{mM h}^{-1} \text{gfw}^{-1}$) in different tissues of Pigeonpea cultivars differing in susceptibility to *Fusarium* wilt

3.6 Spatial and temporal activity of β -1,3-glucanase

Glucanase activity was high in stem and leaf tissues and upon inoculation of the plants with *Fusarium*, a marked increase in glucanase activity in different tissues was observed. In general the maximum enzymatic activity was observed in resistant cultivars at S2 stage. This level of activity was more or less maintained up to S3 stage. This is in contrast to the activity pattern of other enzymes, where the levels showed significant decrease at S3 stage. Both control and challenged plants had higher level of enzymatic activity at later stages of disease development, but the proportionate increase was much higher in resistant cultivars (Fig. 6). At S1 stage, maximum enzymatic activity was up to $0.76 \text{ mM glucose h}^{-1} \text{gfw}^{-1}$ (Asha, control, root) while in cv. Bahar the activity was lowest (0.47 mM). Similarly, there was marked increase in glucanase activity at S2 stage. The resistant cultivars Maruthi showed maximum increase in glucanase activity (17 fold) in leaves. This increase in activity was lower in susceptible cultivars where increase was only up to 1.3 fold in root tissues (Fig. 6).

4. DISCUSSION

Plants have developed a variety of complex mechanisms that involve the biosynthesis and accumulation of metabolites and novel and constitutive proteins that directly or indirectly function in the defense response to pathogens. In the present study, interaction between *Fusarium* and host plants was found to enhance defence responses against wilt disease in resistant cultivars of pigeonpea and the induction of plants' own defence system started only after the infection by respective pathogen, and subsequently resulted in hypersensitive reaction conferring resistance. Also, pigeonpea plants exposed to *Fud* showed enhanced synthesis of pathogenesis related proteins, activities of PAL and other antioxidant enzymes relative to their controls.

4.1 Spatial and temporal activity of anti-oxidant enzymes (PO, PPO and catalase)

The present results revealed increased PO, PPO and catalase activities in pigeonpea tissues after inoculation with *Fud* and the activities were higher in the resistant cultivars at disease initiation stage in comparison to susceptible cultivars. These findings are in conformity with Rath et al. [24] who recorded higher PO activity in leaves of resistant variety. Inoculation with *Fusarium* caused increase in enzyme activity in leaves of both susceptible and resistant varieties, but at severe disease stage, activities recorded significant decline in the susceptible varieties falling even below the control levels in few cases. However, the rate of increase in activity continued up to 30 DAS, upon induction, though the total enzyme concentration was declining at this stage. Xue et al. [25] reported 2-fold increase in total peroxidase in bean tissues after inoculation with binucleate *Rhizoctonia* species in comparison to control.

Other than the site of infection in roots, peroxidase activity also showed marked increase even in stems and leaves. This spatial regulation of enzyme activity in distant tissues has been reported by other workers as well [26]. Tissues of plants attacked by necrotizing pathogens including *Rhizoctonia* and *Colletotrichum* species have been shown to accumulate substantial amount of the enzymes not only at the site of infection, but also in distant tissues. The presence of enzymes in infected plants has been associated with lesion development and limiting the spread of disease [27-29]. The shift in activity of peroxidase can vary, motivating the discussion about its active role in resistance. Peroxidase action could still happen in an indirect way through the activity of subproducts, which possess an antimicrobial activity or by inducing the formation of structural barriers. Higher activity of PPO was observed in pigeonpea resistant cultivars than susceptible cultivars, upon inoculation by *Fusarium* and the differences were prominent in root and stem tissue than in the leaf tissues. In pigeonpea, increased PPO activity proves the fact Asha and Maruthi's resistance is partially conferred by higher induction of polyphenol oxidase (up to 52% over the susceptible varieties) upon *Fusarium* infection. Similar results in chickpea were recorded by Raju et al. [30] who reported higher PPO activities in resistant cultivar ICCV 10 upon infection. The induced PPO-2 isoform in roots and shoots treated with salicylic acid, spermidine and the pathogen have also been implicated in defence responses against the pathogen invasion. However, no induction of PPO was observed in susceptible cultivars. The present results conform Raju et al. [30] where no induction was noticed in susceptible cultivars in their root, shoot and leaves at severe disease stage (S3) upon inoculation. Studies conducted by Constabel et al. [31] and Stewart et al. [32] showed that plant PPOs are induced in response to mechanical wounding, fungal and bacterial infection, and by treatment with signalling molecules such as jasmonic acid/methyl jasmonate (MeJA), systemin and salicylic acid. The PPO over-expressing transgenic tomato plants exhibited high resistance to *Pseudomonas syringae*, the causative agent of speck disease compared to control plants [33]. Localized inoculation of tomato leaflets with *P. syringae* induces a significant increase in PPO activity and leads to systemic resistance to the subsequent infection by *P. syringae* [34]. Thipyapong and Stiffens [35] also reported the PPO catalyzed phenolic oxidation in limiting disease development. Therefore, induction of PPO and PO is quite likely to govern some mechanism in biochemical resistance in resistant cultivars that arose from interaction between *Fusarium* and host plants. However, in un-inoculated condition, no significant changes were observed in PPO level between resistant and susceptible cultivars but significant changes in induction of enzyme from S1 through S3 level were observed in all varieties. This establishes the fact that enzymes are induced at high amount only after pathogen infection and disease initiation. Another important observation was that PPO induction was more in stem tissues, whereas peroxidase induction was higher in leaf tissue. Raju et al. [30] also reported similar spatial distribution of PO and PPO activity in chickpea in response to *F. oxysporum* f. sp. *ciceri* infection.

Catalase induction started at S2 in un-inoculated control plants. Increase in induction was significant in leaf tissues upon infection as compared to controls and it continued up to S3 stage. These results match with Subramanian et al. [36] and El-Khallal [37] who reported that level of H₂O₂ and other enzyme activities increased to the level of tolerance or susceptibility to *Fusarium* wilt. Susceptible cv. BG 256 showed increase in enzymatic activities by 30% over the resistant varieties at S3 stage in leaves after infection. This might be due to the invasion of the pathogen, the host cell wall degrades with the release of particular enzyme at the vicinity of the infection site.

4.2 Spatial and temporal activity of phenyl alanine ammonia lyase

PAL activity increased in response to pathogen treatments and resembles observations on accumulation of phenolics and PAL by Arfaoui et al. [5] in chickpea seedlings treated with *Rhizobium* Pch43 followed by *F. oxysporum* f. sp. *ciceri* race 0. The chickpea seedlings exposed to cell wall protein of *F. oxysporum* f. sp. *ciceri* showed enhanced synthesis of phenols, pathogenesis-related proteins and activities of PAL and peroxidase relative to water treated controls [38]. Stadnik and Buchenauer [39] also reported the enhancement of PAL activity and accumulation of cell wall-bound phenolic compounds in wheat plants treated with BTH, a novel systemic acquired resistance (SAR) inducer in response to powdery mildew infection. Our study also indicates a rapid increase of PAL activity in pathogen treated pigeonpea seedlings, except for the control treatment. Dixon et al. [40] observed an elicitor dose-response effect (isolated from *Colletotrichum lindemuthianum*) in the activity of PAL in bean cells suspension cultures and two maxima of the enzyme activity in response to the application of smaller doses of elicitors.

Invasion of root tissues by the pathogen resulted in decreased activity of PAL in susceptible cultivar, whereas, increased activity of PAL in resistant cultivar might have prevented the fungal invasion and

thus the activity maintained at higher levels during the experimental period. Increased accumulation of PAL with pathogen could re-establish the notion that PAL is synthesized more rapidly in resistant cultivars than that in susceptible cultivars in response to invasion of pathogen.

4.3 Spatial and temporal activity of Chitinase and β -1,3-glucanase

Depolymerisation of cell wall by the combined activities of chitinase and glucanase is reported to kill fungi *in vitro* [28, 29] and cell wall fragments are shown to induce defence reactions in plants by switching on genes responsible for the synthesis of pathogenesis-related proteins [41]. Inhibition of growth of several fungi requires the presence of chitinase and β -1,3-glucanase activities [38]. Ferraris et al. [42] found that infection with *F. oxysporum* f. sp. *lycopersici* caused several fold increases in chitinases, β -1,3-glucanases, glucosidases and N-acetyl-glucosaminidases activities in susceptible varieties and resistant tomato cultivars from 5 to 90 days after inoculation. The present study also proved that chitinase activity was increasing 3.08 and 2.33 fold in root tissue, 2.45 fold and 3.43 fold in stem tissue, 3.15 fold and 3.5 fold in leaf tissues of pigeonpea resistant cultivars.

β -1,3-glucanase activity increased from 1.27-17 fold in the infected resistant cultivars as compared to 1.0-1.5 fold infected susceptible cultivars at disease initiation stage which conforms to Singh et al. [43]. Xue et al. [25] observed relatively higher activity of chitinase in the cotyledons of bean plants than hypocotyls and a significant increase in all cellular fractions of peroxidase, β -1,3-glucanases and chitinase compared with diseased and control plants and glucanase activity was also reported to increase up to 18 fold in induced bean hypocotyls tissues. Similar results were found in the present study where the induction of chitinase activity was lower in root tissues. Induction of β -1,3-glucanase increased up to 17 fold in resistant pigeonpea cultivars. Similar increases, but not as pronounced have been reported from other studies using incompatibility interactions between soybean and *Phytophthora megasperma* f. sp. *glyciensi* [44], between bean cultivars and *C. lindemuthianum* [45] and between *Vigna unguiculata* (L. Walp.) and *Fusarium oxysporum* [46]. Present study showed that chitinase significantly accumulated in different tissues after the onset of diseases. Igratius et al. [47] observed that seed associated chitinases of barley are different from those that found in leaves infected with powdery mildew, while Anuratha et al. [27] isolated an infection related chitinase transcript that was only induced upon infection of rice with the sheath blight pathogen *R. solani*. Both antifungal hydrolases are induced in coordination with other PR proteins in typical systemic acquired resistance responses [48].

Increased activity of defense enzymes in pigeonpea tissues after inoculation with *Fud* is in conformity with Cachinero et al. [49] where defence responses were induced in both pre-induced and non-induced plants infected by pathogens. They concluded that the suppression of *Fusarium* wilt possibly involved an inhibitory effect of pre-induced as well as in induced plant defences on the pathogen. Present work also reveals that in pigeonpea resistant cultivars Asha and Maruthi, the increase in β -1,3-glucanase activity is comparatively lower in root tissue than in leaf tissue. The increase in activity of resistant cultivars was 1.87 and 2.5 fold, respectively, in Asha and Maruthi, whereas in susceptible cultivars, Type 7 and Bahar, it was 1.3 and 1.13 fold, respectively, in root tissue under challenged condition (Fig. 6). Giri et al. [10] also recorded similar lower activity of β -1,3-glucanase in roots of resistant cultivars of chickpea. Benhamou et al. [50] observed accumulation of these enzymes to be faster in incompatible interactions of tomato and *F. oxysporum* f. sp. *lycopersici* or *F. radicle-lycopersici* than in a compatible interaction. In the present study, lower activity of β -1,3-glucanase was observed in resistant variety of pigeonpea in the stem tissue as compared to the susceptible cultivar at S2 stage. Beckman and Roberts [51] in their model for host pathogen interaction suggested callose deposition and lignifications as one of the plants responses against wilt disease and rate of these processes determines the degree of reduction of pathogen invasion. Beffa et al. [52] demonstrated in tobacco that, when expression of β -1,3-glucanase is specifically blocked by the antisense mRNA technique, callose deposits are protected from degradation, resulting in resistance to viral infection. Levels of chitinase and β -1,3-glucanases were also reduced when the infection was confined at S3 stage. The observed decrease in β -1,3-glucanase activity may be associated with the reduction in the pathogen invasion as a result of defence reaction. In the current study, leaf tissue of pigeonpea resistant cv. Maruthi showed 17 fold increase in comparison to control at S2 stage and lower proportionate enhancement was found in root and stem tissue. Moreover, these two enzymes act synergistically in the partial degradation of fungal cell walls. The combination of these two enzymes could strongly inhibit growth of many fungi, including those that could not be inhibited by chitinase or β -1,3-glucanase alone [53]. Chitinases in combination with β -1,3-glucanase protect the

plants from fungal infection by their lytic actions on fungal cell walls or by releasing oligosaccharides signal molecules that can activate an array of plant defences.

The resistant cultivars showed higher peroxidase activities (2-5 fold more) than the susceptible cultivars. In root tissue of Maruthi, 25% increase in peroxidase activity over the control suggests that probably this enzyme strengthens the defence response by enhancing lignifications of cell walls. The induction of PPO activities increased only after the confrontation with the pathogen, suggesting that PPO acts by phenol oxidation. Raju et al. [30] screened different genotypes of chickpea for the induction of PPO isoforms and found that the resistant genotypes against wilt disease expressed multiple isoforms of PPO, while the susceptible genotypes did not. Resistant cv. Maruthi showed 1.5 fold increases in activity at disease initiation stage in the stem tissue, whereas susceptible cv. Type 7 and Bahar showed 8-10% increase upon infection. One interesting finding was that there was almost no change in activity of PPO in Bahar roots at S2 after infection started, supporting the view that susceptible cultivars could not withstand the challenge. On the other hand, catalases implicated in several plant defence mechanism, mainly in the generation of reactive oxygen species have also been associated with induced resistance response. The results presented here indicate that all the three enzymes activities increased in different tissues of host plants after inoculation with *Fud* but catalase activity was pronounced up to S3 stage in contrary to other enzymes. In the leaf tissue of Asha (resistant) and Bahar (susceptible), the same level of activity enhancement was observed. Surprisingly, there was no significant change in Maruthi leaf tissue at S3 stage compared to other varieties, though in the S2 stage, the induction was more in this cultivar compared to the susceptible ones. These effects can be associated with the reason that the susceptible varieties express multiple isoforms of catalase, whereas the resistant genotypes did not.

Priming resistance by inducing these genes could be an efficient and inexpensive way of achieving the control of *Fusarium* wilt in pigeonpea. In case of pigeonpea resistant cultivar Maruthi, the increase was 71.6% in root tissue at S2 stage. In contrast, susceptible cultivar Type 7 did not show significant changes in the stem and leaf tissues in the disease initiation stage. Several hypotheses can be formulated to explain the level of phenolics [6] as well as PAL observed in the plants following *Fud* infection. It may result from the *de novo* mRNA synthesis or accumulation of defence related genes such as PAL. Accumulation of phenolics may also occur, upon challenge by *Fusarium*, from the release of phytoalexins from their preformed conjugated forms. Furthermore, it is likely that these two mechanisms generating phytoalexins can act synergistically. This is very important for future studies focussing on the isolation and transfer of disease resistance genes in the field, as part of an integrated management strategy to protect the plants from further pathogen attack.

4. CONCLUSION

The present results demonstrate that resistant cultivars resulted in a significant increase in chitinase and β -1,3-glucanase and the expression of defence related genes encoding PR proteins can be used as markers for the establishment of resistance. These two enzymes act synergistically on the fungal cell wall resulting in degradation and loss of inner content of cell and indicate the possibilities of their involvement in defence response in pigeonpea against *Fusarium* wilt. A direct role for β -1,3-glucanase in defence of plants against pathogens may be proposed because the substrates for these enzymes are major components of cell walls of many fungi. Induced resistance is multi-component and it is necessary to investigate further other mechanisms involved, either alone or collectively.

REFERENCES

1. FAO. <http://faostat.fao.org/faostat/collections?subset=agriculture>. Last updated Feb. 2010.
2. Nene YL, Sheila VK, Sharma SB (1989) A world list of chickpea and pigeon pea Pathogens. ICRISAT Legume Pathology Progress Report 7. Patancheru, India.
3. Heller J, Tudzynski P. Reactive oxygen species in phytopathogenic fungi: signaling, development, and disease. *Annu Rev Phytopathol* 2011;49:369-390.
4. van Loon LC, Rep M, Pieterse CM. Significance of inducible defense related proteins in infected plants. *Annu Rev Phytopathol* 2006;44:135-162.
5. Arfaoui AE, Hadrami Y, Mabrouk B, Sifi A, Boudabus I, Hadrami EI, Daayf F, Cherif M. A treatment of chickpea with *Rhizobium* isolates enhances the expression of phenylpropanoid defense-related

- 491 genes in response to infection by *Fusarium oxysporum* f. sp. *ciceris*. Plant Physiol Biochem
492 2007;45:470-479.
- 493 6. Datta J, Lal N. Temporal and spatial changes in phenolic compounds in response to *Fusarium* wilt
494 in chickpea and pigeonpea. Cell Mol Biol 2012;58:96-102.
- 495 7. Broetto F, Marchese JA, Leonardo M, Regina M. Fungal elicitor-mediated changes in polyamine
496 content, phenylalanine ammonia-lyase and peroxidase activities in bean cell culture. Gen Appl Plant
497 Physiol 2005;31:235-246.
- 498 8. Veronese, P, Ruiz, MT, Coca, MA, Lee, H, Ibeas, JI, Damz, B, Pardo, JM, Hasegawa, PM,
499 Bressan, RA, Narasimhan, ML. In defense against pathogen: both plant sentinels and foot soldiers
500 need to know the enemy. Plant Physiol 2003;131:1580-1590.
- 501 9. Ramachandra KK, Vasanthi NS, Shetty HS. Induction of β -1,3-glucanase in seedlings of pearl
502 millet in response to infection by *Sclerospora graminicola*. Eur J Plant Pathol 2000;106:267-274.
- 503 10. Giri P, Harsulkar AM, Patankar AG, Gupta VS, Sainani MN, Deshpande VV, Ranjekar PK.
504 Association of induction of protease and chitinase in chickpea roots with resistance to *Fusarium*
505 *oxysporum* f. sp. *ciceri*. Plant Pathol 1998;47:693-699.
- 506 11. Stevenson PC, Turner HC, Haware MP. Phytoalexin accumulation in the roots of chickpea (*Cicer*
507 *arietinum* L.) seedlings associated with resistance to *Fusarium* wilt (*Fusarium oxysporum* f. sp.
508 *ciceris*). Physiol Mol Plant Pathol 1997;50:167-178.
- 509 12. Oliveira JTA, Barreto ALH, Vasconcelos IM, Eloy YRG, Gondim DMF, Fernandes CF, Freire-Filho
510 FR. Role of Antioxidant Enzymes, Hydrogen Peroxide and PR-Proteins in the Compatible and
511 Incompatible Interactions of Cowpea (*Vigna unguiculata*) Genotypes with the Fungus *Colletotrichum*
512 *gloeosporioides*. J Plant Physiol Pathol 2014;2:3. <http://dx.doi.org/10.4172/2329-955X.1000131>
- 513 13. Pareek SS, Ravi I, Sharma V. Induction of β -1,3-glucanase and chitinase in *Vigna aconitifolia*
514 inoculated with *Macrophomina phaseolina*. J Plant Interact 2014;9:434-439, DOI:
515 10.1080/17429145.2013.849362
- 516 14. Gharbi Y, Barkallah M, Bouazizi E, Hibar K, Gdoura R, Triki MA. Lignification, phenols
517 accumulation, induction of PR proteins and antioxidant-related enzymes are key factors in the
518 resistance of *Olea europaea* to *Verticillium* wilt of olive. Acta Physiol Plant 2017;39:43.
519 <https://doi.org/10.1007/s11738-016-2343-z>
- 520 15. Tullu A, Muehlbauer FJ, Siomn CJ, Mayer MS, Kumar J, Kaiser WJ, Draft JM. Inheritance and
521 linkage of a gene for resistance to race 4 of *Fusarium* wilt and RAPD markers in chickpea. Euphytica
522 1988;102:227-232.
- 523 16. Kar M, Mishra D. Catalase, Peroxidase and Polyphenoloxidase activities during Rice leaf
524 senescence. Plant Physiol 1976;57:315-319.
- 525 17. Braber JM. Catalase and peroxidase in primary bean leaves during development and
526 senescence. Z Pflanzenphysiol 1980;97:135-144.
- 527 18. Zucker M. Induction of Phenylalanine ammonia-lyase by light and its relation to chlorogenic acid
528 synthesis in potato tuber tissue. Plant Physiol 1965;40:779-784.
- 529 19. Chen A, Mayor R, Deloach J. Purification and characterization of chitinase from a stable fly
530 *Stomoxys calcitrans*. Arch Biochem Biophys 1982;216:314-321.
- 531 20. Tsukamoto T, Koga D, Ide A, Ishibashi T, Horinomatsoslig M, Yagishita K, Imoto T. Purification
532 and some properties of chitinase from yam *Dioscorea opposita* Thumb. Agr Biol Chem 1984;48:227-
533 232.
- 534 21. Reissig JL, Strominger JL, Leloir LF. A modified colorimetric method for the estimation of *N*-
535 acetylamino sugars. J Biol Chem 1955;217:959-966.
- 536 22. Koga D, Sueshige M, Orikono K, Utsmi T, Tanaka S, Yamada T, Ede A. Efficiency of chitinolytic
537 enzymes in the formation of *Trichoderma matsutake* protoplasts. Agr Biol Chem 1988;52:2091-2093.
- 538 23. Somogyi M. Notes on sugar determination. J Biol Chem 1952;195:19-23.
- 539 24. Rathi YPS, Bhatt A, Singh US. Biochemical changes in pigeonpea (*Cajanus cajan* (L.) Millsp.)
540 leaves in relation to resistance against sterility mosaic disease. J Biosci 1986;10:467-474.
- 541 25. Xue L, Charest LM, Jabaji Hare SH. Systemic Induction of peroxidases, 1,3- β -glucanases,
542 chitinases, and resistance in bean plants by binucleate *Rhizoctonia* species. Phytopathology
543 1988;88:359-365.
- 544 26. El-Argawy E, Adss IA. Quantitative gene expression of Peroxidase, Polyphenol oxidase and
545 Catalase as molecular markers for resistance against *Ralstonia solanacearum*. Amer J Mol Biol
546 2016;6:88-100. 33.
- 547 27. Anuratha CS, Zen KC, Cole KC, Mew T, Muthukrishnan S. Induction of chitinases and β -1, 3-
548 glucanases in *Rhizoctonia solani*-infected rice plants: Isolation of an infection-related chitinases DNA
549 clone. Plant Physiol 1996;97:39-46.

- 550 28. Mauch F, Mauch MB, Boller T. Antifungal hydrolyses in pea tissue. I. Purification and
551 Characterization of Two Chitinases and Two β -1,3-glucanases differentially Regulated during
552 Development and in Response to fungal Infection. *Plant Physiol* 1988;87:325-333.
- 553 29. Mauch F, Mauch MB, Boller T. Antifungal hydrolyses in pea tissue. II. Inhibition of fungal growth
554 by combinations of chitinase and β -1,3-glucanases. *Plant Physiol* 1988;87:448-453.
- 555 30. Raju S, Jayalakshmi SK, Sreeramulu K. Comparative study on the induction of defense related
556 enzymes in two different cultivars of chickpea (*Cicer arietinum* L.) genotypes by salicylic acid,
557 spermine and *Fusarium oxysporum* f. sp. *ciceris*. *Aust J Crop Sci* 2008;2:121-140.
- 558 31. Constabel CP, Yip L, Patton JJ, Christopher ME. Polyphenol oxidase from hybrid poplar, cloning
559 and expression in response to wounding and herbivory. *Plant Pathol* 2000;124:285-295.
- 560 32. Stewart RJ, Sawyeer BJB, Bucheli CS, Robinson SP. Polyphenol oxidase is induced by chilling
561 and wounding in pineapple. *Aust J Plant Physiol* 2001;28:181-191.
- 562 33. Li L, Stiffens JC. Over expression of polyphenol oxidase in transgenic tomato plants results in
563 enhanced bacterial disease resistance. *Planta* 2002;215:239-247.
- 564 34. Stout MJ, Fidantsef AL, Duffey SS, Bostock RM. Signal interactions in pathogen and insect attack:
565 systemic plant-mediated interactions between pathogens and herbivores of the tomato, *Lycopersicon*
566 *esculentum*. *Physiol Mol Plant Pathol* 1999;54:115-130.
- 567 35. Thipyapong P, Stiffens JC. Tomato Polyphenol oxidase differential response of the PPO F
568 promoter to injuries and wound signals. *Plant Physiol* 1997;115:409-418.
- 569 36. Subramanian S, Moziah M, Sariah M, Puad MP, Xavier R. Bioassay method for testing *Fusarium*
570 wilt disease tolerance in transgenic banana. *Sci Hortic* 2006;108:378-389.
- 571 37. El-Khalal SM. Induction and modulation of resistance in Tomato plants against *Fusarium* wilt
572 disease by bioagent fungi (Arbuscular Mycorrhiza) and/or hormonal elicitors (Jasmonic Acid and
573 Salicylic Acid): 2- Changes in the antioxidant enzymes, phenolic compounds and pathogenesis-
574 related proteins. *Aust J Basic & Appl Sci* 2007;1:717-732.
- 575 38. Saikia R, Singh BP, Kumar R, Arora DK. Detection of pathogenesis-related proteins- chitinase and
576 β -1,3-glucanase in induced chickpea. *Curr Sci* 2005;89:659-663.
- 577 39. Stadnik MJ, Buchenauer H. Inhibition of phenylalanine ammonia-lyase suppresses the resistance
578 induced by benzothiadazole in wheat *Blumeia graminis* f. sp. *tritici*. *Physiol Mol Plant Pathol*
579 2000;57:25-34.
- 580 40. Dixon RA, Dey PM, Murphy DL, Whitehead IM. Dose responses for *Colletotrichum*
581 *lindemuthianum* elicitor-mediated enzyme induction in French bean cell suspension cultures. *Planta*
582 1981;151:272-280.
- 583 41. Ryan CA, Farmer EE. Oligonucleotide signals in plants: A current assessment. *Annu Rev Plant*
584 *Physiol Plant Mol Biol* 1991;42:651-674.
- 585 42. Ferraris L, Abbattista Gentile I, Matta A. Variations of phenols concentration as a consequence of
586 stresses that induce resistance to *Fusarium* wilt of tomato. *Z Pflanzenk Pflanzen* 1987;94:624-629.
- 587 43. Singh R, Sindhu A, Singhal HR. Biochemical basis of resistance in chickpea (*Cicer arietinum* L.)
588 against *Fusarium* wilt. *Acta Phytopathol Entomol Hung* 2003;38:13-19.
- 589 44. Yi SY, Hwang BK. Differential induction and accumulation of β -1,3-glucanase and chitinase
590 isoforms in soybean hypocotyls and leaves after compatible and incompatible infection with
591 *Phytophthora megasperma* f. sp. *glycinea*. *Physiol Mol Plant Pathol* 1996;48:200-207.
- 592 45. Daugrois JH, Lafitte C, Barthe JP, Touze A. Induction of 1,3- β -glucanase and chitinase activity in
593 compatible and incompatible interactions between *Colletotrichum lindemuthianum* and bean cultivars.
594 *J Phytopathol* 1990;130:225-231.
- 595 46. Vieira FA, Carvalho AO, Vitoria AP, Retamal CA, Gomes VM. Differential expression of defense-
596 related proteins in *Vigna unguiculata* (L. Walp.) seedlings after infection with *Fusarium oxysporum*.
597 *Crop Prot* 2012;29:440-447.
- 598 47. Igratius SMJ, Chopra RK, Muthukrishnan S. Effects of fungal infection and wounding on the
599 expression of chitinases and β -1,3-glucanases in near-isogenic lines of barley. *Physiol Plant*
600 1994;90:584-592.
- 601 48. Ryals JA, Neuenschwander UH, Willits MG, Molina A, Steiner HY, Hunt MD. Systemic acquired
602 resistance. *The Plant Cell* 1996;8:1809-1819.
- 603 49. Cachinero JM, Hervas A, Jimenez-Diaz RM, Tena M. Plant defense reactions against *Fusarium*
604 wilt in chickpea induced by incompatible race 0 of *Fusarium oxysporum* f. sp. *ciceri* and non host
605 isolates of *F. oxysporum*. *Plant Pathol* 2002;51:765-776.
- 606 50. Benhamou N, Joosten MHAJ, de Wit PJG. Subcellular localization of chitinase and of its potential
607 substrate in tomato root tissues infected by *Fusarium oxysporum* f. sp. *radices-lycopersici*. *Plant*
608 *Physiol* 1990;97:1494-1500.

- 609 51. Beckman CH, Roberts EM. On the nature and genetic basis for resistance and tolerance to fungal
610 wilt diseases of plants. *Adv Bot Res* 1995;21:36-72.
- 611 52. Beffa RS, Hofer RM, Thomas M, Meins F Jr. Decreased susceptibility to viral disease of β -1, 3-
612 glucanase deficient plants generated by antisense transformation. *The Plant Cell* 1996;8:1001-1011.
- 613 53. Vogeli U, Meins F, Boller T. Coordinated regulation of chitinase and β -1,3-glucanase in bean
614 leaves. *Planta* 1988;174:364-372.