

Original Research Article

A comparative study of High Sensitive C-Reactive Protein (hs-CRP) and urinary calcium/creatinine ratio in normal pregnancy and pregnancy induced hypertension in Gujarat, India

ABSTRACT

Aim To measure & correlate serum hs-CRP and urinary calcium creatinine ratio in between pregnancy induced hypertension and normal pregnancy.

Material & Methods ~~The study was approved by local institutional ethics committee.~~ It was conducted in total 150 subjects attending Department of Obstetrics & Gynecology at Sir Takhtsinhji Hospital, Bhavnagar, Gujarat, which were categorized in two groups. Group A: 75 patients of pregnancy induced hypertension; Group B: 75 normal pregnant women as a control group. They were primarily diagnosed based on measurement of blood pressure, clinical examination followed by biochemical investigations like hs-CRP, urinary calcium & creatinine, RBS, urea, creatinine, uric acid, CK-MB. Written informed consent was obtained from all participants before enrolling in the study.

Results No significant change in the levels of biological parameters like RBS, urea, creatinine and CK-MB was observed in both the study groups ($p > 0.05$) except serum uric acid shows significant difference ($p < 0.001$). The levels of hs-CRP were found to be significantly higher in patients of pregnancy induced hypertension (7.81 ± 3.681) as compared to the normal pregnant women (3.11 ± 1.972) ($p < 0.0001$). ~~This difference was statistically highly significant.~~ UCa/Cr ratio had more sensitivity and specificity than hs-CRP for prediction of pregnancy induced hypertension compared to normal pregnancy.

Conclusion UCa/Cr ratio can be used as a more sensitive and specific test to detect pregnancy induced hypertension.

Key Words *hs-CRP, urinary calcium creatinine ratio, pregnancy induced hypertension*

27 INTRODUCTION

28 Pregnancy induced hypertension (PIH) is one of the most common leading cause of maternal
29 and perinatal morbidity and mortality in females & it complicates approximately one out of every ten
30 pregnancies. Early detection of PIH is more important to save mother and fetal life. PIH is associated
31 with increased risks of serious outcomes such as premature delivery, intra uterine growth retardation
32 (IUGR), perinatal mortality & morbidity, acute renal failure, acute hepatic failure, intra-partum
33 haemorrhage, post-partum haemorrhage, maternal mortality & morbidity following complication such
34 as haemolysis (H), elevated liver enzymes (EL), low platelets (LP) (HELLP syndrome), disseminated
35 intravascular coagulation (DIC) and seizures.^[1,2,3,4]

36 Pre-eclampsia (PE) is most common complication of pregnancy induced hypertension. The
37 incidence of PE in hospital practice varies widely from 5 to 15% in India. The incidence of PE in
38 primigravidae is about 10% and in multigravidae 5%. The etiology of endothelial dysfunction in PE is
39 not known, but it has been postulated to be part of an exaggerated maternal inflammatory response to
40 pregnancy.^[5,6,7,8]

41 There has been an increasing interest for the detection of pregnancy disorders before the
42 symptoms actually occur. Endothelial dysfunction is accompanied by elevated levels of inflammatory
43 markers. Indeed, such marker levels have been shown to be much higher in women with PE
44 compared to normal pregnant females.^[9]

45 Serum high sensitive C-reactive protein (hs-CRP) is one of the suitable markers for low grade
46 inflammation evaluation. This marker rises subsequent to stress which are accompanied by
47 endothelial dysfunction and lead to peripheral vascular remodelling, decreased compliance and
48 vascular stiffness. Inflammation is considered to have a crucial role in pathophysiologic mechanism of
49 PE and endothelial dysfunction is accompanied by elevated levels of inflammatory markers.^[10]

50 Urinary calcium creatinine ratio (UCa/Cr) is a valuable marker for prediction of PIH.
51 Physiological changes during normal pregnancy leads to increase in glomerular filtration rate those
52 results in increasing of creatinine clearance and calcium secretion in urine. But in women developing
53 PE because of vasospasm and decrease of renal flow, creatinine clearance decreases, followed by
54 increase in serum creatinine level.^[12]

55 Anil bargale et al. concluded that there was significantly increased level of serum hs-CRP in
56 PE patients as compared to normal pregnant women. They also found positive significant correlation
57 between hs-CRP levels and blood pressure.^[13]

58 Kanak kanti mandal et al. observed that the level of serum hs-CRP was higher in
59 preeclamptic patients than the normal pregnant women & in preeclamptic patients the cause of
60 increased hs-CRP value may be due to superimposed infection along with PE.^[14]

61 N.V. Lakshmi et al. reported that urinary calcium creatinine ratio was used in prediction of
62 pregnancy induced hypertension & there was significant lowering of urinary excretion of calcium in
63 cases who developed PIH. PIH developed more in cases with $UCa/Cr < 0.04$.^[1]

64 Amandeep Kaur et al. demonstrated that urinary calcium excretion in preeclamptic females
65 was significantly lower than that observed in case of age matched normal pregnant females and
66 normal non pregnant females which rules out the possibility of decreased resorption of calcium at the
67 level of renal tubules, while urinary creatinine excretion in normal pregnant females was found to be
68 more than that observed in case of normal non-pregnant females, which were responsible for the
69 decreased urinary calcium: urinary creatinine ratio in PE compared to normal pregnancies.^[5]

70 Keeping in view the different outcomes of the researchers, pregnancy induced hypertension
71 remains most significant and intriguing unsolved problem in obstetrics. In light of the above facts the
72 present study is designed to determine the level of serum hs-CRP and UCa/Cr ratio and to correlate
73 these parameters between normal pregnancy and PIH.

74

75 **Objectives**

- 76 • To compare serum high sensitive C-reactive protein level, urinary calcium and urinary creatinine
77 level in normal pregnancy and pregnancy induced hypertension.
- 78 • To compare random blood sugar, urea, creatinine, uric acid, creatine kinase-MB level in normal
79 pregnancy and pregnancy induced hypertension.
- 80 • To study the correlation of high sensitive C-reactive protein, urinary calcium/creatinine ratio in
81 between normal pregnancy and pregnancy induced hypertension.

82

MATERIAL & METHODS

The present study was conducted at Department of Biochemistry, Government Medical College and Sir Takhtsinhji General Hospital, Bhavnagar in which 75 cases of pregnancy induced hypertension and 75 normal pregnant women with normal blood pressure as a control group were included. Informed consent was taken from them. All cases were admitted to the Gopnath maternity ward (obstetric ward) of Sir T. General Hospital, Bhavnagar. They were primarily diagnosed for pregnancy induced hypertension by clinical examination followed by Biochemical investigations. The subjects in control group were selected by measuring blood pressure and inclusion-exclusion criteria were followed.

Case Group - Patients of pregnancy induced hypertension = 75

Control Group - Normal pregnant women = 75

Inclusion criteria: (For cases)

- Pregnant women with gestational age >20 week.
- Patient is a case of hypertension -
Pre-eclampsia, eclampsia and gestational hypertension (systolic blood pressure greater than 140 mm Hg or diastolic blood pressure 90 mm Hg)
- Singleton pregnancy

Exclusion criteria: (For both Groups)

- Not willing to participate in study
- Rheumatoid arthritis/Cardiovascular disease/Renal disorder/Diabetes mellitus
- Any recent or present fever or infectious disease
- Multiple pregnancy
- Smoking
- Any vaginal bleeding
- Previous history of PIH/Chronic hypertension

Laboratory Investigations

Following laboratory investigations were done in study group and control group. Venous blood sample and urine sample will be collected respectively as given below and analyzed on fully auto analyzer ILAB-650 (Instrumentation Laboratory, Italy) at NABL accredited Clinical Biochemistry Section, Laboratory Services Sir T. Hospital, Bhavnagar.

114 From Venous blood:

- 115 • high sensitive C-reactive protein
- 116 • Random blood sugar
- 117 • Blood urea
- 118 • Serum creatinine
- 119 • Uric acid
- 120 • Creatine kinase - MB

121 From Urine Sample

- 122 • Urinary calcium
- 123 • Urinary creatinine

124
125 For all investigations, SOPs were used. Like estimation of high sensitive c- reactive protein (HS-CRP)
126 by turbidimetric immunoassay method ^[15] urinary calcium by modified arsenazo method ^[16] creatinine
127 by Modified Jaffe's Kinetic method ^[17] urea estimation by urease-gldh kinetic method, uric acid by
128 uricase, POD method, CK-MB estimation by immuno-inhibition method ^[18,19].

129 Data were analyzed by using GraphPad instat version 3.0. In data analysis, comparison of these
130 parameters were carried out by applying unpaired t-test and Fisher's exact test. Interpretation was
131 done according to p-value. *p < 0.05 - significant and **p < 0.001 - highly significant. All the subjects
132 were divided in following groups:

133 Group A: Patients of pregnancy induced hypertension (Case group)

134 Group B: Normal pregnant women who volunteered to participate in the present study. (Control
135 group).

136

137

RESULTS

The present study included 75 patients of pregnancy induced hypertension and 75 normal pregnant women with normal blood pressure as a control group in which hs-CRP and urinary calcium, urinary creatinine, urinary calcium creatinine ratio were measured on fully automated chemistry analyzer ILab-650 (Instrumentation Laboratory, USA).

Mean $\pm$ SD of age among cases was 24.65 ± 4.180 years which include 13 patients (17%) ≤ 20 years, 39 patients (52%) between 21-25 years, 17 patients (25%) between 26-30 years and 6 patients (8%) ≥ 31 years of age. The maximum number of patients belongs to age group between 21-25 years. In controls Mean $\pm$ SD of age was 24.52 ± 4.078 years which include 10 normal pregnant women (13%) ≤ 20 years, 45 normal pregnant women (60%) between 21-25 years, 16 normal pregnant women (21%) between 26-30 years and 4 normal pregnant women (5%) ≥ 31 years of age. The maximum number of normal pregnant women belongs to age group between 21-25 years. There was no statistically significant difference of age between two groups.

Table 1 shows the mean level of systolic blood pressure was 153.89 ± 12.512 in Group A and 113.81 ± 7.599 in Group B, while diastolic blood pressure level was 97.14 ± 9.799 in Group A and 73.70 ± 6.768 in Group B. In patients of PIH has significantly higher level of systolic and diastolic blood pressure than normal pregnant women ($p < 0.0001$), difference between them were statistically highly significant ($p < 0.0001$) where, comparison of two groups was done by unpaired t-test.

As per Table 2, the other parameter concern only serum uric acid shows significant difference ($p < 0.001$) in between two groups where as other biochemical parameters like RBS, urea, creatinine and CK-MB did not show significant difference in their values ($p > 0.05$) in both the groups (Ref. Table No.4). As evident from table 5 mean and SD of serum hs-CRP was 7.81 ± 3.681 in group A and 3.11 ± 1.972 in Group B, difference between them were statistically highly significant ($p < 0.0001$) where, comparison of two groups was done by unpaired t-test. Mean and SD of urinary calcium was 6.95 ± 1.703 in group A and 9.79 ± 3.370 in Group B. Patients of PIH have significantly lower urinary calcium level than normal pregnant women, difference between them were statistically highly significant ($p < 0.0001$). While mean and SD of urinary creatinine was 157.61 ± 53.159 in Group A and 103.88 ± 40.817 in Group B. Urinary creatinine level was significantly higher in patients of PIH as compared to normal pregnant women, difference between them were statistically highly significant ($p < 0.0001$) where, comparison of two groups was done by unpaired t-test. Moreover, mean and SD of

urinary calcium creatinine ratio was 0.047 ± 0.023 in group A and 0.105 ± 0.047 in Group B. Patients of PIH have significantly lower UCa/Cr ratio than normal pregnant women, difference between them were statistically highly significant ($p < 0.0001$) where, comparison of two groups was done by unpaired t-test.

Table 3 and 4 shows the Sensitivity, Specificity, Positive predictive value & Negative predictive value of serum hs-CRP and urinary calcium creatinine ratio by using Fisher's exact test. As evident from table 4 at the cutoff level of 5 mg/L for hs-CRP the sensitivity of hs-CRP in relation to pregnancy induced hypertension is 77 % and the specificity is 82 %. Positive predictive value is 81 %, while the negative predictive value is 78 %. As evident from table 4, at the cutoff level of 0.057 the sensitivity of UCa/Cr ratio in relation to pregnancy induced hypertension is 80 % and the specificity is 86 %. Positive predictive value is 85 %, while the negative predictive value is 81 %. Hence, UCa/Cr ratio had more sensitivity and specificity than hs-CRP for prediction of pregnancy induced hypertension.

TABLE 1: COMPARISON OF BLOOD PRESSURE LEVELS BETWEEN GROUP A AND GROUP B

STATISTICS	SYSTOLIC BP		DIASTOLIC BP	
	GROUP A	GROUP B	GROUP A	GROUP B
	PIH	NPW	PIH	NPW
Mean	153.89	113.81	97.14	73.70
Standard Deviation	12.512	7.599	9.799	6.768
Standard Error of Mean	1.445	0.8775	1.131	0.7815
Minimum	140	100	80	60
Maximum	200	130	130	90
Significance	$p < 0.0001$		$p < 0.0001$	

PIH- pregnancy induced hypertension, NPW-normal pregnant women

TABLE 2: COMPARISON OF BIOCHEMICAL PARAMETERS BETWEEN GROUP A AND GROUP B

Parameter	Biological Reference Interval	Group A PIH			Group B NPW			Statistical Significance
		MIN.	MAX.	MEAN $\pm$ SD	MIN.	MAX.	MEAN $\pm$ SD	
RBS	Up to 140 mg/dl	68	135	95.96 $\pm$ 16.665	66	123	92.65 $\pm$ 13.596	t=1.331 p=0.1851
Urea	15-45 mg/dl	10	42	20.88 $\pm$ 7.876	12	49	19.09 $\pm$ 6.927	t=1.475 p=0.1423
Creatinine	0.6-1.1 mg/dl	0.4	1.2	0.71 $\pm$ 0.218	0.5	1.0	0.66 $\pm$ 0.146	t=1.967 p=0.0511
UA	2.6-6.0 mg/dl	3.3	11.6	6.32 $\pm$ 1.655	2.4	7.0	4.66 $\pm$ 1.046	t=7.348 p<0.001
CK-MB	18-51 U/L	16	51	31.746 $\pm$ 8.320	20	53	34.49 $\pm$ 8.703	t=1.976 p=0.050
Urinary Calcium		3.1	10.8	6.92 $\pm$ 1.70	2.8	24.0	9.79 $\pm$ 3.37	t=8.548 p<0.0001
Urinary Creatinine		52	286	157.6 $\pm$ 53.1	44	203	103.9 $\pm$ 40.8	t=11.751 p<0.0001
UCa/Crratio		0.02	0.16	0.04 $\pm$ 0.02	0.03	0.23	0.10 $\pm$ 0.04	t=7.831 p<0.0001

*p < 0.05 - significant, **p< 0.001 - highly significant, #p $\geq$ 0.05 - not significant

TABLE-3: SERUM HS-CRP LEVELS AND UCA/CRRATIO IN GROUP A AND GROUP B

	Group A PIH (N=75)	Group B NPW (N=75)	Total (N=150)
hs-CRP(mg/l)			
≤ 5	58 (39%)	13 (9%)	71 (47%)
> 5	17 (11%)	62 (41%)	79 (53%)
UCa/Cr ratio			
≤ 0.057	60 (40%)	10 (7%)	70 (47%)
>0.057	15 (10%)	65 (43%)	80 (53%)

TABLE-4: OVERALL PREDICTIVE VALUE OF HS-CRP AND UCA/CR RATIO IN STUDY GROUPS

	HS-CRP	UCA/CR RATIO
Sensitivity	77 %	80 %
Specificity	82 %	86 %
Positive Predictive Value	81 %	85 %
Negative Predictive Value	78 %	81 %

TABLE 5: COMPARISON OF HS-CRP LEVELS BETWEEN GROUP A AND GROUP B

Statistics	hs-CRP (R.I. = 0.3 – 8.8 mg/l) (n=75)	
	Group A PIH	Group B NPW
Mean	7.81	3.11
Standard Deviation	3.681	1.972
Standard Error of Mean	0.4250	0.2277
Minimum	1.5	0.6
Maximum	18.9	9.0
Significance	p<0.0001	

DISCUSSION

Pregnancy induced hypertension (PIH) is associated with high maternofoetal mortality and morbidity in both underdeveloped and developed countries. Approximately 70% of hypertensive disorders are due to pre-eclampsia (PE). In modern obstetric practice, hypertensive disorders of pregnancy are understood to encompass a clinical spectrum of abnormalities ranging from minimal elevation in blood pressure to severe hypertension with multiorgan dysfunction. The worldwide incidence of PE is 3-4% of all pregnancies.^[20] In the United States, PE accounts for 15% to 17.6% of

maternal deaths. In India the incidence was reported as 8-10% of pregnancies being 10% in primigravida, 5% in multigravida.^[2]

While the exact causes of PE are not well understood, certain factors may increase a woman's risk of developing PIH or PE. The current hypothesis regarding the etiology of PE focuses on mal-adaptation of the immune responses and defective trophoblast invasion. Thus, an excessive maternal inflammatory response, perhaps directed against foreign fetal antigens, results in a chain of events including shallow trophoblast invasion, defective spiral artery remodeling, placental infarction and release of pro-inflammatory cytokines in the systemic circulation.^[2,9]

It has been shown that CRP is elevated in women with PE.^[21] Determination of hs-CRP has been suggested to be more sensitive than conventional measurement of CRP and provides better sensitivity in confirmation of inflammation. Recently, several studies have conducted to elucidate a relationship between PE and serum hs-CRP levels.^[22]

Renal excretion of calcium increases during pregnancy, maximum excretion levels reached during the third trimester. Urinary excretion of calcium levels in non-pregnant women is in the range about 100 - 250 mg/day, in pregnant women the range is between 350-620 mg/day. There is a decrease in urinary calcium levels in pre-eclampsia. Decreased renal filtration and increased tubular reabsorption may result in hypocalciuria.^[23]

The aetiology of hypocalciuria in preeclamptic patients is unknown. It has been speculated that hypocalciuria may result from decreased dietary intake, decreased intestinal absorption, increased calcium uptake by the fetus and placenta, or intrinsic renal tubular dysfunction.^[24]

Findings of the present study shows the mean age of study group A was 24.65 ± 4.180 years and group B was 24.52 ± 4.078 years without any significance difference between them.

The mean and SD of systolic blood pressure was high 153.89 ± 12.512 in patients of PIH and 113.81 ± 7.599 in normal pregnant women, while diastolic blood pressure was high 97.14 ± 9.799 in patients of PIH and 73.70 ± 6.768 in normal pregnant women with statistically significant difference ($p < 0.0001$) (Table-1).

As per the other parameter concern only serum uric acid shows significant difference ($p < 0.001$) in between group A and Group B where as other biochemical parameters like RBS, Urea, Creatinine and CK-MB did not show significant deference in their values ($p > 0.05$) in both the groups.

The mean and SD of serum hs-CRP was 7.81 ± 3.681 in patients of PIH and 3.11 ± 1.972 in normal

pregnant women, difference between them were statistically highly significant ($p < 0.0001$). The mean and SD of urinary calcium was 6.92 ± 1.703 in group A and 9.79 ± 3.370 in Group B. Patients of PIH have significantly lower urinary calcium level than normal pregnant women ($p < 0.0001$). While mean and SD of urinary creatinine was 157.61 ± 53.159 in Group A and 103.9 ± 40.817 in Group B. Urinary creatinine level was significantly higher in patients of PIH as compared to normal pregnant women ($p < 0.0001$). The mean and SD of urinary calcium creatinine ratio was 0.047 ± 0.023 in group A and 0.105 ± 0.047 in Group B. Patients of PIH had significantly lower UCa/Cr ratio than normal pregnant women with statistically significant difference ($p < 0.0001$) (Table-2).

The sensitivity and specificity of serum hs-CRP at the cutoff level of 5 mg/L for prediction of pregnancy induced hypertension was 77% and 82% respectively with 81% positive predictive value and 78% negative predictive value (Table-3,4). The sensitivity and specificity of UCa/Cr ratio at the cutoff level of 0.057 for prediction pregnancy induced hypertension was 80% and 86% respectively with 85% positive predictive value and 81% negative predictive value (Table-3,4).

UCa/Cr ratio had more sensitivity and specificity than hs-CRP for prediction of pregnancy induced hypertension compared to normal pregnancy (Table-4).

From above findings it is indicated that serum hs-CRP level was significantly increased and UCa/Cr ratio was significantly decreased in pregnancy induced hypertension compared to normal pregnancy. According to Vijayalakshmi et al.^[10] serum hs-CRP level significantly elevated in pre-eclampsia suggesting pre-eclampsia to be an exaggerated inflammatory condition when compared to normal pregnancy. Rashmi Sinha et al.^[24] reported that UCa/Cr ratio at 0.04 in spot urine sample being a good test for prediction of pre-eclampsia can be recommended as a screening test in all asymptomatic pregnant women. In order to justify our work, we are mentioning work of some previous researchers whose findings are supported by observation of present study.^[25-27]

CONCLUSION

In the present study it is concluded that serum hs-CRP level was significantly increased and UCa/Cr ratio was significantly decreased in pregnancy induced hypertension compared to normal pregnancy. Serum hs-CRP at the cutoff level 5 mg/L can be useful in identifying pregnant women at risk for pregnancy induced hypertension. UCa/Cr ratio < 0.057 in spot urine sample can be used as a good test for prediction of pregnancy induced hypertension. However, in patients of PIH serum hs-CRP

show less predictive value as compared to UCa/Cr ratio, therefore UCa/Cr ratio can be used as a more sensitive and specific test to detect pregnancy induced hypertension which is a non-invasive and inexpensive test.

REFERENCES

1. Dr. N.V. Lakshmi, Dr. P. Kiranmai, Dr. K. Ambika Devi and Dr. J. Rama Rao. Role of Urinary Calcium Creatinine Ratio in Prediction of Pregnancy Induced Hypertension. *Int. J Pharm Bio Sci.* 2013 July;4(3):(B)1021-1026.
2. Nicole R. Anderson, Megan Undeberg, Karen MS Bastianelli. Pregnancy-Induced Hypertension and Preeclampsia: A Review of Current Antihypertensive Pharmacologic Treatment Options. *Austin J Pharmacol Ther.* 2013;1(1):2373-6208.
3. Manjusha Sajith, Vandana Nimbargi, Amit Modi, Ronak Sumariya, Atmaram Pawar. Incidence of pregnancy induced hypertension and prescription pattern of antihypertensive drugs in pregnancy. *Apr 2014;5(4):163-170.*
4. Khosravi, S., Dabiran, S., Lotfi, M. and Asnavandy, M. Study of the Prevalence of Hypertension and Complications of Hypertensive Disorders in Pregnancy. *Open Journal of Preventive Medicine.* 2014;(4):860-867.
5. Amandeep Kaur et al., Serum and Urinary Calcium and Creatinine Levels in the 3rd Trimester of Normal Pregnant and Pre-eclamptic Females. *International Journal of Recent Scientific.* July 2015;6(7):4908-4913.
6. Nanda K, Sadanand G, Muralidhara Krishna C S, K L Mahadevappa. C - reactive protein as a predictive factor of preeclampsia. *Int J Biol Med Res.* 2012;3(1): 1307-1310.
7. Gandhi MR, Jani PS, Patel UM, Kakani CR, Thakor NC, Gupta N. Perinatal outcome in pregnancy induced hypertension cases at GMERS Medical College, Dharpur-Patan, North Gujarat region, India: a prospective study. *Int J Adv Med.* 2015;2:152-5.
8. Dutta. Text book of OBSTETRICS. Hypertensive Disorders in Pregnancy. 7th edition; p. 219-241.

9. Z Tavana, J Zolghadri, G Madadi. The Relationship between Maternal serum Highly Sensitive C - reactive protein, Leptin and Hypertensive Disorders of Pregnancy. The Internet Journal of Endocrinology. 2010;6(2);1-7.
10. Vijayalakshmi P et al. Study of serum hsCRP and lipid profile in pre-eclampsia. International Journal of Recent Trends in Science and Technology. April 2015; 14(3):605-609.
11. Samira Behboudi- Gandevari, Narges Alian Moghadam, Lida Mogadam- banaem, Bitamohamadi and Mohamad Asghari. Association of high-sensitivity C-reactive protein serum levels in early pregnancy with the severity of preeclampsia and fetal birth weight. J. Perinat. Med. 40. (2012);601–605.
12. Azita fath Negad Kazemi, Fahime shhatie, Nilofar Satarjade, Mehrangiz Ebrahimi Mameghani. The Predictive value of Urinary calcium to creatinine Ratio, Roll-over test and BMI in early diagnosis of Pre-eclampsia. Research journal of Biological sciences. 2010;5(2):183-186.
13. Anil Bargale, Jayashree V. Ganu, Dhiraj J. Trivedi, Nitin Nagane, Rakesh Mudaraddi, Aparna Sagare. Serum hs-CRP and Uric Acid as Indicator of Severity in Preeclampsia. International Journal of Pharma and Bio Sciences. July-sept 2011;2(3);B:340-345.
14. Dr. Kanak Kanti Mandal, Dr. Arpita Das, Dr. H. Lakhipyari Devi, Dr. N. Omita, Prof. N. Nabakishore Singh, Prof. W. G. Singh. Serum high sensitivity C - reactive protein as predictor of Preeclampsia. IOSR Journal of Dental and Medical Sciences. Feb 2016;15(2);Ver. XII:26-31.
15. Quantia CRP-US product insert. Tulip diagnostic. Estimation of serum hs-CRP by turbidimetric immunoassay method.
16. Calcium product insert, Agappe. Estimation of urinary calcium by Arsenazo III method.
17. Creatinine product insert, Agappe. Estimation of serum creatinine by Jaffe's kinetic method.
18. DGKC, J. Clin. Chem. Clin. Bioch. 1977;15:255.
19. Di. witt, C. Trendelenburg, J. clin. chemie, clin. Bioch. 1982;20:235.
20. Alice Wang, Sarosh Rana and S. Ananth Karumanchi. Preeclampsia: The Role of Angiogenic Factors in Its Pathogenesis. Physiology. 2009;24:147-158.

- 329 21.Mehdi Farzadnia, Hossein Ayatollahi, Maliheh Hasan-Zade, Hamid Reza Rahimi. A
330 comparative study of vascular cell adhesion molecule-1 and high-sensitive C-reactive protein
331 in normal and preeclamptic pregnancies. *Interventional Medicine & Applied*
332 *Science*.2012;4(2):26-30.
- 333 22.Hossein Ayatollahi, Maliheh Hasanzade, Mahdi Farzadnia, Mahdi Khabbaz Khoob, Atefeh
334 Rahmanian. Serum Level of High Sensitive C - reactive protein in Normal and Preeclamptic
335 Pregnancies. *Iranian Journal of Pathology*. 2007;2(3): 100-104.
- 336 23.Swapna V. S, Triveni Jambale, Jayaprakash Murthy D. Study of Urinary Calcium and Urinary
337 Creatinine Levels and Urinary Calcium/Creatinine Ratio in Gestational Hypertensive Patients.
338 *Journal of Evolution of Medical and Dental Sciences*.2015;4(53):9145-9150.
- 339 24.Rashmi Sinha, Indu Bhushan. Study of Urinary Calcium/Creatinine Ratio (CCR) In a Spot
340 Sample of Urine for Early Prediction of Preeclampsia.*IOSR Journal of Dental and Medical*
341 *Sciences*. May 2016;15(5);Ver. VIII:101-104.
- 342 25.Ifeoma Udenze & Casimir Amadi, Nicholas Awolola, Christian Chigozie Makwe. The role of
343 cytokines as inflammatory mediators in preeclampsia. *Pan African Medical*
344 *Journal*.2015;20:219. Available at [http://www.panafrican-med-](http://www.panafrican-med-journal.com/content/article/20/219/full/)
345 [journal.com/content/article/20/219/full/](http://www.panafrican-med-journal.com/content/article/20/219/full/)
- 346 26.Svetlana Duricic, Marina Stojanov, Ivana Obradovic, Aleksandar Glisic, Darko Plecas.
347 Fibronectin and C - reactive protein in Pregnancy Induced Hypertension. *Jugoslav Med*
348 *Biochem*.2003;22:325-328.
- 349 27.Menha Swellama, Nervana Samya, Susan Abdl Wahab and Mohamed Saeed Ibrahim.
350 Emerging role of endothelial and inflammatory markers in preeclampsia.*Disease Markers* 26.
351 2009:127–133.