

Effect of Pretreatment with Aqueous Leaf Extract of *Vitex doniana* on Cadmium-Induced Toxicity to Rats

ABSTRACT

Aim: To determine the effect of pretreatment with aqueous leaf extract of *Vitex doniana* on the cytotoxic response of the antioxidant defence systems in the liver and kidneys of rats exposed to a mild dose of cadmium.

Materials and methods: Sixteen Male Wistar strain rats (180-220 g b.wt) were divided into four groups: Group A served as the control and rats were exposed to and maintained on a normal rat diet and tap water throughout study. Group B was maintained on a normal rat diet and then exposed to a single daily oral dose of cadmium (3 mg CdCl₂/kg) in drinking water for five days. Groups C were pretreated with 200 mg/kg *Vitex doniana* leaf extract for fifteen days and after that received a single daily oral dose of cadmium (3 mg CdCl₂/kg) in drinking water for five days. Group D was exposed to only 200 mg/kg *Vitex doniana* leaf extract for fifteen days. At the end of the experiment membrane lipid peroxidation, glutathione contents and activities of antioxidant enzymes catalase, superoxide dismutase, and cadmium content were determined in the liver and kidney samples

Results: The results showed that pretreatment with aqueous leaf extract of *Vitex doniana* effectively countered Cd-induced membrane lipid peroxidation, depletion of the non-enzymic antioxidants, glutathione, and induction of the antioxidant enzymes catalase and superoxide dismutase in the liver and kidney as well as effectively reduced cadmium accumulation in the liver and kidney and cadmium-induced liver tissue cell injury. The protective effect of aqueous leaf extract of *Vitex doniana* against Cd-induced lipid peroxidation and tissue glutathione depletion was more pronounced in the kidney than in the liver.

Conclusion: The aqueous leaf extract of *Vitex doniana* significantly reversed Cd-induced deleterious alterations in the liver and kidney tissue of the rats. The active antioxidant principle was not determined but is presumed to be polyphenols.

Keywords: Cadmium, *Vitex doniana*, polyphenols, pretreatment, endogenous antioxidants, lipid peroxidation.

1. INTRODUCTION

Cadmium, a toxic heavy metal, a common industrial and environmental pollutant, released into the environment from natural and anthropogenic sources has been shown to manifest its toxicity in humans and animals mainly by virtue of its bioaccumulation in target tissues, stimulation of the generation of free radicals and interference with the utilization of essential metals, all of which culminate in oxidative stress [1, 2, 3, 4, 5]. Oxidative stress is a condition associated with an increased rate of cellular damage induced by an imbalance between reactive oxygen species (ROS) formation and scavenging by antioxidants [6, 7]. It has been demonstrated that Cd induces oxidative

44 stress by stimulating the production of free radicals such as hydroxyl radicals [5], superoxide anions,
45 nitric oxide and hydrogen peroxide [8], resulting in membrane lipid peroxidation and oxidative
46 deterioration of proteins and DNA and the process initiating various pathological conditions in humans
47 and animals [8].

48 Humans become exposed to Cd pollutants through the food chain and the air. Following oral
49 exposure, Cd is absorbed and delivered to the liver by endogenous intestinal protein metallothionein,
50 and from the liver, it is rapidly redistributed to other organs with the kidney as the main target organ
51 for Cd toxicity [9]. Public health interest in the toxic effects of environmental Cd was awakened by
52 epidemiological evidence linking industrial Cd waste pollution of marine food sources to the outbreak
53 of *it Itai disease* in Japan, a disease characterized by, among others, severe bone disorders and renal
54 tubular lesions [10]. Studies with experimental animals have shown that exposure to Cd results in
55 toxic lesions in many species, with Cd showing various mechanisms of toxicity in particular species
56 under different experimental conditions [8]. The severity of Cd intoxication of target organs is
57 dependent on the route, dose, and duration of exposure and it manifests in various forms ranging
58 from acute toxicosis to cancer. For this reason, it has been postulated that ingestion of antioxidant
59 could counter Cd oxidative damage to target tissue cells. The role of antioxidants is to neutralize the
60 excess of free radicals, to protect the cells from toxic effects and to contribute to disease prevention
61 [7, 11]. This has been demonstrated in experimental cadmium intoxication where certain
62 micronutrients antioxidants notably, vitamin E and selenium, have reportedly reversed or prevented
63 cadmium toxicity to target tissue cells [12, 13, 14, 15, 16, 17, 18].

64 Apart from micronutrient supplements, some medicinal plant extracts have been reported to exhibit a
65 protective role against ROS and lipid peroxidation induced by xenobiotics including environmental
66 pollutants like Cd [7, 19, 20, 21, 22]. *Vitex doniana* leaf has been in used for many decades in the
67 treatment of many illnesses, and as food in Nigeria and some part of Africa [23]. This plant is usually
68 common, relatively cheap and with promising health-boosting potentials. There is evidence that *Vitex*
69 *doniana* is an important source of polyphenols which are known to act as strong antioxidants against
70 ROS induced oxidative stress [7, 24]. The study of the biological effects of polyphenols has become
71 an area of interest in the light of recent advances in the field of nutrition and medical sciences. We
72 report on the protective effect of *Vitex doniana* leaf extract on Cd-induced oxidative damage to rat

73 hepatic and renal tissues as determined by tissue Cd bioaccumulation profiles, oxidative stress
74 biomarkers such as membrane lipid peroxidation, antioxidant defence enzymes activities, non-
75 enzymic tissue antioxidant levels, and biomarkers of cellular tissue damage.

76 2. MATERIALS AND METHODS

77 2.1 Collection of Leaf Samples and Preparation of Extract

78 Fresh leaf of *Vitex doniana* was obtained in the vicinity of the Faculty of Medical Sciences, University
79 of Jos, Jos, Plateau State, Nigeria. They were identified and authenticated by Mr Joseph in Federal
80 College of Forestry Jos, Plateau State. The leaf was washed with distilled water to remove dust
81 particles and shade dried at room temperature under continuous ventilation for two weeks, and the
82 dried leaf was pounded into a fine powder using a pestle and a mortar. Sixty grams of *Vitex doniana*
83 powder were weighed and soaked overnight in 500 ml of distilled water. The mixture was shaken on a
84 mechanical shaker for 3 hours and filtered through a Whatman No.1 filter paper. The resulting
85 aqueous extract of *Vitex doniana* was concentrated and evaporated to dryness using a Rotary
86 Evaporator at 40°C and stored in the refrigerator pending use. The extract was further reconstituted in
87 distilled water at appropriate concentration before administering to experimental animals. The protocol
88 for the treatment of rats was approved by Animal Ethical Committee of the University of Jos with
89 reference number: UJ/FPS/ F17-00379.

90 2.2 Animal Treatment

91 Wistar Strain male rats (b.wt. 180-220 g) obtained from the Animal House Unit, University of Jos,
92 were used in the study. They were maintained on a standard rat diet, 'Vital Feed' (purchased from
93 Grand Cereals and Oil Mills Ltd, Kuru, Nigeria) and tap water as drinking water, *ad libitum*. The
94 respective working doses of *Vitex doniana* extract and Cd (as CdCl₂) administered orally to
95 experimental animals in this study was first determined in a pilot study. The Cd dose used was the
96 lowest in the graded concentration of the Cd salt tolerated by the rats with quantifiable tissue
97 biochemical changes without the fatality. On the other hand, the *Vitex doniana* extract dose used was
98 the concentration that produced significant antioxidant defence capability compared to control rats fed
99 normal diet alone.

Rats were weighed and distributed evenly, 3 rats/cages into four standard plastic-metal rat cages, labeled A-D, respectively. Rats in the four groups were fed the standard 'Vital feed' rat diet and drinking water *ad libitum*. However, each rat in groups C and D received twice daily, an oral supplement of aqueous extract of *Vitex doniana* at a dose of 200 mg *Vitex doniana* extract/kg b.wt/day, administered by means of a needle-free Syringe. The daily oral *Vitex doniana* supplementation was carried out for 15 days. Thereafter, rats in group B and C were each given one single oral dose CdCl₂ in aqueous solution (3 mg CdCl₂/kg b.wt) daily for 5days.

2.3 Tissue Collection and Preparation

At the end of the feeding experiment, on day 11, rats under anaesthesia were sacrificed by decapitation and, in each case, the liver and kidneys were excised and washed in ice-cold normal saline to remove adhering blood particles. Homogenates of liver and kidney samples of each rat were prepared separately by homogenizing 1 g portion in ice-cold 50 mM Tris-HCl buffer, pH7.4 (1:10, w/v) in an Akia homogenizer. The homogenates were centrifuged at 2,400 xg for 10 min in a refrigerated low-speed centrifuge and the supernatant (S1) fractions were collected with Pasteur Pipette into plastic vials and stored at 2 °f pending biochemical analysis. The rest of the kidney and liver samples were used for determination of Cd content.

2.4 Phytochemical Analysis

Phytochemical screening of the leaves of *Vitex doniana* was done using standard method of Harborne [25].

2.5 Biochemical Analysis

Membrane lipid peroxidation, non-enzymic tissue antioxidants (ascorbic acid and glutathione) concentration and antioxidant enzymes (superoxide dismutase, glutathione peroxidase and catalase) activity were determined in the liver and kidney supernatant fractions and red blood cells. Lipid peroxidation measured by malondialdehyde (MDA) content was assayed by the thiobarbituric acid reaction according to the method described by [26]. The concentration of reduced glutathione (GSH) in the supernatant fraction of the tissue homogenates was assessed by the Ellman reaction [27] using the method described by [28] Catalase (CAT) activity was determined by measuring the rate of decomposition of hydrogen peroxide at 570 nm as described by [29]. Superoxide dismutase (SOD)

activity was assayed as described by [30]. Tissue cell injury enzyme biomarkers; serum alkaline phosphatase (ALP), alanine aminotransferases (ALT) and aspartate aminotransferases (AST) were determined according to the method of [31].

2.6 Tissue Cadmium Determination

The cadmium contents of the liver and kidney tissue were estimated with inductively couple plasma optical emission spectrophotometer (ICP OES) optima 2000DV after wet digestion. 1g portion of the tissue was digested with 20 ml HNO₃-HClO₄ mixture (1:4 v/v) at 100 °C and the resultant digest diluted to 100 ml with deionized water [32].

2.7 Statistical Analysis

Statistical analysis of numerical data (expressed as mean $\pm$ SD) was done using the statistical package for the social sciences software (SPSS) programme. One way analysis of variance (ANOVA) with post hoc analysis was used to assess the differences between the experimental groups and statistical significance was considered at $p < 0.05$.

3. RESULTS

3.1 Tissue Cadmium Content

The results of tissue cadmium determination are summarized in Table1. It can be observed from the table that cadmium was detected in both the liver and kidney of both the control and treated rats, but the mean cadmium contents of both the liver and kidney of rats exposed to Cd alone was significantly higher ($p < 0.05$) than those of the corresponding tissues in both the control and extract treated groups. However, the liver and kidney of rats exposed to Cd following pretreatment with aqueous extract of *Vitex doniana* (group C) had a significantly lower ($p < 0.05$) Cd content than the corresponding tissues of rats exposed to Cd alone (group B). This suggests that pretreatment with aqueous leaf extract of *Vitex doniana* impaired Cd bioaccumulation in the liver and kidney. Also, it can be observed from the table that the mean kidney tissue Cd content of rats is significantly higher ($p < 0.05$) than the liver, suggesting that the kidney is more active in bioaccumulation of Cd than the liver.

Table 1 Effect of pretreatment with aqueous leaf extract of *Vitex domain* on cadmium content of rat liver and kidneys

Group	Treatment	Cadmium concentration ($\mu\text{g/g}$ tissue)	
		Liver	Kidney
A	Control	0.14 $\pm$ 0.30	0.19 $\pm$ 0.10
B	Cd	1.60 $\pm$ 0.29 ^a	2.60 $\pm$ 0.25 ^a
C	VDE + Cd	0.42 $\pm$ 0.10 ^{ab}	0.65 $\pm$ 0.24 ^{ab}
D	VDE	0.12 $\pm$ 0.10 ^a	0.16 $\pm$ 0.10 ^{ab}

155 Data are expressed as mean $\pm$ SD, n= 4 for each group. VDE = *Vitex doniana* extract
 156 values are significantly different from control (p<0.05)
 157 ^b values are significantly different from the group treated with Cd alone (p<0.05)
 158

159 3.2 Membrane Lipid Peroxidation

160 The results of the effect of aqueous leaf extract of *Vitex doniana* on lipid peroxidation are summarized
 161 in Table 2. It can be observed from the table that there was a significant increase (P < 0.05) in the
 162 mean concentration of malondialdehyde (MDA) in rats given Cd alone (group C) when compared with
 163 the control (group A), suggesting that ingestion of Cd-induced lipid peroxidation as determined by
 164 MDA. However, pretreatment with aqueous leaf extract of *Vitex doniana* leads to a significant
 165 reduction (P < 0.05) in MDA level in the tissues examined. The mean MDA level is generally higher in
 166 the kidney than in the liver except in group A, where the reverse is the case.

167 Table 2 Effect of pretreatment with aqueous leaf extract of *Vitex domain* on cadmium-induced lipid
 168 peroxidation in the liver and kidneys of rat

Group	Treatment	Malondialdehyde concentration (nmol/g tissue)	
		Liver	Kidney
A	Control	28.53 $\pm$ 2.82	24.12 $\pm$ 1.29

B	Cd	63.62±1.66 ^a	73.81±1.89 ^a
C	VDE + Cd	38.04±1.90 ^{ab}	57.40±1.65 ^{ab}
D	VDE	19.35±1.91 ^a	15.75±1.260 ^{ab}

Data are expressed as mean ± SD, n= 4 for each group. VDE = *Vitex doniana* extract values are significantly different from control (p<0.05)

^b values are significantly different from the group treated with Cd alone (p<0.05)

3.3 Non-enzymic Tissue Antioxidants

The result of tissue glutathione determination is summarized in Tables 3. The level of glutathione was significantly much lower in the liver and kidney of rats exposed to Cd alone (group B) than in the corresponding tissue of the control (group A), suggesting that exposure to Cd markedly depleted glutathione stores of the liver and kidney. However, pretreatment with aqueous leaf extract of *Vitex doniana* had a sparing effect on tissue glutathione and inhibited its depletion by Cd. The mean glutathione content of the liver is generally higher than that of the kidney, suggesting that the liver has higher glutathione reserves than the kidney.

Table 3. Effect of pretreatment with aqueous leaf extract of *Vitex domain* on tissue glutathione content of rats exposed to Cd.

Group	Treatment	Glutathione concentration (µmol/g tissue)	
		Liver	Kidney
A	Control	2.88±0.82	1.24±0.29
B	Cd	1.09±0.63 ^a	0.70±0.10 ^a
C	VDE + Cd	1.92±0.19 ^{ab}	0.90±0.15 ^{ab}
D	VDE	2.90±0.70 ^b	1.50±0.12 ^{ab}

Data are expressed as mean $\pm$ SD, n= 4 for each group. VDE = *Vitex doniana* extract
 values are significantly different from control (p<0.05)
^b values are significantly different from the group treated with Cd alone (p<0.05)

3.4 Antioxidant Enzymes

The results of the antioxidant enzymes enzyme catalase and superoxide dismutase in the liver and kidney are summarized in Table 4. In both, the liver and kidneys, the mean activity of each of the antioxidant enzymes were significantly higher (p<0.05) in rats exposed to cadmium alone (group B) than in the control rats (group A). This would imply that exposure to cadmium-induced the activity of each of the two antioxidant enzymes, catalase and superoxide dismutase in the liver and kidney. However, pretreatment with aqueous leaf extract of *Vitex doniana* moderated the antioxidant enzyme-inducing the effect of cadmium on catalase and superoxide dismutase in both tissues examined. Catalase and superoxide dismutase activities in the liver and kidney of rats pretreated with aqueous leaf extract of *Vitex doniana* were significantly enhanced.

Table 4. Effect of pretreatment with aqueous leaf extract of *Vitex domain* on the activities of antioxidant enzymes catalase and superoxide dismutase in the liver and kidney of rats exposed to cadmium

Group	Treatment	Catalase activity ($\mu\text{mol/g}$ tissue)		SOD activity ($\mu\text{mol/g}$ tissue)	
		Liver	Kidney	Liver	Kidney
A	Control	6.10 $\pm$ 0.71	7.64 $\pm$ 0.66	9.23 $\pm$ 1.23	44.58 $\pm$ 3.1.81
B	Cd	12.45 $\pm$ 0.73 ^a	17.21 $\pm$ 0.75 ^a	55.05 $\pm$ 0.38 ^a	75.56 $\pm$ 1.81 ^a
C	VDE + Cd	8.20 $\pm$ 0.61 ^{ab}	14.54 $\pm$ 0.26 ^{ab}	36.76 $\pm$ 0.97 ^{ab}	62.55 $\pm$ 1.20 ^{ab}
D	VDE	5.10 $\pm$ 0.15 ^{ab}	5.23 $\pm$ 0.55 ^{ab}	8.02 $\pm$ 1.21 ^{ab}	33.94 $\pm$ 2.40 ^{ab}

Data are expressed as mean $\pm$ SD, n= 4 for each group. VDE = *Vitex doniana* extract
 values are significantly different from control (p<0.05)
^b values are significantly different from the group treated with Cd alone (p<0.05)

3.5 Tissue Cell Injury Enzyme Biomarkers

The results are summarized in Table 5. The mean ALT, AST and ALP activities in the liver of rats exposed to cadmium alone were significantly higher ($p<0.05$) when compared to the control rats. This would imply that exposure to Cd caused serious tissue cell damage; hence, more of these cellular enzymes are released. However, pretreatment with aqueous leaf extract of *Vitex doniana* significantly lower ($p<0.05$) the activities of cell injury biomarker enzymes in the tissues examined when compared to rats exposed to cadmium alone. The results suggest that the use of *Vitex doniana* extract caused a reduction in the level of normal metabolic oxidative damage to liver cells.

Table 5: Activities of tissue marker enzymes aspartate and alanine aminotransferases, and alkaline phosphatase in the liver of rats exposed to cadmium following pretreatment with aqueous leaf extract of *Vitex doniana*

Group	Treatment	Liver enzyme activities ($\mu\text{g tissue}$) (umol/g tissue)		
		AST	ALT	ALP
A	Control	133.50 $\pm$ 1.19	123.00 $\pm$ 0.83	81.20 $\pm$ 2.56
B	Cd	176.50 $\pm$ 6.64 ^{ab}	128.50 $\pm$ 3.70 ^b	160.10 $\pm$ 7.97 ^{ab}
C	VDE + Cd	167.50 $\pm$ 4.30 ^a	121.50 $\pm$ 2.64 ^a	91.10 $\pm$ 5.94 ^a
D	VDE	131.53 $\pm$ 3.46 ^b	121.10 $\pm$ 2.89 ^{ab}	101.20 $\pm$ 4.62 ^{ab}

Data are expressed as mean $\pm$ SD, n= 4 for each group. VDE = *Vitex doniana* extract values are significantly different from control ($p<0.05$)

^b values are significantly different from the group treated with Cd alone ($p<0.05$)

3.6 Phytochemical Constituents of *Vitex doniana*

The results of phytochemical screening of *Vitex doniana* are summarized in Table 6. The signs +, ++ and +++ represents the presence of phytochemicals in the plant extract in the trace, moderate and abundance respectively. It can be observed from the table that phytochemicals were detected at varying quantity in the aqueous leaf extract of *Vitex doniana*. The screening revealed that alkaloids, saponin, tannins, and cardiac glycosides were moderately present while flavonoids and phenols are abundantly present in the plant extract

Table 6 Results of phytochemical screening of *Vitex doniana*

Phytochemical	Alkaloids	Cardiac glycosides	Flavonoids	Phenols	Saponins	Tannins
VDE	++	++	+++	+++	+	++

VDE = *Vitex doniana* extract, + = present in trace, ++ = moderately present, +++ = abundantly present

4. DISCUSSION

Cadmium is a toxic metal that promotes early oxidative stress in animals and humans and afterwards contributes to the development of serious pathological conditions because of its long retention in some tissues [33, 34]. In the current study, cadmium administration to rats leads to a significantly increased accumulation of cadmium in the liver and kidneys of rats treated with cadmium. The generally higher concentration of Cd in the kidney than in the liver is consistent with its being the major target organ for cadmium toxicity [9]. The presence of cadmium in trace quantities in the liver and kidney of rats on the control diet has also been observed by other workers [14, 32, 35]. It is a testimony of the ubiquity of cadmium in the food chain and the environment. However, pretreatment with aqueous leaf extract of *Vitex doniana* alters the pattern of Cd disposition and markedly reduced the level of bioaccumulation of Cd in the rat liver and kidney. This is in agreement with the previous studies which showed that oral intake of cadmium induces its accumulation in these tissues [17]. The decreased cadmium concentration in the liver and kidneys could be explained by its redistribution to other tissues and organs as well as by the formation of cadmium-extract complexes or by interfering with free radical chain initiation and progression of Cd-induced oxidative damage. The decreased accumulation of cadmium in liver and kidneys after pretreatment with aqueous leaf extract of *Vitex doniana* before exposure to cadmium, indicate that extract of *Vitex doniana* diminished the toxic effects of cadmium.

Cd-induced oxidative stress is characterized by increased lipid peroxidation and altered nonenzymatic and enzymatic antioxidant system. The results of this study showed that exposure to cadmium causes a significant increase in membrane lipid peroxidation in the liver and kidneys of rats but pretreatment with aqueous leaves extract of *Vitex doniana* was effective in the prevention of oxidative damage induced by cadmium, which resulted in significantly lower degree of lipid peroxidation in the liver and

255 kidneys. This is consistent with the previous studies in which natural plant products effectively
256 reduced lipid peroxidation induced in response to various toxicants [36, 37, 38]. This can be explained
257 by the important role of polyphenols in preventing lipid peroxidation and in the protection of the
258 integrity and functioning of tissues and cells. This is in accord with reported protective effects of
259 antioxidant nutrients against Cd-induced oxidative stress and lipid peroxidation in the liver and kidney
260 [12, 15, 17].

261 Reduced glutathione (GSH) is an essential constituent of the endogenous antioxidant defence
262 system, and it functions as a direct free-radical scavenger as well as reduces intracellular reactive
263 oxygen species [39]. It is usually used up in the course of destroying the oxy-radicals leading to the
264 depletion of its tissue reserves in the process [16]. In the present study, exposure to cadmium-
265 induced a significant depletion of glutathione in the liver and kidney of rat which was effectively
266 prevented by pretreatment with aqueous leaf extract of *Vitex doniana*. The observed sparing effect of
267 aqueous leaf extract of *Vitex doniana* on Cd-induced depletion of glutathione in the rat liver and
268 kidney is consistent with a protective role for aqueous leaf extract of *Vitex doniana* against Cd toxicity.
269 This would appear to suggest that sparing effect of aqueous leaf extract of *Vitex doniana* pre-
270 supplementation or tissue glutathione was most effective in the kidney. The diminished level of liver
271 GSH in Cd-treated rats in the present study might be due to its reductive defence role in maintaining
272 an oxidant/antioxidant balance during cadmium-toxicity

273 There is an increasing body of evidence suggesting that cadmium induces alterations in the activities
274 of endogenous antioxidant enzymes with catalase and superoxide dismutase (SOD) being among the
275 first enzymatic antioxidant defences for the body against oxidant-induced cytotoxic challenge [5, 40].
276 Upon ingestion, many xenobiotics, including oxidants, such as 2, 4- dinitrophenyl hydrazine [41],
277 induce the enzymes that metabolize them. Furthermore, when cells are oxidatively challenged,
278 antioxidant enzymes levels increase as a protective mechanism [18]. In the current study, the results
279 obtained regarding the activities of SOD and catalase in the liver and kidney indicate that exposure to
280 mild doses of cadmium significantly induced enzyme activity which was effectively reversed by pre-
281 treatment with aqueous leaf extract of *Vitex doniana*. This is in accord with reported protective effects
282 of antioxidant nutrients against Cd-induced oxidative stress and alteration of levels of antioxidant

enzymes in the liver and kidney [16, 17, 18, 42]. The overall result suggests that the extract acts in a similar mode to that of Livolin forte [43].

Measurement of the activities of “marker” enzymes or biomarkers in tissues and body fluids can be used in assessing the degree of assault and the toxicity of a chemical compound on organs/tissues long before it is revealed by histological techniques [44, 45]. In this study, a marked increase in the activities of aminotransferases and alkaline phosphatase in the liver tissue of Cd-exposed rats are indications that the liver has been injured. However, pretreatment with aqueous leaf extract of *Vitex doniana* led to a marked decrease in the activities of these marker enzymes suggesting a marked reduction in the degree and rate of tissue cell injury. This is in accord with reported protective effects of the protective effect of carrot juice against cadmium-induced toxicity in the liver [18].

5. CONCLUSION

From the findings of this study on the effect of pretreatment with *Vitex doniana* leaf extract on all the indicators of the cytotoxic response to cadmium exposure, it can be concluded that an organism well nourished with the natural antioxidant components in the plant *Vitex doniana* at the time of Cd exposure has a more effective defence capability against Cd-induced oxidative damage to tissue than otherwise as determined by the various parameters. Although the study did not specifically identify the active principle but based on the well reported antioxidant properties of *Vitex doniana* due to the presence of polyphenols, polyphenols are presumed as the active antioxidant principle.

Consent: NA

REFERENCES

- [1] Nath R, Prasad P, Paliwal VK, Chopra RK, Molecular basis of cadmium toxicity, Progress in Food & Nutrition Science, 1984, 8(1-2), 109-63.
- [2] Sarker S, Yadar P, Trivedi R, Bansal AK, Bhatnagar D, Cadmium-induced lipid peroxidation and the status of the antioxidant system in rat tissues, Journal of Trace Elements in Medicine and Biology, 1995, 9(3), 144-149.
- [3] Valko M, Morris H, Cronin MT. Metals, Toxicity and Oxidative Stress. Current Medicinal Chemistry, 2005, 12, 1161-1208.
- [4] Eriyamremu GE, Ojimogho SE, Asagba SO, Osagie VE. Palm oil induced changes in ocular tissue lipid peroxidation, antioxidant enzymes and ATPases of rabbits in cadmium toxicity, 2008, 3155-3158.

- 313 [5] Patra RC, Amiya KR, Swarup D. Oxidative stress in lead and cadmium toxicity and its
314 amelioration, *Veterinary Medicine International*, 2011,1-2.
- 315 [6] Zikic RV, Stajn AS, Ognjanovic BI, Saicic ZS, Kostic MM, Pavlovic SZ, Petrovic VM. The
316 effect of cadmium and selenium on the antioxidant enzyme activities in rat heart. *Journal of*
317 *Environmental Pathology, Toxicology and Oncology*, 1998,17, 259–264.
- 318 [7] Roopha PD, Padmalatha C. Effect of Herbal Preparation on Heavy Metal (Cadmium) Induced
319 Antioxidant System in Female Wistar Rats, *Journal of Medical Toxicology*, 2012, 8(2): 101–
320 107.
- 321 [8] Waisberg M, Joseph P, Hale B, Beyersmann D. Molecular and cellular mechanisms of
322 cadmium carcinogenesis: a review, *Toxicology* 2003, 192, 95-117.
- 323 [9] Asagba SO. Role of diet in absorption and toxicity of oral cadmium-A review of literature,
324 *African Journal of Biotechnology*, 2009, 8(25), 7428-436.
- 325
- 326 [10] Tomohito A, Akira K, Koji B, Shinsuke M, Shingo M. Effects of water management on Cd and
327 As content in rice grain.19th World Congress of Soil Science, Soil Solutions for a Changing
328 World. Brisbane, Australia, 2010, 103-106.
- 329 [11] Douglas RM, Chalker EB, Treacy B. Vitamin C for preventing and treating the Cochrane
330 Database of Systematic Reviews, 2000 2:10-98.
- 331 [12] Tandon SK, Singh S, Prasad S, Khandekar K, Dwivedi VK, Chatterjee M, Mathur N, Reversal
332 of cadmium induced oxidative stress by chelating agent, antioxidant or their combination in
333 rat. *Toxicology Letters*, 2003, 145, 211-217.
- 334 [13] Nahed SH, Sahar MA, Reverse effect of vitamin E on oxidative stress, derivatives and
335 conductivity changes of hemoglobin induced by exposure to cadmium, 2007,437-443.
- 336 [14] Ognjanovic BI, Markovic SD, Pavlovic SZ, Zikic RV, Stajn AS, Saicic ZS. Effect of chronic
337 cadmium exposure on antioxidant defense system in some tissues of rats: protective effect of
338 selenium, *Physiological Research*, 2008, 57, 403-411.
- 339 [15] Mehmet KL, Burhan A, Meryem A, Yeter TT, Cevat A, Hamdi U, Vitamin E protects against
340 oxidative damage caused by cadmium in the blood of rats, 2009, 154-160.
- 341 [16] Swaran JSG. Structural, chemical and biological aspects of antioxidants for strategies against
342 metal and metalloid exposure. *Oxidative Medicine and Cellular Longevity*, 2009, 2(4),191-
343 206.
- 344 [17] Mafulul SG, Okoye ZSC, Protective effect of pre-supplementation with selenium on cadmium-
345 induced oxidative damage to some rat tissues. *International Journal of Biological and*
346 *Chemical Sciences*, 2012, 6(3), 1128-1138.
- 347 [18] Embugushiki RE, Mafulul SG, Okoye ZSC. Protective effect of carrot juice pretreatment on
348 cadmiuminduced oxidative cytotoxic damage to some rat tissues. *IOSR Journal of Pharmacy*
349 *and Biological Sciences*, 2013, 7(6), 55-62.
- 350 [19] Ahmed RS, Seth V, Banerjee BD. Influence of dietary ginger (*Zingiber officinale* Rosa on
351 antioxidant defense system in rat: comparison with ascorbic acid. *Indian Journal of*
352 *Experimental Biology*, 2000; 38:604–606.
- 353 [20] Aqil F, Ahmed I, Mehmood Z. Anti-oxidants and free radical scavenging properties of twelve
354 traditionally used Indian medicinal plants. *Turkish Journal of Biology*, 2006; 30:177–183.
- 355
- 356
- 357
- 358
- 359

- 360 [21] Quideau ST, Deffieux D, Douat-Casassus C, Pouységu L. Plant polyphenols: chemical
361 properties, biological activities, and synthesis. *Angewandte Chemie International Edition*,
362 2011; 50: 586-621.
- 363 [22] Brzóska MM, Borowska S, Tomczyk M. Antioxidants as a potential preventive and therapeutic
364 strategy for cadmium. *Current Drug Targets*, 2016, 17, 1350-1384
- 365 [23] Muhammad AI, Wudil AM, Yunusa I, Mukhtar ZG, Sharif AA, Kabara HT. Oral administration
366 of aqueous bark extract of *Vitex doniana* affects serum electrolytes levels in Albino rats. *Point*
367 *Journal of Medical Research*, 2015, 1(1): 001-005.
- 368 [24] Noda Y, Metal AK. Hydroxyl and superoxide anion radical scavenging activities of natural
369 souuce anti-oxidants using the computerized JES-FR 30 ESR spectrophotometer
370 system. *Biochemistry and Molecular Biology International*, 1997;42:35–44.
- 371 [25] Harborne JB. *Phytochemical methods: a guide to modern techniques of plant analysis*.
372 Chapman and Hall Ltd., London, 1973, 49–188
- 373 [26] Ohkawa H, Ohishi N, yagi K. Assay for lipid peroxides in animal tissues by thiobarbituric acid
374 reaction. *Analytical Biochemistry*, 1979, 95:351-358.
- 375 [27] Ellman GL. Tissue sulfhydryl groups. *Archives of Biochemistry and Biophysics* 1959, 82 (1):
376 70–7
- 377 [28] Beutler E, Duron O, Kelly BM. Improved method for the determination of blood glutathione,
378 *Journal of Laboratory and Clinical Medicine*, 1963,61, 882–888.
- 379 [29] Sinha KA. Colorimetric assay of catalase. *Analytical Biochemistry*, 1971, 47:389–394
- 380 [30] Heikkila RE, Cabbat F. 1976. A sensitive assay for superoxide dismutase based on the
381 autoxidation of 6-hydroxydopamine. *Analytical Biochemistry* 75: 356-362.
- 382 [31] Reitman S, Frankel S (1957) A colorimetric method for the determination of serum glutamic
383 oxalacetic and glutamic pyruvic transaminases. *American Journal of Clinical Pathology* 1957,
384 28:56–63.
- 385 [32] Asagba SO, Eriyamremu GE, Adaikpoh MA, Ezeoma A.2004. Levels of lipid peroxidation,
386 superoxide dismutase and Na⁺/K⁺-ATPase in some tissues of rats exposed to a Nigerian diet
387 and cadmium. *Biological Trace Element Research*, 2004, 100 (1): 075-086.
- 388 [33] Bagchi D, Bagchi M, Stohs SJ, Ray SD, Kuszynski CA, Pruess HG. Free radicals and grape
389 seed proanthocyanidin extract: importance in human health and disease prevention.
390 *Toxicology* 2000; 148: 187–97.
- 391 [34] Kanter M, Aksu B, Akpolat M, Tarladacalisir YT, Aktas C, Uysal H. Vitamin E protects against
392 oxidative damage caused by cadmium in the blood of rats. *European Journal of General*
393 *Medicine*, 2009, 6(3), 154-160.
- 394 [35] Asagba SO, Eriyamremu GE, Onyeneke EC, Suru M. 2006. Influence of a Nigerian-like diet on
395 calcium, phosphate and alkaline phosphatase levels in the plasma and bone of cadmium
396 exposed rats. *Journal of Medical Science* 6(5): 758-764.
- 397 [36] Hasan SK, Sultana S. Geraniol attenuates 2- acetylaminofluorene induced oxidative stress,
398 inflammation and apoptosis in the liver of Wistar rats. *Toxicology Mechanisms and Methods*,
399 2015, 25:559–73.
- 400
401
402

- 403 [37] Mohamed MA, Eldin IM, Mohammed AE, Hassan HM. Effects of Lawsonia inermis L.
404 (Henna) leaves methanolic extract on carbon tetrachloride induced hepatotoxicity in rats.
405 Journal of Intercultural Ethnopharmacology, 2015; 5:22–6.
- 406 [38] Kumar M, Kaur P, Chandel M, Singh AP, Jain A, Kaur S. Antioxidant and hepatoprotective
407 potential of Lawsonia inermis L. leaves against 2-acetylaminofluorene induced hepatic
408 damage in male Wistar rats , BMC Complementary and Alternative Medicine, 2017, 17(56), 2-
409 11.
- 410 [39] Oyinloye BE, Adenowo AF, Osunsanmi FO, Ogunyinka BI, Nwozo SO, Kappo AP. Aqueous
411 extract of *Monodora myristica* ameliorates cadmium-induced hepatotoxicity in male rats.
412 SpringerPlus, 2016, 5:641, 1-7.
- 413 [40] El-Sokkary GH, Nafady AA, Shabash EH. Melatonin ameliorates cadmium-induced oxidative
414 damage and morphological changes in the kidney of rat, The Open Neuroendocrinology
415 Journal, 2009, 2, 1-9.
- 417 [41] Maduka HCC, Okoye ZSC. The effect of *sacoglottis gabonensis* stem bark extract, a Nigeria
418 alcoholic beverage additive, on the natural antioxidant defences during 2,4-
419 dinitrophenylhydrazine-induced membrane peroxidation *in vivo*. Vascular pharmacology 202,
420 39:21-31.
- 422 [42] Deepti G, Shabad P, Dua KK. Chronic cadmium toxicity in rats: treatment with combined
423 administration of vitamins, amino acids, antioxidants and essential metals. Journal of Food
424 and Drug Analysis 2009,18 (6), 464-470.
- 426 [43] Olukiran OS, Akomolafe RO, Bamitale KD, Ajayi AO, Okonji RE, Bejide RA. Protective and
427 curative effects of Livolin forte® on carbon tetrachlorideinduced liver damage in Wistar rats.
428 Journal of experimental and integrative medicine, 2014, 4(1):57–65
- 429 [44] Yakubu MT, Bilbis LS, Lawal M, Akanji MA. Effect of Repeated Administration of sildenafil
430 citrate on selected enzyme activities of liver and kidney of male albino rats. Nigerian Journal
431 Pure & Applied Science, 2003, 18, 1395–1400.
- 432 [45] Nafiu MO, Akanji MA, Yakubu MT. Effect of Aqueous Extract of *Cochlospermum Planchonii*
433 Rhizome on Some Kidney and Liver Functional Indices of Albino Rats. African Journal
434 of Traditional, Complementary and Alternative medicine ,2011, 8(1), 22–26.