

Original Research Article

The relation of Vitamin D Status and Presence of Acute Coronary Syndrome in patients.

Abstract

Background: Cardiovascular disease (CVD) consists of a variety of heart diseases, illnesses and events that impact the heart & circulatory system, including high blood pressure & coronary artery disease (CAD). Vitamin D is one of the fat-soluble vitamins also known as sunshine Vitamin due to its synthesis in the body following exposure to ultraviolet (UV) B rays. The aim of this study was estimation of serum vitamin D in patients with acute coronary syndrome.

Materials and Method: The present study included 50 cases of diagnosed acute coronary artery disease patients aged between 20 to 60 years and 50 apparently healthy controls and TMT negative matched for age and sex. Serum vitamin D was measured by ELISA method.

Results: The present study shown that significantly decreased levels of serum vitamin D ($p < 0.001$) in acute coronary syndrome patients as compared with healthy controls.

Conclusion: Decreased levels of vitamin D are risk factor for acute coronary syndrome, and also novel marker of CHD.

Study Design: Prospective Observational Study.

Keywords: Vitamin D, Coronary Artery Disease, Deficiency & Serum.

INTRODUCTION

Vitamin D is one of the fat-soluble vitamins also known as sunshine Vitamin due to its synthesis in the body following exposure to ultraviolet (UV) B rays, however it is unique in a way that it acts as a prohormone and mediates its functions by binding to a member of nuclear receptor super family, the Vitamin D receptor (Aggarwal et al., 2016)^[1]. The active form of vitamin D 1,25-dihydroxyvitamin D (1,25(OH)₂D), acts as a steroid hormone by binding to the vitamin D receptor (VDR), which is present in many cells throughout the body, including cardiomyocytes, vascular smooth muscle, and endothelium (Nibbelink et al., Wu-Wong et al., Merke et al.)^[2,3,4].

Vitamin D deficiency is prevalent in most parts of the world. Vitamin D has long been known to be an essential part of bone metabolism, although recent evidence suggests that vitamin D plays a key role in the pathophysiology of other diseases, including cardiovascular disease. Emerging evidence indicates that vitamin D deficiency, cardiovascular disease and endothelial dysfunction are linked together (Zahra Dana Siadat et al. 2012)^[5]. Endothelial dysfunction is an important antecedent event in the development of CHD and atherosclerosis (McGreevy et al. 2011, Syal et al., 2012)^[6&7]. Vitamin D is known to affect vascular endothelium directly or indirectly through up-regulation of the renin-angiotensin system or

via induction of smooth muscle proliferation and a proinflammatory state. A variety of possible biological mechanism (Blood pressure elevation, Insulin resistance, Inflammation, Obesity, Endothelial dysfunction, Vascular remodeling due to hyper-parathyroidism) have been proposed by which, vitamin D deficiency may cause cardiovascular events (Pittas et al., 2007, Saneei et al., 2013, Chitalia et al., 2012)^[8,9,10]. In addition, vitamin D has anti-inflammatory effects and prevents cholesterol removal by macrophage and foam cell formation on vessels walls. Also found relation has been seen between vitamin D serum level and coronary artery calcification (Judd et al., 2008)^[11].

According the Endocrine Society Practice Guideline (2011)

Reference value of vitamin D

Vitamin D level	ng/mL
Deficiency	<20
Insufficiency	20-29
Sufficiency	30-100

MATERIAL & METHOD

The present study was conducted in the Department of Biochemistry in association with Department of Medicine of Major S.D. Singh Medical College Fatehgarh Farrukhabad. The period of study was from May 2016 to April 2017.

The present study included 50 cases of diagnosed acute coronary artery disease patients aged between 20 to 60 years and 50 apparently healthy controls and TMT negative matched for age and sex. Informed written consent was taken from all the subjects.

Inclusion criteria:

Acute coronary syndrome patients,

Age 20-60 years.

Exclusion criteria:-

1. Vitamin D supplementation.
2. Congestive heart failure
3. Significant chronic liver disease
4. Chronic kidney disease
5. Diabetes mellitus
6. Endocrine dysfunction
7. Malabsorption syndrome
8. Renal Rickets
9. Pregnant Female

Anthropometric measurements:

Anthropometric indices including height and weight were taken while subjects were in the standing position and wearing light clothing without shoes. Body weight and height were measured in kilograms and in centimeters, respectively. Body mass index (BMI) was calculated as weight (kilo- grams) divided by height squared (in square meters).

83 Collection of Samples:-

84 After overnight fasting 5 ml of venous blood sample was drawn by venepuncture
85 from a peripheral vein under all aseptic precautions in a disposable syringe. The blood thus
86 collected in clean dry glass tubes was allowed to stand for 30 minutes at room temperature
87 for the retraction of clot. Then it was centrifuged at 3000 r.p.m. for 10 minutes to separate the
88 serum. The serum sample was stored at -20°C in the refrigerator for analysis. Care was taken
89 to avoid haemolysis of the sample. Then serum was analyzed for lipid profile and vitamin D.

91 OBSERVATION & RESULTS

92 **Table 1: Baseline characteristics of study population**

Baseline characteristics	Case (Mean±SD)	Control (Mean±SD)	P value
Age	47.8±10.3	45.4±11.1	0.266
BMI	26.89±4.75	23.93±3.42	0.001
Pulse	82.7±17.4	76.98±4.04	0.025
SBP	127.8±24.6	117.60±8.47	0.007
DBP	86.80±13.0	75.80±4.76	0.000

93 Table 1 show basic characteristics of study subjects, mean age of the cases were 47.8±10.3
94 and that of controls were 45.4±11.1. Mean BMI of cases were 26.89±4.75 and of control
95 were 23.93±3.42. SBP and DBP were show significant correlation.

96 **Table 2: Comparison of serum vitamin D level in study population**

Vitamin D Level	Case	
	No.	%
Deficiency	29	58
Insufficiency	11	22
Sufficiency	10	20
Total	50	100

97 Table 2: show that 58% of cases were vitamin D deficient while 22% were insufficient.

99 Graph 1: Comparison of serum vitamin D level in study population





Table 3: The difference of mean of 25(OH) D serum levels between the two study groups

Parameter	Case (Mean±SD)	Control (Mean±SD)	p value
Vitamin D (25-OHD)	20.9±11.8	37.28±8.80	0.000

Table 3 shows that mean value of vitamin D in cases was 20.9±11.8 while mean value of vitamin D in controls was 37.28±8.80, the difference was statistically significant.

Graph 2: The difference of mean of vitamin D serum levels between the two study groups



DISCUSSION

Vitamin D has increasingly been recognized to play a role in a broad range of bodily functions beyond bone health, including CV health. In parallel, the recognition of the

presence of widespread vitamin D deficiency is increasing. The consequences of this growing epidemic of vitamin D deficiency are still not well understood. However, recent reports of the associations of vitamin D deficiency with multiple CV conditions have been of great interest, suggesting the need for prospective validation and extended observations. In present study we evaluated vitamin D in patients of ST segment elevated MI.

In a research conducted by Thomas on the survivors of the Framingham study, the risk of cardiovascular disease was estimated 1.62 (Kim et al., 2004)^[12]. The findings of both of these studies suggested that vitamin D deficiency can be associated with a high risk of cardiovascular disease. Several mechanisms may explain the association between vitamin D and cardiovascular disease. Decholecalciferol regulates renin-angiotensin axis through the suppression of the renin gene. A change of 25- hydroxycholecalciferol causes changes in the smooth muscle of the vascular wall and also inflammation and thrombosis and that could explain cardiovascular complications (Zehnder et al., 2004 and Daynes et al., 1990)^[13,14].

VITAMIN D

In the current study vitamin D deficiency was observed in 58% and insufficiency in 22% of total patients of ACS on taking total count to 80%. Whereas, only 20 % of the population had adequate Vitamin D levels in our analysis. The results of the current study are in accordance with the study of lee JH et al.^[15] wherein they reported very high prevalence up to 75% as vitamin D deficient and 21% as insufficient, making a total of 96% of patients with abnormally low vitamin D levels who presented with coronary artery disease.

CONCLUSION

The conclusion of the present study moderate to severe vitamin D deficiency is a new risk factor for developing cardiovascular disease. Vitamin D deficiency might be considered as a surrogate marker of poor health status. Because measurement of vitamin D levels is easily done and vitamin D supplements are readily available and inexpensive; so, further large scale prospective studies and clinical trials are needed to establish the cause effect relationship and to evaluate the efficacy of vitamin D supplementation in prevention of CAD.

REFERENCES

1. Aggarwal R, Akhthar T, and Jain SK. Coronary artery disease and its association with Vitamin D deficiency J Midlife Health. 2016; Apr-Jun; 7(2): 56–60.
2. Nibbelink KA, Tishkoff DX, Hershey SD, Rahman A, Simpson RU. 1,25(OH)₂-vitamin D₃ actions on cell proliferation, size, gene expression, and receptor localization, in the HL-1 cardiac myocyte. J Steroid Biochem Mol Biol Mar. 2007;103:533–7.
3. Wu-Wong JR, Nakane M, Ma J, Ruan X, Kroeger PE. Effects of Vitamin D analogs on gene expression profiling in human coronary artery smooth muscle cells. Atherosclerosis. 2006;186:20–28.
4. Merke J, Milde P, Lewicka S, Hugel U, Klaus G, Mangelsdorf DJ, et al. Identification and regulation of 1,25-dihydroxyvitamin D₃ receptor activity and biosynthesis of 1,25-dihydroxyvitamin D₃. Studies in cultured bovine aortic endothelial cells and human dermalcapillaries. J Clin Invest 1989;83(6):1903-1915.
5. Zahra Dana Siadat, Keyvan Kiani, Masoumeh Sadeghi, Amir Sina Shariat, Ziba Farajzadegan, and Maryam Kheirmand Association of vitamin D deficiency and coronary

160 artery disease with cardiovascular risk factors J Res Med Sci. 2012 Nov; 17(11): 1052–
161 1055.

162 6. Cora McGreevy and David Williams New Insights About Vitamin D and Cardiovascular
163 Disease Ann Intern Med. 2011;155:820-826.

164 7. Syal SK et al Vitamin D Deficiency, Coronary Artery Disease, and Endothelial
165 Dysfunction: Observations From a Coronary Angiographic Study in Indian Patients
166 Volume 24 - Issue 8 - August 2012 (/issue/3452)

167 8. Pittas AG, Harris SS, Stark PC, et al. The effects of calcium and vitamin D
168 supplementation on blood glucose and markers of inflammation in nondiabetic adults.
169 Diabetes care 2007;30(4):980-986.

170 9. Saneei P, Salehi-Abargouei A, Esmailzadeh A. Serum 25-hydroxy vitamin D levels in
171 relation to body mass index: a systematic review and meta-analysis. Obes Rev
172 2013;14(5):393–404.

173 10. Chitalia N, Recio-Mayoral A, Kaski JC, et al. Vitamin D deficiency and endothelial
174 dysfunction in non-dialysis chronic kidney disease patients. Atherosclerosis 2012;
175 220(1):265-268.

176 11. Judd S, Tangpricha V. Vitamin D deficiency and risk for cardiovascular disease.
177 Circulation. 2008;117:503–11.

178 12. Kim DH, Sabour S, Sagar UN, Adams S, Whellan DJ. Prevalence of hypovitaminosis D
179 in cardiovascular diseases (from the National Health and Nutrition Examination Survey
180 2001 to 2004). Am J Cardiol 2008; 102(11): 1540-4.

181 13. Zehnder D, Bland R, Chana RS, Wheeler DC, Howie AJ, Williams MC, et al. Synthesis
182 of 1,25-dihydroxyvitamin D(3) by human endothelial cells is regulated by inflammatory
183 cytokines: a novel autocrine determinant of vascular cell adhesion. J Am Soc Nephrol
184 2002; 13(3): 621-9.

185 14. Daynes RA, Araneo BA, Dowell TA, Huang K, Dudley D. Regulation of murine
186 lymphokine production in vivo. III. The lymphoid tissue microenvironment exerts
187 regulatory influences over T helper cell function. J Exp Med 1990; 171(4): 979-96.

188 15. Lee JH, Gadi R, Spertus JA, Tang F, O'Keefe JH. Prevalence of vitamin D deficiency in
189 patients with acute myocardial infarction. Am J Cardiol 2011; 107(11):1636-8.