

Original Research Article

The relation of Vitamin D Status and Presence of Acute Coronary Syndrome in patients.

Abstract

Background: Cardiovascular disease (CVD) is consists of a variety of heart disease, illnesses and events that impact the heart & circulatory system, including high blood pressure & coronary artery disease (CAD). Vitamin D is one of the fat soluble vitamins also known as sunshine Vitamin due to its synthesis in the body following exposure to ultraviolet (UV) B rays. The aim of the study is to find the status of Vitamin D & there relation to Acute Coronary Syndrome

Materials and Method: The present study included 50 cases of diagnosed acute coronary artery disease patients aged between 20 to 60 years and 50 apparently healthy controls and TMT negative matched for age and sex. Serum vitamin D was the measurement by ELISA method.

Results: The present study showed that significantly decreased levels of serum vitamin D ($p < 0.001$) in acute coronary syndrome patients as compared with healthy controls.

Conclusion: Decreased levels of vitamin D are a risk factor for the acute coronary syndrome, and also novel marker of CHD.

Study Design: Prospective Observational Study.

Keywords: Vitamin D, Coronary Artery Disease, Deficiency & Serum.

INTRODUCTION

Vitamin D is one of the fat soluble vitamins also known as sunshine Vitamin due to its synthesis in the body following exposure to ultraviolet (UV) B rays, however it is unique in a way that it acts as a prohormone and mediates its functions by binding to a member of nuclear receptor super family, the Vitamin D receptor (Aggarwal et al., 2016)^[1]. The active form of vitamin D 1,25-dihydroxyvitamin D (1,25(OH)2D), acts as a steroid hormone by binding to the vitamin D receptor (VDR), which is present in many cells throughout the body, including cardiomyocytes, vascular smooth muscle, and endothelium (Nibbelink et al., Wu-Wong et al., Merke et al.)^[2,3, 4].

Vitamin D deficiency is prevalent in most parts of the world. Vitamin D has long been known to be an essential part of bone metabolism, although recent evidence suggests that vitamin D plays a key role in the pathophysiology of other diseases, including cardiovascular disease. Emerging evidence indicates that vitamin D deficiency, cardiovascular disease and endothelial dysfunction are linked together (Zahra Dana Siadat et al. 2012)^[5]. Endothelial dysfunction is an important antecedent event in the development of CHD and atherosclerosis (McGreevy et al. 2011, Syal et al., 2012)^[6&7]. Vitamin D is known to affect vascular

endothelium directly or indirectly through up-regulation of the renin-angiotensin system or via induction of smooth muscle proliferation and a proinflammatory state. A variety of possible biological mechanism (Blood pressure elevation, Insulin resistance, Inflammation, Obesity, Endothelial dysfunction, Vascular remodeling due to hyper-parathyroidism) have been proposed by which, vitamin D deficiency may cause cardiovascular events (**Pittas et al., 2007, Saneei et al., 2013, Chitalia et al., 2012**)^[8,9,10]. In addition, vitamin D has anti-inflammatory effects and prevents cholesterol removal by macrophage and foam cell formation on vessels walls. Also found relation has been seen between vitamin D serum level and coronary artery calcification (**Judd et al., 2008**)^[11].

According the Endocrine Society Practice Guideline (2011)

Reference value of vitamin D

Vitamin D level	ng/mL
Deficiency	<20
Insufficiency	20-29
Sufficiency	30-100

MATERIAL & METHOD

The present study was conducted in the Department of Biochemistry in association with Department of Medicine of Major S.D. Singh Medical College Fatehgarh Farrukhabad. The period of study was from May 2016 to April 2017.

The present study included 50 cases of diagnosed acute coronary artery disease patients aged between 20 to 60 years and 50 apparently healthy controls and TMT negative matched for age and sex. **Informed written consent was taken from all the subjects.**

Inclusion criteria:

Patient having Acute coronary syndrome with Chest Pressure

Acute Ischaemia

Sweating

Jaw Pain

Chest Burning

Exclusion criteria:-

1. Vitamin D supplementation.
2. Congestive heart failure
3. Significant chronic liver disease
4. Chronic kidney disease
5. Diabetes mellitus
6. Endocrine dysfunction
7. Malabsorption syndrome
8. Renal Rickets
9. Pregnant Female

Anthropometric measurements:

Anthropometric indices including height and weight were taken while subjects were in the standing position and wearing light clothing without shoes. Body weight and height

were measured in kilograms and in centimeters, respectively. Body mass index (BMI) was calculated as weight (kilo- grams) divided by height squared (in square meters).

Collection of Samples:-

After overnight fasting, 5 ml of venous blood sample was drawn by venepuncture from a peripheral vein under all aseptic precautions in a disposable syringe. The blood thus collected in clean dry glass tubes was allowed to stand for 30 minutes at room temperature for the retraction of the clot. Then it was centrifuged at 3000 r.p.m. for 10 minutes to separate the serum. The serum sample was stored at -20°C in the refrigerator for analysis. Care was taken to avoid haemolysis of the sample. Then serum was analyzed for lipid profile and vitamin D.

Data analysis:-

The Statistical software namely SPSS 20.0 is used for the analysis of the data. Microsoft Word and Excel software have been used to generate graphs, tables etc.

OBSERVATION & RESULTS

Table 1: Baseline characteristics of study population

Baseline characteristics	Case (Mean±SD)	Control (Mean±SD)	P value
Age	47.8±10.3	45.4±11.1	0.266
BMI	26.89±4.75	23.93±3.42	0.001
Pulse	82.7±17.4	76.98±4.04	0.025
SBP	127.8±24.6	117.60±8.47	0.007
DBP	86.80±13.0	75.80±4.76	0.000

Table 1 shows basic characteristics of study subjects, mean age of the cases were 47.8±10.3 and that of controls were 45.4±11.1. Mean BMI of cases were 26.89±4.75 and of control were 23.93±3.42. SBP and DBP were shown significant correlation.

Table 2: Comparison of serum vitamin D level in study population

Vitamin D Level	Case	
	No.	%
Deficiency	29	58
Insufficiency	11	22
Sufficiency	10	20

Total	50	100
-------	----	-----

Table 2: show that 58% of cases were vitamin D deficient while 22% were insufficient.

Graph 1: Comparison of serum vitamin D level in study population





Table 3: The difference of mean of 25(OH) D serum levels between the two study groups

Parameter	Case (Mean±SD)	Control (Mean±SD)	p value
Vitamin D (25-OHD)	20.9±11.8	37.28±8.80	0.000

Table 3 shows that mean value of vitamin D in cases was 20.9±11.8 while mean value of vitamin D in controls was 37.28±8.80, the difference was statistically significant.

Graph 2: The difference of mean of vitamin D serum levels between the two study groups



DISCUSSION

Vitamin D has increasingly been recognized to play a role in a broad range of bodily functions beyond bone health, including CV health. In parallel, the recognition of the presence of widespread vitamin D deficiency is increasing. The consequences of this growing epidemic of vitamin D deficiency are still not well understood. However, recent reports of the associations of vitamin D deficiency with multiple CV conditions have been of great interest, suggesting the need for prospective validation and extended observations. In the present study, we evaluated vitamin D in patients of ST-segment elevated MI.

In a research conducted by Thomas on the survivors of the Framingham study, the risk of cardiovascular disease was estimated 1.62 (Kim et al., 2004)^[12]. The findings of both of these studies suggested that vitamin D deficiency can be associated with a high risk of cardiovascular disease. Several mechanisms may explain the association between vitamin D and cardiovascular disease. Decholecalciferol regulates renin-angiotensin axis through the suppression of the renin gene. A change of 25- hydroxycholecalciferol causes changes in the smooth muscle of the vascular wall and also inflammation and thrombosis and that could explain cardiovascular complications (Zehnder et al., 2004 and Daynes et al., 1990)^[13,14].

VITAMIN D

In the current study, vitamin D deficiency was observed in 58% and insufficiency in 22% of total patients of ACS on taking total count to 80%. Whereas, only 20 % of the population had adequate Vitamin D levels in our analysis. The results of the current study are in accordance with the study of lee JH et al.^[15] wherein they reported very high prevalence up to 75% as vitamin D deficient and 21% as insufficient, making a total of 96% of patients with abnormally low vitamin D levels who presented with coronary artery disease.

CONCLUSION

The conclusion of the present study moderate to severe vitamin D deficiency is a new risk factor for developing cardiovascular disease. Vitamin D deficiency might be considered as a surrogate marker of poor health status. Because measurement of vitamin D levels is easily done and vitamin D supplements are readily available and inexpensive; so, further large-scale prospective studies and clinical trials are needed to establish the cause-effect relationship and to evaluate the efficacy of vitamin D supplementation in prevention of CAD.

Ethical Disclaimer:

As per international standard or university standard written ethical permission has been collected and preserved by the authors.

REFERENCES

1. Aggarwal R, Akhthar T, and Jain SK. Coronary artery disease and its association with Vitamin D deficiency J Midlife Health. 2016; Apr-Jun; 7(2): 56–60.

- 163 2. Nibelink KA, Tishkoff DX, Hershey SD, Rahman A, Simpson RU. 1,25(OH)₂-vitamin
164 D₃ actions on cell proliferation, size, gene expression, and receptor localization, in the
165 HL-1 cardiac myocyte. *J Steroid Biochem Mol Biol Mar.* 2007;103:533–7.
- 166 3. Wu-Wong JR, Nakane M, Ma J, Ruan X, Kroeger PE. Effects of Vitamin D analogs on
167 gene expression profiling in human coronary artery smooth muscle cells. *Atherosclerosis.*
168 2006;186:20–28.
- 169 4. Merke J, Milde P, Lewicka S, Hugel U, Klaus G, Mangelsdorf DJ, et al. Identification
170 and regulation of 1,25-dihydroxyvitamin D₃ receptor activity and biosynthesis of 1,25-
171 dihydroxyvitamin D₃. Studies in cultured bovine aortic endothelial cells and human
172 dermalcapillaries. *J Clin Invest* 1989;83(6):1903-1915.
- 173 5. Zahra Dana Siadat, Keyvan Kiani, Masoumeh Sadeghi, Amir Sina Shariat, Ziba
174 Farajzadegan, and Maryam Kheirmand Association of vitamin D deficiency and coronary
175 artery disease with cardiovascular risk factors *J Res Med Sci.* 2012 Nov; 17(11): 1052–
176 1055.
- 177 6. Cora McGreevy and David Williams New Insights About Vitamin D and Cardiovascular
178 Disease *Ann Intern Med.* 2011;155:820-826.
- 179 7. Syal SK et al Vitamin D Deficiency, Coronary Artery Disease, and Endothelial
180 Dysfunction: Observations From a Coronary Angiographic Study in Indian Patients
181 Volume 24 - Issue 8 - August 2012 (/issue/3452)
- 182 8. Pittas AG, Harris SS, Stark PC, et al. The effects of calcium and vitamin D
183 supplementation on blood glucose and markers of inflammation in nondiabetic adults.
184 *Diabetes care* 2007;30(4):980-986.
- 185 9. Saneei P, Salehi-Abargouei A, Esmailzadeh A. Serum 25-hydroxy vitamin D levels in
186 relation to body mass index: a systematic review and meta-analysis. *Obes Rev*
187 2013;14(5):393–404.
- 188 10. Chitalia N, Recio-Mayoral A, Kaski JC, et al. Vitamin D deficiency and endothelial
189 dysfunction in non-dialysis chronic kidney disease patients. *Atherosclerosis* 2012;
190 220(1):265-268.
- 191 11. Judd S, Tangpricha V. Vitamin D deficiency and risk for cardiovascular disease.
192 *Circulation.* 2008;117:503–11.
- 193 12. Kim DH, Sabour S, Sagar UN, Adams S, Whellan DJ. Prevalence of hypovitaminosis D
194 in cardiovascular diseases (from the National Health and Nutrition Examination Survey
195 2001 to 2004). *Am J Cardiol* 2008; 102(11): 1540-4.
- 196 13. Zehnder D, Bland R, Chana RS, Wheeler DC, Howie AJ, Williams MC, et al. Synthesis
197 of 1,25-dihydroxyvitamin D(3) by human endothelial cells is regulated by inflammatory
198 cytokines: a novel autocrine determinant of vascular cell adhesion. *J Am Soc Nephrol*
199 2002; 13(3): 621-9.
- 200 14. Daynes RA, Araneo BA, Dowell TA, Huang K, Dudley D. Regulation of murine
201 lymphokine production in vivo. III. The lymphoid tissue microenvironment exerts
202 regulatory influences over T helper cell function. *J Exp Med* 1990; 171(4): 979-96.
- 203 15. Lee JH, Gadi R, Spertus JA, Tang F, O'Keefe JH. Prevalence of vitamin D deficiency in
204 patients with acute myocardial infarction. *Am J Cardiol* 2011; 107(11):1636-8.