

A comparative study of High Sensitive C-Reactive Protein (hs-CRP) and urinary calcium/creatinine ratio in normal pregnancy and pregnancy induced hypertension in Gujarat, India

ABSTRACT

Background Pregnancy-induced hypertension (PIH) is a condition characterized by high blood pressure during pregnancy. It is the most common leading cause of maternal and perinatal morbidity and mortality in females. Serum high sensitive C-reactive protein (hs-CRP) is one of the suitable markers for low grade inflammation evaluation. Urinary calcium-creatinine ratio (UCa/Cr) is a valuable marker for prediction of PIH.

Aim To measure & correlate serum hs-CRP and urinary calcium-creatinine ratio in between pregnancy-induced hypertension and normal pregnancy.

Material & Methods In present study total 150 subjects attending Department of Obstetrics & Gynecology at Sir Takhtsinhji Hospital, Bhavnagar, Gujarat, were included which were categorized in two groups. Group A: 75 patients of pregnancy-induced hypertension; Group B: 75 normal pregnant women as a control group. They were primarily diagnosed based on measurement of blood pressure, clinical examination followed by biochemical investigations like hs-CRP, urinary calcium & creatinine, RBS, urea, creatinine, uric acid, CK-MB. Written informed consent was obtained from all participants before enrolling in the study.

Results No significant change in the levels of biological parameters like RBS, urea, creatinine and CK-MB was observed in both the study groups ($p>0.05$) except serum uric acid shows a significant difference ($p<0.001$). The levels of hs-CRP were found to be significantly higher in patients of pregnancy-induced hypertension (7.81 ± 3.681) as compared to the normal pregnant women (3.11 ± 1.972) ($p<0.0001$). UCa/Cr ratio had more sensitivity and specificity than hs-CRP for prediction of pregnancy-induced hypertension compared to normal pregnancy.

27 **Conclusion** UCa/Cr ratio can be used as a more sensitive and specific test to detect pregnancy
28 induced hypertension.

29 **Key Words** *hs-CRP, urinary calcium creatinine ratio, pregnancy induced hypertension*

30

31 INTRODUCTION

32 Pregnancy induced hypertension (PIH) is one of the most common leading cause of maternal
33 and perinatal morbidity and mortality in females & it complicates approximately one out of every ten
34 pregnancies. Early detection of PIH is more important to save mother and fetal life. PIH is associated
35 with increased risks of serious outcomes such as premature delivery, intra uterine growth retardation
36 (IUGR), perinatal mortality & morbidity, acute renal failure, acute hepatic failure, intra-partum
37 haemorrhage, post-partum haemorrhage, maternal mortality & morbidity following complication such
38 as haemolysis (H), elevated liver enzymes (EL), low platelets (LP) (HELLP syndrome), disseminated
39 intravascular coagulation (DIC) and seizures.^[1,2,3,4]

40 Pre-eclampsia (PE) is most common complication of pregnancy induced hypertension. The
41 incidence of PE in hospital practice varies widely from 5 to 15% in India. The incidence of PE in
42 primigravidae is about 10% and in multigravidae 5%. The etiology of endothelial dysfunction in PE is
43 not known, but it has been postulated to be part of an exaggerated maternal inflammatory response to
44 pregnancy.^[5,6,7,8]

45 There has been an increasing interest for the detection of pregnancy disorders before the
46 symptoms actually occur. Endothelial dysfunction is accompanied by elevated levels of inflammatory
47 markers. Indeed, such marker levels have been shown to be much higher in women with PE
48 compared to normal pregnant females.^[9]

49 Serum high sensitive C-reactive protein (hs-CRP) is one of the suitable markers for low-grade
50 inflammation evaluation. This marker rises subsequent to stress which are accompanied by
51 endothelial dysfunction and lead to peripheral vascular remodelling, decreased compliance and
52 vascular stiffness. Inflammation is considered to have a crucial role in pathophysiologic mechanism of
53 PE and endothelial dysfunction is accompanied by elevated levels of inflammatory markers.^[10]

54 Urinary calcium-creatinine ratio (UCa/Cr) is a valuable marker for prediction of PIH.
55 Physiological changes during normal pregnancy leads to increase in glomerular filtration rate those
56 results in increasing of creatinine clearance and calcium excretion in urine. But in women developing
57 PE because of vasospasm and decrease of renal flow, creatinine clearance decreases, followed by
58 increase in serum creatinine level.^[12]

Anil bargale et al. concluded that there was significantly increased level of serum hs-CRP in PE patients as compared to normal pregnant women. They also found positive significant correlation between hs-CRP levels and blood pressure.^[13]

Amandeep Kaur et al. demonstrated that urinary calcium excretion in preeclamptic females was significantly lower than that observed in case of age matched normal pregnant females and normal non pregnant females which rules out the possibility of decreased resorption of calcium at the level of renal tubules, while urinary creatinine excretion in normal pregnant females was found to be more than that observed in case of normal non-pregnant females, which were responsible for the decreased urinary calcium: urinary creatinine ratio in PE compared to normal pregnancies.^[5]

Keeping in view the different outcomes of the researchers, pregnancy induced hypertension remains most significant and intriguing unsolved problem in obstetrics. With the background of above facts the present study is designed to determine and correlate the level of serum hs-CRP and UCa/Cr ratio between normal pregnancy and PIH.

OBJECTIVES FOR THE PRESENT STUDY WERE

- To compare serum high sensitive C-reactive protein level, urinary calcium and urinary creatinine level in normal pregnancy and pregnancy induced hypertension.
- To compare random blood sugar, urea, creatinine, uric acid, creatine kinase-MB level in normal pregnancy and pregnancy induced hypertension.
- To study the correlation of high sensitive C-reactive protein, urinary calcium/creatinine ratio in between normal pregnancy and pregnancy induced hypertension.

MATERIAL & METHODS

The present study was conducted at Department of Biochemistry, Government Medical College and Sir Takhtsinhji General Hospital, Bhavnagar in which 75 cases of pregnancy induced hypertension and 75 normal pregnant women with normal blood pressure as a control group were included. **Informed consent was taken from them.** All cases were admitted to the Gopnath maternity ward (obstetric ward) of Sir T. General Hospital, Bhavnagar. They were primarily diagnosed for pregnancy induced hypertension by clinical examination followed by Biochemical investigations. The

90 subjects in control group were selected by measuring blood pressure and by following inclusion-
91 exclusion criteria

92 Case Group - Patients of pregnancy induced hypertension = 75

93 Control Group - Normal pregnant women = 75

94 **Inclusion criteria: (For cases)**

- 95 • Pregnant women with gestational age >20 week.
- 96 • Patient is a case of hypertension -
- 97 Pre-eclampsia, eclampsia and gestational hypertension (systolic blood pressure greater than
- 98 140 mm Hg or diastolic blood pressure 90 mm Hg)
- 99 • Singleton pregnancy

100 **Exclusion criteria: (For both Groups)**

- 101 • Not willing to participate in study
- 102 • Rheumatoid arthritis/Cardiovascular disease/Renal disorder/Diabetes mellitus
- 103 • Any recent or present fever or infectious disease
- 104 • Multiple pregnancy
- 105 • Smoking
- 106 • Any vaginal bleeding
- 107 • Previous history of PIH/Chronic hypertension

108 **Laboratory Investigations**

109 . Venous blood sample was collected in plain and fluoride vacutainer and urine sample was
110 collected in sterile urine container. All the samples were analyzed on fully auto analyzer ILAB-650
111 (Instrumentation Laboratory, Italy) at NABL accredited Clinical Biochemistry Section, Laboratory
112 Services Sir T. Hospital, Bhavnagar.

113 Following laboratory investigations were done in study group and control group

114 From Venous blood:

- 115 • high sensitive C-reactive protein
- 116 • Random blood sugar
- 117 • Blood urea
- 118 • Serum creatinine
- 119 • Uric acid

- Creatine kinase - MB

From Urine Sample

- Urinary calcium
- Urinary creatinine

For all the investigations, SOPs were used. Estimation of high sensitive c- reactive protein (hs-CRP) was done by turbidimetric immunoassay method ^[14] urinary calcium by modified arsenazo method ^[15] creatinine by Modified Jaffe's Kinetic method ^[16] , RBS by Glucose oxidase peroxidase, urea estimation by urease-gldh kinetic method, uric acid by uricase, CK-MB estimation by immuno-inhibition method ^[17,18]. Internal quality control and results of external quality control for the above parameters were within range. No test was done from hemolyzed sample, repeat sample was ordered.

Biological reference interval of RBS is upto 140mg/dl, urea is 15-45mg/dl, creatinine is 0.6-1.1mg/dl, uric acid is 2.6-6.0 mg/dl, CKMB is 18-51 U/L and hs CRP is 0.3 – 8.8 mg/l

Data were analyzed by using GraphPad instat version 3.0. In data analysis, comparison of these parameters were carried out by applying unpaired t-test and Fisher's exact test. Interpretation was done according to p-value. *p < 0.05 - significant and **p < 0.001 - highly significant. All the subjects were divided in following groups:

Group A: Patients of pregnancy induced hypertension (Case group)

Group B: Normal pregnant women who volunteered to participate in the present study. (Control group).

RESULTS

The present study included 75 patients of pregnancy induced hypertension and 75 normal pregnant women with normal blood pressure as a control group in which hs-CRP and urinary calcium, urinary creatinine, urinary calcium creatinine ratio were measured.

Mean $\pm$ SD of age among cases was 24.65 $\pm$ 4.180 years which include 13 patients (17%) $\leq$ 20years, 39 patients (52%) between 21-25 years, 17 patients (25%) between 26-30 years and 6 patients (8%) $\geq$ 31 years of age. The maximum number of patients belongs to age group between 21-25 years. In controls Mean $\pm$ SD of age was 24.52 $\pm$ 4.078years which include 10 normal pregnant women (13%) $\leq$

20years, 45 normal pregnant women (60%) between 21-25 years, 16 normal pregnant women (21%) between 26-30 years and 4normal pregnant women (5%) $\geq$ 31 years of age. The maximum number of normal pregnant women belongs to age group between 21-25 years. There was no statistically significant difference of age between two groups.

Table 1 shows the mean level of systolic blood pressure was 153.89 ± 12.512 in Group A and 113.81 ± 7.599 in Group B, while diastolic blood pressure level was 97.14 ± 9.799 in Group A and 73.70 ± 6.768 in Group B. Patients of PIH has significantly higher level of systolic and diastolic blood pressure than normal pregnant women ($p < 0.0001$), difference between them were statistically highly significant ($p < 0.0001$)

As per Table 2, only serum uric acid shows significant difference ($p < 0.001$) in between two groups where as other biochemical parameters like RBS, urea, creatinine and CK-MB did not show significant difference in their values($p > 0.05$) in both the groups. As evident from table 5 mean and SD of serum hs-CRP was 7.81 ± 3.681 in group A and 3.11 ± 1.972 in Group B, difference between them were statistically highly significant ($p < 0.0001$). Mean and SD of urinary calcium was 6.95 ± 1.703 in group A and 9.79 ± 3.370 in Group B. Patients of PIH have significantly lower urinary calcium level than normal pregnant women, difference between them were statistically highly significant ($p < 0.0001$). While mean and SD of urinary creatinine was 157.61 ± 53.159 in Group A and 103.88 ± 40.817 in Group B. Urinary creatinine level was significantly higher in patients of PIH as compared to normal pregnant women, difference between them were statistically highly significant ($p < 0.0001$). Moreover, mean and SD of urinary calcium creatinine ratio was 0.047 ± 0.023 in group A and 0.105 ± 0.047 in Group B. Patients of PIH have significantly lower UCa/Cr ratio than normal pregnant women, difference between them were statistically highly significant ($p < 0.0001$) where, comparison of two groups was done by unpaired t-test.

Table 3 and 4 shows the Sensitivity, Specificity, Positive predictive value & Negative predictive value of serum hs-CRP and urinary calcium creatinine ratio by using Fisher's exact test. As evident from table 4 at the cutoff level of 5 mg/L for hs-CRP the sensitivity of hs-CRP in relation to pregnancy induced hypertension is 77 % and the specificity is 82 %. Positive predictive value is 81 %, while the negative predictive value is 78 %. As evident from table 4, at the cutoff level of 0.057 the sensitivity of UCa/Cr ratio in relation to pregnancy induced hypertension is 80 % and the specificity is 86 %. Positive predictive value is 85 %, while the negative predictive value is 81 %. Hence, UCa/Cr

ratio had more sensitivity and specificity than hs-CRP for prediction of pregnancy induced hypertension.

TABLE 1: COMPARISON OF BLOOD PRESSURE LEVELS BETWEEN GROUP A AND GROUP B

STATISTICS	SYSTOLIC BP		DIASTOLIC BP	
	GROUP A	GROUP B	GROUP A	GROUP B
	PIH	NPW	PIH	NPW
Mean	153.89	113.81	97.14	73.70
Standard Deviation	12.512	7.599	9.799	6.768
Standard Error of Mean	1.445	0.8775	1.131	0.7815
Significance	p<0.0001		p<0.0001	

PIH- pregnancy induced hypertension, NPW-normal pregnant women

TABLE 2: COMPARISON OF BIOCHEMICAL PARAMETERS BETWEEN GROUP A AND GROUP B

Parameter	Biological Reference Interval	Group A PIH	Group B NPW	Statistical Significance
		MEAN $\pm$ SD	MEAN $\pm$ SD	
RBS	Up to 140 mg/dl	95.96 $\pm$ 16.665	92.65 $\pm$ 13.596	t=1.331 p=0.1851
Urea	15-45 mg/dl	20.88 $\pm$ 7.876	19.09 $\pm$ 6.927	t=1.475 p=0.1423
Creatinine	0.6-1.1 mg/dl	0.71 $\pm$ 0.218	0.66 $\pm$ 0.146	t=1.967 p=0.0511
UA	2.6-6.0 mg/dl	6.32 $\pm$ 1.655	4.66 $\pm$ 1.046	t=7.348 p<0.001
CK-MB	18-51 U/L	31.746 $\pm$ 8.320	34.49 $\pm$ 8.703	t=1.976 p=0.050
Urinary Calcium		6.92 $\pm$ 1.70	9.79 $\pm$ 3.37	t=8.548 p<0.0001
Urinary Creatinine		157.6 $\pm$ 53.1	103.9 $\pm$ 40.8	t=11.751 p<0.0001
UCa/Crratio		0.04 $\pm$ 0.02	0.10 $\pm$ 0.04	t=7.831 p<0.0001

*p < 0.05 - significant, **p< 0.001 - highly significant, #p $\geq$ 0.05 - not significant

TABLE-3: SERUM HS-CRP LEVELS AND UCA/CRRATIO IN GROUP A AND GROUP B

	Group A PIH (N=75)	Group B NPW (N=75)	Total (N=150)
hs-CRP(mg/l)			
≤ 5	58 (39%)	13 (9%)	71 (47%)
> 5	17 (11%)	62 (41%)	79 (53%)
UCa/Cr ratio			
≤0.057	60 (40%)	10 (7%)	70 (47%)
>0.057	15 (10%)	65 (43%)	80 (53%)

TABLE-4: OVERALL PREDICTIVE VALUE OF HS-CRP AND UCA/CR RATIO IN STUDY GROUPS

	HS-CRP	UCA/CR RATIO
Sensitivity	77 %	80 %
Specificity	82 %	86 %
Positive Predictive Value	81 %	85 %
Negative Predictive Value	78 %	81 %

TABLE 5: COMPARISON OF HS-CRP LEVELS BETWEEN GROUP A AND GROUP B

Statistics	hs-CRP (R.I. = 0.3 – 8.8 mg/l) (n=75)	
	Group A PIH	Group B NPW
Mean	7.81	3.11
Standard Deviation	3.681	1.972

Standard Error of Mean	0.4250	0.2277
Significance	p<0.0001	

DISCUSSION

Pregnancy induced hypertension (PIH) is associated with high maternofetal mortality and morbidity in both underdeveloped and developed countries. Approximately 70% of hypertensive disorders are due to pre-eclampsia (PE). In modern obstetric practice, hypertensive disorders of pregnancy are understood to encompass a clinical spectrum of abnormalities ranging from minimal elevation in blood pressure to severe hypertension with multiorgan dysfunction. The worldwide incidence of PE is 3-4% of all pregnancies.^[19] In the United States, PE accounts for 15% to 17.6% of maternal deaths. In India the incidence was reported as 8-10% of pregnancies being 10% in primigravida, 5% in multigravida.^[2]

While the exact causes of PE are not well understood, certain factors may increase a woman's risk of developing PIH or PE. The current hypothesis regarding the etiology of PE focuses on mal-adaptation of the immune responses and defective trophoblast invasion. Thus, an excessive maternal inflammatory response, perhaps directed against foreign fetal antigens, results in a chain of events including shallow trophoblast invasion, defective spiral artery remodeling, placental infarction and release of pro-inflammatory cytokines in the systemic circulation.^[2,9]

It has been shown that CRP is elevated in women with PE.^[20] Determination of hs-CRP has been suggested to be more sensitive than conventional measurement of CRP and provides better sensitivity in confirmation of inflammation. Recently, several studies have conducted to elucidate a relationship between PE and serum hs-CRP levels.^[21]

Renal excretion of calcium increases during pregnancy, maximum excretion levels reached during the third trimester. Urinary excretion of calcium levels in non-pregnant women is in the range about 100 - 250 mg/day, in pregnant women the range is between 350-620 mg/day. There is a decrease in urinary calcium levels in pre-eclampsia. Decreased renal filtration and increased tubular reabsorption may result in hypocalciuria.^[22]

The aetiology of hypocalciuria in preeclamptic patients is unknown. It has been speculated that hypocalciuria may result from decreased dietary intake, decreased intestinal absorption, increased calcium uptake by the fetus and placenta, or intrinsic renal tubular dysfunction.^[23]

The mean and SD of systolic blood pressure was high 153.89 ± 12.512 in patients of PIH and 113.81 ± 7.599 in normal pregnant women, while diastolic blood pressure was high 97.14 ± 9.799 in patients of PIH and 73.70 ± 6.768 in normal pregnant women with statistically significant difference ($p < 0.0001$) (Table-1).

Only serum uric acid shows significant difference ($p < 0.001$) in between group A and Group B. The mean and SD of serum hs-CRP was 7.81 ± 3.681 in patients of PIH and 3.11 ± 1.972 in normal pregnant women, difference between them were statistically highly significant ($p < 0.0001$).

According to Vijayalakshmi et al.^[10] serum hs-CRP level significantly elevated in pre-eclampsia suggesting pre-eclampsia to be an exaggerated inflammatory condition when compared to normal pregnancy.

Yasmin Ashraf et al.^[23] reported that serum hs-CRP level measured in preeclamptic group were significantly higher 14.80 ± 0.679 mg/l compared to control group 4.978 ± 0.336 mg/l with p-value 0.0001.

Zaima Ali et al.^[24] in 2013 discovered that The hs-CRP level was significantly increased in third trimester in pregnancies complicated by PE as compared to normotensive pregnant women and concluded that exaggerated systemic inflammation is characteristic of PE

In our study patients of PIH have significantly lower urinary calcium level than normal pregnant women ($p < 0.0001$). Urinary creatinine level was significantly higher in patients of PIH as compared to normal pregnant women ($p < 0.0001$). The mean and SD of urinary calcium creatinine ratio was 0.047 ± 0.023 in group A and 0.105 ± 0.047 in Group B. Patients of PIH had significantly lower UCa/Cr ratio than normal pregnant women with statistically significant difference ($p < 0.0001$) (Table-2).

N.V. Lakshmi et al.^[1] reported that urinary calcium creatinine ratio was used in prediction of pregnancy induced hypertension & there was significant lowering of urinary excretion of calcium in cases who developed PIH. PIH developed more in cases with $UCa/Cr < 0.04$.

Rashmi Sinha et al.^[25] reported that UCa/Cr ratio at 0.04 in spot urine sample being a good test for prediction of pre-eclampsia can be recommended as a screening test in all asymptomatic pregnant women.

Babli Yadav et al.^[26] observed in 2014 that significant fall in urine calcium creatinine ratio in PIH study group compared to normal pregnant women

Swapna V. S et al. [22] reported that Urinary calcium and creatinine level were significantly decreased in gestational hypertensive patients compared to normotensive pregnant women

The sensitivity and specificity of serum hs-CRP for prediction of pregnancy induced hypertension was 77% and 82% respectively with 81% positive predictive value and 78% negative predictive value (Table-3,4). The sensitivity and specificity of UCa/Cr ratio for prediction pregnancy induced hypertension was 80% and 86% respectively with 85% positive predictive value and 81% negative predictive value (Table-4).

From above findings it is indicated that serum hs-CRP level was significantly increased and UCa/Cr ratio was significantly decreased in pregnancy-induced hypertension compared to normal pregnancy.

UCa/Cr ratio had more sensitivity and specificity than hs-CRP for prediction of pregnancy induced hypertension compared to normal pregnancy (Table-4).

CONCLUSION

In the present study, it is concluded that serum hs-CRP level was significantly increased and UCa/Cr ratio was significantly decreased in pregnancy-induced hypertension compared to normal pregnancy. Serum hs-CRP at the cutoff level 5 mg/L can be useful in identifying pregnant women at risk for pregnancy induced hypertension. UCa/Cr ratio <0.057 in spot urine sample can be used as a good test for prediction of pregnancy induced hypertension. However, in patients of PIH serum hs-CRP show less predictive value as compared to UCa/Cr ratio, therefore UCa/Cr ratio can be used as a more sensitive and specific test to detect pregnancy induced hypertension which is a non-invasive and inexpensive test.

Ethical Disclaimer:

As per international standard or university standard written ethical permission has been collected and preserved by the authors.

288 **REFERENCES**

- 289 1. Dr. N.V. Lakshmi, Dr. P. Kiranmai, Dr. K. Ambika Devi and Dr. J. Rama Rao. Role of Urinary
290 Calcium Creatinine Ratio in Prediction of Pregnancy Induced Hypertension. *Int. J Pharm Bio*
291 *Sci.* 2013 July;4(3):(B)1021-1026.
- 292 2. Nicole R. Anderson, Megan Undeberg, Karen MS Bastianelli. Pregnancy-Induced
293 Hypertension and Preeclampsia: A Review of Current Antihypertensive Pharmacologic
294 Treatment Options. *Austin J Pharmacol Ther.* 2013;1(1):2373-6208.
- 295 3. Manjusha Sajith, Vandana Nimbargi, Amit Modi, Ronak Sumariya, Atmaram Pawar. Incidence
296 of pregnancy induced hypertension and prescription pattern of antihypertensive drugs in
297 pregnancy. *Apr* 2014;5(4):163-170.
- 298 4. Khosravi, S., Dabiran, S., Lotfi, M. and Asnavandy, M. Study of the Prevalence of
299 Hypertension and Complications of Hypertensive Disorders in Pregnancy. *Open Journal of*
300 *Preventive Medicine.* 2014;(4):860-867.
- 301 5. Amandeep Kaur et al., Serum and Urinary Calcium and Creatinine Levels in the 3rd Trimester
302 of Normal Pregnant and Pre-eclamptic Females. *International Journal of Recent Scientific.*
303 July 2015;6(7):4908-4913.
- 304 6. Nanda K, Sadanand G, Muralidhara Krishna C S, K L Mahadevappa. C - reactive protein as a
305 predictive factor of preeclampsia. *Int J Biol Med Res.* 2012;3(1): 1307-1310.
- 306 7. Gandhi MR, Jani PS, Patel UM, Kakani CR, Thakor NC, Gupta N. Perinatal outcome in
307 pregnancy induced hypertension cases at GMERS Medical College, Dharpur-Patan, North
308 Gujarat region, India: a prospective study. *Int J Adv Med.* 2015;2:152-5.
- 309 8. Dutta. Text book of OBSTETRICS. Hypertensive Disorders in Pregnancy. 7th edition; p. 219-
310 241.
- 311 9. Z Tavana, J Zolghadri, G Madadi. The Relationship between Maternal serum Highly Sensitive
312 C - reactive protein, Leptin and Hypertensive Disorders of Pregnancy. *The Internet Journal of*
313 *Endocrinology.* 2010;6(2);1-7.
- 314 10. Vijayalakshmi P et al. Study of serum hsCRP and lipid profile in pre-eclampsia. *International*
315 *Journal of Recent Trends in Science and Technology.* April 2015; 14(3):605-609.

11. Samira Behboudi- Gandevari, Narges Alian Moghadam, Lida Mogadam- banaem, Bitamohamadi and Mohamad Asghari. Association of high-sensitivity C-reactive protein serum levels in early pregnancy with the severity of preeclampsia and fetal birth weight. *J. Perinat. Med.* 40. (2012);601–605.
12. Azita fath Negad Kazemi, Fahime shhatie, Nilofar Satarjade, Mehrangiz Ebrahimi Mameghani. The Predictive value of Urinary calcium to creatinine Ratio, Roll-over test and BMI in early diagnosis of Pre-eclampsia. *Research journal of Biological sciences*. 2010;5(2):183-186.
13. Anil Bargale, Jayashree V. Ganu, Dhiraj J. Trivedi, Nitin Nagane, Rakesh Mudaraddi, Aparna Sagare. Serum hs-CRP and Uric Acid as Indicator of Severity in Preeclampsia. *International Journal of Pharma and Bio Sciences*. July-sept 2011;2(3);B:340-345.
14. Macy E M, *Clin Chem*, 1997;43:52-58
15. Kessler G, *Clin Chem*, 1964;10(8); 686-706
16. Kaplam A, *Clin Chem*, 1984; 1261-1266
17. DGKC, *J. Clin. Chem. Clin. Bioch.* 1977;15:255.
18. Di. witt, C. Trendelenburg, *J. clin. chemie, clin. Bioch.* 1982;20:235.
19. Alice Wang, Sarosh Rana and S. Ananth Karumanchi. Preeclampsia: The Role of Angiogenic Factors in Its Pathogenesis. *Physiology*. 2009;24:147-158.
20. Mehdi Farzadnia, Hossein Ayatollahi, Maliheh Hasan-Zade, Hamid Reza Rahimi. A comparative study of vascular cell adhesion molecule-1 and high-sensitive C-reactive protein in normal and preeclamptic pregnancies. *Interventional Medicine & Applied Science*. 2012;4(2):26-30.
21. Hossein Ayatollahi, Maliheh Hasan-zade, Mahdi Farzadnia, Mahdi Khabbaz Khoob, Atefeh Rahmanian. Serum Level of High Sensitive C - reactive protein in Normal and Preeclamptic Pregnancies. *Iranian Journal of Pathology*. 2007;2(3): 100-104.
22. Swapna V. S, Triveni Jambale, Jayaprakash Murthy D. Study of Urinary Calcium and Urinary Creatinine Levels and Urinary Calcium/Creatinine Ratio in Gestational Hypertensive Patients. *Journal of Evolution of Medical and Dental Sciences*. 2015;4(53):9145-9150.

- 344 23.Yasmin Ashraf, Nabila Roohi, Aasia Sharif, Samina Ashraf, & Sadaf Ilyia. Elevated Levels of C
345 - reactive protein in Preeclamptic Women Following 20th Week of Pregnancy. Biologia
346 (Pakistan) December. 2015;61(2):307-311.
- 347 24.Zaima Ali, Saima Zaki, Ambreen Tauseef, Ayesha Akmal. C Reactive Protein levels are
348 elevated in the Third Trimester in Preeclamptic pregnant Women. P J M H S.Jan-Mar
349 2013;7(1):188-190
- 350 25.Rashmi Sinha, Indu Bhushan. Study of Urinary Calcium/Creatinine Ratio (CCR) In a Spot
351 Sample of Urine for Early Prediction of Preeclampsia. IOSR Journal of Dental and Medical
352 Sciences. May 2016;15(5);Ver. VIII:101-104
- 353 26.Dr. Babli Yadav, Dr. Sangita Paneri, Dr. Sumitra Yadav. Evaluation of Oxidative Stress and
354 Urinary Calcium Creatinine Ratio in Pregnancy Induced Hypertension. Global Journal of
355 Medical research: E Gynecology and Obstetrics.2014;14(1);Ver.1.0:1-2.
356