

Nutrients and Constituents Relevant to Antioxidant, Antimicrobial and Anti-Breast Cancer Properties of *Salvia officinalis* L.

ABSTRACT

Objective: Comprehensive assessment of chemical composition of sage leaves (*Salvia officinalis* L.) in order to highlight constituents relevant to antioxidant, antimicrobial and anticancer potentials, in addition to determining safe dose to facilitate its application in functional food and dairy products.

Methods: High Performance Liquid Chromatography (HPLC) was employed to determine constituents such as amino acids, fatty acids and phenolic compounds content. Antioxidant activity was characterized using, α -diphenyl- β -picrylhydrazyl (DPPH) and reducing power methods. The antimicrobial potentials were examined against nine pathogenic strains. MDA-MB-231 cell line was used to assess anticancer activity.

Results: Sage was found to be a good source of calcium, iron, zinc and vitamins B6 and B12. Performed HPLC analysis indicated rich content of essential amino acids (lysine, phenylalanine and leucine), polyunsaturated fatty acids and phenolic compounds (quercetin and cinnamic), which interpreted its high antioxidant powers. Sage revealed antimicrobial and antioxidant potentials that could nominate it as applicable food bio-preservative, not to mention its anticancer effect as an extra pharmacological feature. Sage performed anti-proliferative activity with IC_{50} of 300 μ g/ mL, against MDA-MB-231 cell line.

Conclusion: Obtained results emphasis sage leaves content of variable nutrients and active compounds that reflected on its vast nutritional and pharmacological potentials such as; antioxidant, antimicrobial and cytotoxic effect against breast MDA-MB-231 cell line.

Key words: Sage (*Salvia officinalis*); HPLC; Antioxidant potentials; Antimicrobial properties; MDA-MB-231 cell line.

1. INTRODUCTION

Salvia officinalis “sage”, which called Maramia in Levant, is initially derived from the Latin *salvere* (to save), pointing out to its healing powers (Hamidpour et al., 2014). *Salvia* recorded a long history of uses in nutritional, culinary, pharmacological and medicinal purposes. Abundant bioactive compounds content reported in sage seem to be responsible for its antimicrobial, antioxidant and pharmacological properties, including cytotoxicity against human cancer cells (Xavier and Pereira-Wilson, 2016). *S. officinalis* in food preparation was frequently reported as bio-preservative against Gram+ and Gram- bacteria (Al-terehi et al., 2013; Tajkarimi et al., 2010). Furthermore, studies highlighted the effect of *Salvia* in preventing and controlling various diseases naturally in a safe manner, and German Commission E (a governmental regulatory agency) has accepted the use of *S. officinalis* for a number of medical applications (Ghorbani and Esmaeilizadeh, 2017).

Exploring natural plants as anticancer agents, is one of the fastest growing interest of research (Seca and Pinto, 2018). Phytochemicals such as polyphenols, alkaloids, triterpenes and saponins, are biologically active compounds that have shown promising anticancer properties in both *in vitro* and *in vivo* based on anti-proliferative activity on tumor cells and antioxidant scavenging activity (Xavier and Pereira-Wilson, 2016; Xavier et al., 2009). In addition, they found to possess antimicrobial, anti-mutagenic, anti-inflammatory, and anti-cholinesterase properties. Currently, the demand has increased in food and pharmaceutical industries for these compounds as alternative natural approaches seeking safety (Yadav et al., 2017).

The objective of the present study was the assessment of nutritional and chemical compositions of sage leaves (*Salvia officinalis* L.) in order to highlight constituents relevant to antioxidant, antimicrobial and anticancer potentials against human breast cancer MDA-MB-231 cell line, and determine the safe dose to facilitate its application in functional foods and dairy products.

2. MATERIAL AND METHODS

2.1. Plant material and preparation

Common sage leaves (*Salvia officinalis* L.) was obtained from a reliable spice dealer, Alexandria, Egypt. To prepare sage aqueous extract (SAE), dried grounded leaves was extracted in Milli-Q water with ratio (1:20 w/v), stirred for 3 h at temperature (60°C) and centrifuged (Pro-Centrifuge, Centurion Science Limited, UK) for 20 min at 5500 rpm. The aqueous extract was filtered, freezed and then dried using vacuum freeze-dryer (Model FDF 0350, Korea) according to Garcia (Garcia et al., 2016) with modifications.

The plant leaves has been checked with

<http://www.theplantlist.org/tpl1.1/search?q=salvia+officinalis>

2.2. Proximate analysis

2.2.1. Nutrients content

Percent of moisture and ash content was determined as described by (AOAC, 1990). The protein content was determined by Kjeldahl method as described by (AOAC, 1990), and calculated according to equation; Protein= N x 6.25 to convert nitrogen content into protein content.

The crude fat content was determined using an Ankom^{XT10} fat extractor (Ankom Technology Corp., Macedon, NY). Crude Fat contained within a food or feed sample can be calculated using the following formula:

$$\% \text{ Crude fat} = \frac{100 \times (W2 - W3)}{W1}$$

Where: W1 = Original weight of sample, W2 = Weight of pre-dried sample and filter bag, W3 = Weight of dried sample and filter bag after extraction.

Crude fibre content was determined using Crude Fiber Analysis in Feeds - Filter Bag Technique A2000 - AOCS Approved Procedure Ba 6a-05 (Ankom Technology Corp., Macedon, NY). Crude fibre percent according to the formula:

$$\%Crude\ fibre = \frac{W3 - W1}{W2} \times 100$$

Where: W1 = Weight of pre-dried filter bag, W2 = Original weight of sample, W3 = Weight of filter bag contains organic residue remaining after digesting <https://www.ankom.com>

Total carbohydrate by difference was calculated by the following formula;

$$100 - (\text{weight in g [protein + fat + water + ash] in 100 g of food}) \text{ (WHO/FAO, 2002)}$$

It should be clear that carbohydrate estimated in this fashion includes fibre.

2.2.2. Electrolytes and minerals concentrations

Concentrations of electrolytes; sodium and potassium and minerals; zinc, iron, magnesium and calcium were determined using Atomic Absorption Spectrometry analytical technique according to (Beatty and Kerber, 1993). Concentration of phosphorus was determined using simplified colorimetric ammonium paramolybdate-vanadate method (yellow method), the colour intensity was measured at wave length 420 nm (T80 UV/VIS Spectrophotometer, PG Instruments Ltd) according to (Olsen and Sommers, 1982). Phosphorus concentrations were calculated using the formula;

$$\%P = P \text{ from st. curve} \times \frac{V1}{W_{sample}} \times \frac{1}{V2} \div 10\,000$$

Where: V1: Volume of stock standard solution of sample (50 mL), V2: Volume of stock standard solution of sample used in measurements (5mL), W_{sample}: Weight of sample (≈0.5 gm).

2.2.3. B Vitamins content

Concentration of some B Vitamins; Niacin Vitamin B3, Pyridoxine Vitamin B6 and Vitamin B12 was determined quantitatively via High Performance Liquid Chromatography (HPLC) at Food Safety and Quality Control Lab (FSQCLab), Faculty of Agriculture, Cairo University, Egypt, according to (Agilent Application Note, Publication number 5989-9313EN 2008). Agilent 1260 infinity HPLC Series (Agilent, USA), equipped with Quaternary pump, a

Kinetex XB-C18 column 100 mm X 4.6 mm (Phenomenex, USA), operated at 35°C. The separation was achieved using a binary linear elution gradient with (A) 25 mM NaH₂PO₄ pH 2.5 (v/v), (B) methanol. The injected volume was 20 µL. VWD detector was used for at 220 nm for vitamins B3, B6 and B12.

2.3. High Performance Liquid Chromatography (HPLC) of main components

For determining amino acid content, acid hydrolysis was carried out according to Csomos (Csomos and Simon-Sarkadi, 2002). Free amino acids extraction was carried out according to Emam (Emam, 2011). Amino acid analysis was carried out using Automatic Amino Acid Analyzer (AAA 400 INGOS Ltd). amino acids required pattern values are according to Food and Agriculture Organization (FAO) (FAO, 1985), while Amino Acid Score (AAS%) or Chemical Score (CS) was calculated as “Percentage of adequacy” as follows;

$$\text{Amino Acid Score} = \frac{\text{mg of amino acid in 1 g test protein}}{\text{mg of amino acid in requirement pattern}} \times 100$$

Fatty acids was determined via saponified and unsaponified fat separation according to AOAC (AOAC, 2005).

Phenolic compounds of sage leaves was identified according to Croci (Croci et al., 2009), using the following standards; gallic acid, caffeic acid, coumaric, syringic acid, vanillin, cinnamic acid, pyrogallol, catechin, quercetin and rutin.

2.4. Phytochemical, phenolic and flavonoid content

Phytochemical screening analysis for SAE were; Wagner’s method, Fehling’s test, foam test and Salkowki’s test and were performed according to Wadood (Wadood et al., 2013).

Total phenolic content assay was performed on SAE according to Singleton (Singleton et al., 1999) using Folin-Ciocalteu reagent. The total phenolic content was expressed as gallic acid equivalents (GAE mg/g). All measurements were performed in triplicates.

Total flavonoid contents of SAE was determined using colorimetric method as described by Sakanaka (Sakanaka et al., 2005). Results expressed in term of standard used for the quantification of flavonoids. All samples were performed in triplicate.

2.5. Pharmacological potentials

2.5.1. Antioxidant activity

Antioxidant scavenging activity of SAE was determined using two different methods; DPPH (2,2-diphenyl-1-picrylhydrazyl) assay that was performed as described by Brand-Williams (Brand-Williams et al., 1995), and reducing power assay as described by Ferreira (Ferreira et al., 2007).

2.5.2. Antimicrobial screening

Antimicrobial potentials of SAE was screened against nine pathogenic strains using agar well diffusion assay as described by Hamad (Hamad et al., 2017). The nine tested pathogenic organisms were; five Gram-positive strains; *Staph. aureus* NCTC10788, *Staph. pyogenes* EMCC1772, *St. mutants* EMCC1815, *B. subtilis* DB100 and *Clostridium botulinum* ATCC3584, three Gram-negative strains; *E. coli* BA12296, *E. coli* ATCC25922 and *K. pneumonia* ATCC12296 and one yeast strain; *Candida albicans* ATCCMYA2876. The pathogenic strains were obtained from Microbiological Recourses Center (Cairo MERCIN), Egypt Microbial Culture Collection (EMCC), Ain Shams University, Cairo, Egypt, and were maintained at Food Technology Dept., City of Scientific Research and Technological Applications. The minimum inhibitory concentration (MIC) was assessed within the concentrations of 50, 25, 12.5 and 6.25 mg/ mL of sage extract.

2.5.3. Cytotoxicity and anticancer assay

This test was used to evaluate cytotoxicity. Membrane of viable cells capture the neutral red, therefore after subsequent lysis, absorbance can be used as a measure of cell viability (Borenfreund and Puerner, 1985). The stock solution of NR (3.3 g/l) was diluted to 1/100 in

the cell culture medium and the extracting solution consisted of 50% (v/v) ethanol in Milli-Q water with 1% (v/v) acetic acid. MDA-MB-231 Triple negative breast cancer cells were grown for 48h in presence of SAE, (50–600µg) after incubation 150 µL of freshly prepared neutral red solution pre-warmed to 37 °C was added to each well and all plates were incubated at 37 °C for additional 4 h. The cells were washed twice with PBS, 150µl of destain solution were added to each well and plates were shaken for 15 min. The absorbance was determined at 540 nm using a Microplate Elisa Reader while, light microscope (Olympus IX70-58F2 Multi-parameter Fluorescence Microscope) was used for monitoring the morphological changes in treated MDA-MB-231 cells after 48 h of incubation.

2.6. Statistical analysis

Statistical analysis was performed using Analytical Software SPSS® 13.0 (Statistical Package for the Social Sciences) (2005).

3. RESULTS

3.1. Proximate composition

Table (1) illustrated proximate composition of sage leaves (*Salvia officinalis*) including, nutrients, minerals and B vitamins. Proximate nutrient composition of sage (*Salvia officinalis*), indicated that sage is a high-carbohydrate high-fiber nutrient source plant. Another advantage is the relatively high protein (17g/100 g) with low-fat content (3.46 g/100g). High ash content indicated sage richness in minerals. The sage showed remarkable concentration of iron (84.037 mg/100gm), calcium (894.273 mg/100g) and zinc (5.472 mg/100g). On the other hand, sage showed poor content of sodium and potassium (19.893, 34.605 mg/100g). Vitamin B concentrations of sage were (3.258, 1.545, 0.329 mg/100g) of vitamins B3, B6 and B12 respectively.

3.2. HPLC analysis

3.2.1. Amino acids profile

Amino acid content of sage (*Salvia officinalis*) are represented in (Table 2). Obtained results showed that the main essential amino acid (EAA) in sage was lysine with content of (10.44 g/100g). Sage also provided considerable content of phenylalanine and leucine (0.74 and 0.45 g/ 100g respectively), limiting content of valine, threonine, histidine and isoleucine. On the other hand, sage exhibited generous content of non-essential amino acids, glycine, glutamic, alanine, and aspartic, (1.02, 0.83, 0.75 and 0.53 g/ 100g respectively).

3.2.2. Fatty acids content

Fatty acids (FA) profile of sage (*Salvia officinalis*) was exhibited in (Table 3). Sage showed USFAs/ SFAs ratio of 0.32 due to the superiority of SFAs. The highest content of SFAs belonged to tridecylic (C13:0) (11.33 g/100g of total FA), followed by equal amounts of lauric (C12:0) and myristic (C14:0) (7.6 g/100g of total FA). UFAs 18:3 ω -3, C18:2 ω -6 and C18:1 ω -9 was present with values of (6.46, 4.40 and 3.13 g/100g of total FA respectively).

3.2.3. Phenolic compounds profile

Table (4) and Fig. (2) demonstrated phenolic compounds concentrations of sage aqueous extracts. Within the ten used phenolic compound standards, three compounds were not detected in SAE; gallic acid, catechin and pyrogallol. Quercetin and cinnamic acid showed the highest concentrations (604.7575 and 390.3971 μ g/mL respectively) representing (0.6756 and 0.4361%) of sage TPC (Table 5). The rest of tested phenolic compounds standards; caffeic acid, syringic acid, coumaric acid, vanillin and rutin were existed in little concentrations (0.6132, 0.5134, 0.5134, 0.0856 and 0.0766 μ g/mL respectively).

3.3. Pharmacological potentials

3.3.1. Phenolic and flavonoids content and antioxidant powers

Qualitative screening of sage phytochemicals results showed presence of a wide range of constituents include, carbohydrate (reducing sugars), volatile oil, phenolic compounds,

flavonoids/ flavones, tannins, steroids, terpenoids and amino acids which are mostly responsible for antioxidant, antimicrobial and cytotoxic activities.

Table (5) illustrated phenolic, flavonoid content and antioxidant scavenging potentials of sage aqueous extract and Sage showed a total phenolic content of 88.57 ± 3.14 mg GAE/ g, while total flavonoids was 37.13 ± 1.83 mg catechol equivalent/ g. Antioxidant and scavenging potentials of aqueous extract of sage; DPPH results represented with inhibitory concentration at which 50% of DPPH radicals are scavenged (IC_{50} mg/mL) and reducing power represented with effective concentration at which the absorbance is 0.5 (EC_{50} mg/mL). Sage leaves water extract showed IC_{50} and EC_{50} values (27.52 and 239.52 mg/ mL respectively).

3.3.2. Antimicrobial activity

Table (6) presented inhibition zone diameter in mm of different concentrations of sage (*Salvia officinalis*) aqueous extract and Minimal Inhibitory Concentration (MIC) for each strain. Results revealed that sage leaves extract owned antimicrobial activity against all tested pathogenic strains with varied MICs. According to MICs, the highest antimicrobial interaction of sage extract was against *Staph. aureus* NCTC10788 with inhibition zone diameter started from 20 mm with concentration of 50 mg/mL, and ended with 6 mm with concentration of 6.25 mg/mL, which was the lowest MIC amongst the five Gram-positive strains. Whilst, sage showed MIC of 12.5 mg/mL against the other four Gram-positive strains with inhibition zone diameters started from 19 – 13.5 mm with concentration of 50 mg/mL.

Antifungal activity of sage aqueous extract was obvious against *Candida albicans* ATCCMYA2876 with MIC of 6.25 mg/ mL, the inhibition zone diameter reached 22 mm with concentration of 50 mg/mL.

3.3.3. Cytotoxicity and anticancer potentials

Figure (1) showed anti-proliferative activity of sage (*Salvia officinalis*) aqueous extract (SAE) against MDA-MB-231 cell line via neutral red assay (IC_{50} of 300 μ g/mL). Characteristics of

dying cells clearly appeared in sage extract treated cells indicating its effect on cell viability with an IC_{50} of 300 μ g. Figure 2 (a and b), illustrated morphological changes in MDA-MB-231 Triple negative breast cancer cell following treatment with sage aqueous extract (SAE) with concentration of 300 μ g/ mL for 48 h. Light microscopy observation of SAE-treated cells revealed morphological changes induced by the extract. As shown on figure 2b, MDA-MB-231 treated cells with concentration of 300 μ g/ mL SAE, underwent visible morphological changes after 48 h of incubation, where treated cells appeared smaller and rounded (characteristics of dying cells) (Fig.2b, arrows), in addition to decrease of neutral red uptake in a concentration-dependent manner.

4. DISCUSSION

Plant-fiber-enriched foods are reported to improve glucose and lipid metabolism, they lower serum cholesterol concentrations in patients with diabetes, in addition to multiple effects on different cardiovascular risk factors (De Natale et al., 2009). The electrolytes results emphasized that sage contains 3 times more calcium and 5.5 times more zinc than milk. USDA (USDA, 2016) reported different values for sage mineral except for zinc and sodium content. Alterations in essential trace elements like Fe, Zn and Ca may play an important role in the pathogenesis of complex diseases including cancers. Dietary calcium intake found to be associated with a reduced incidence of cancer, while iron, is involved in many processes including DNA, RNA and protein synthesis, and Zn may represent an independent risk factor for cancer and therefore a possible target for prevention (Al Faris and Ahmad, 2011; Singh et al., 2016). Obtained vitamin results indicated that sage could be considered a good source of vitamins B6 and B12 and encourage suggestions of using as food supplement especially for vegetarians and elderly. What is most important that, Vitamin B6 and B12 was reported to sensitize a large panel of cancer cell lines to apoptosis (Pandey et al., 2013; Saghiri et al., 2017). Obtained values of B vitamins fluctuated between agreement and disagreement with

(Lovik et al., 2009; USDA, 2016). Nutritional profile may aid to explain the current increased demand on *Salvia* species for nutritional and pharmacological industries (Yadav et al., 2017). Many biological functions requiring lysine; include synthesis of connective tissues and carnitine, conversion of fatty acids to energy that support healthy growth in children and maintenance of healthy immune function especially in the management of Herpes simplex virus (HSV) infections (Hall et al., 2018). Lysine also found to enhance calcium absorption preventing Osteoporosis. On food technology wise; lysine is involved in the browning reaction applied to foods such as pastries and desserts (Braun and Cohen, 2014). Through obtained results, it is recommended to mix sage with foods rich in limited EAA. Apart from amino acid, starvation diets theories - that stimulates autophagy in cancer- which is complex and likely context-dependent, the maintenance of the intracellular level of amino acids is crucial for cellular homeostasis. Deficiency or imbalance of essential amino acids in the diet was reported to produce profound depression in the immune resistance of the host animal to tumors (Lashinger et al., 2016; Saito et al., 2018).

Increased intake of lauric, myristic or palmitic acid was reported to raise serum total, LDL and HDL cholesterol levels, and lowered triglyceride levels, while lauric acid alone reduced the total cholesterol to HDL cholesterol and LDL cholesterol to HDL cholesterol ratios (Mensink, 2016). Unsaturated fatty acids (UFAs) was reported to exert antineoplastic activity by inducing apoptotic cell death in human cancer cells through increasing the sensitivity of tumor cells to conventional therapies (D'Eliseo and Velotti, 2016). In this context, USDA (USDA, 2016) reported vitiated values about sage. Reported laboratory studies indicated clear role for omega-3 PUFAs in preventing cancer development at various stages (de Aguiar Pastore Silva et al., 2015; Eltweri et al., 2017). Generally, it is difficult to match results of nutrients and constituents with earlier researchers since *Salvia* species have various

compositions depending on the genetic, climatic, seasonal, and environmental factors (Hamidpour et al., 2014).

Caffeic acid and its derivatives including rosmarinic acid was reported to be the most representative phenolic compounds in *Lamiaceae*, *S. officinalis* extracts (Olah et al., 2004). Anticancer activities of cinnamic acid derivatives include induction of apoptosis by irreversible DNA damage leading to cell death was reported (Niero and Machado-Santelli, 2013). Since phenolic compounds are the key for several therapeutic properties, quantitative identification of sage leaves phenolic compounds indicated pharmacological potentials to be useful as a source for novel applications in functional food and pharmaceutical industries (Bahadori et al., 2017).

Earlier studies on sage reported the presence of obtained phytochemicals classes of active compounds (Ghorbani and Esmailizadeh, 2017; Hamidpour et al., 2014). Obtained results of phenolic and flavonoid content were in context with Özcan (Özcan and Özkan, 2015), while it disagreed with Gîrd and Pop (Gîrd et al., 2014; Pop et al., 2014). Sage was reported to possess the greatest antioxidant capacity amongst fresh herbs, due to its phenolic and flavonoids content which related for multiple health benefits, including reduction in cancer risk and modification of tumor behavior (Kaefer and Milner, 2011; Scapin et al., 2016).

Activity of sage against filamentous fungi and yeasts such as *Candida albicans* was previously reported by Sirirat and Jelena (Sirirat and Jelena, 2010), and thus may be useful in the treatment of different kinds of fungi in human. Antimicrobial results can nominate sage as a promising natural antimicrobial agent which is in accordance to the results of (Zomorodian et al., 2015). Reasonable antimicrobial and antioxidant activities encouraged further investigation of anti-breast cancer activities of sage extract.

Alterations of the MDA-MB-231 cell surface or of the sensitive lysosomal membrane lead to lysosomal fragility and other changes that gradually become irreversible. The main advantage

of neutral red assay is that it detects only viable cells (Babich and Borenfreund, 1987). Previous researches reported the antitumor properties of sage extract which probably in correlation to its **announced** antioxidant activity as in other plant extracts (Garcia et al., 2016; Rupasinghe et al., 2016). Sage bioactive compounds, Linalyl acetate (Ly), Terpeniol (Te) and Camphor (Ca), has been reported to have synergistic effect on two human colon cancer cell lines HCT-116 (p53 (+/+)) and p53 (-/-). Apoptosis by Ly + Te + Ca, in p53 (+/+) cells (Itani et al., 2008). Despite progress in cancer research, the biological active components and antitumor mechanisms have not yet been exactly determined, but mitochondrial membrane potential decay is one of the accompanying processes of the intrinsic apoptotic pathway activation (Jantová et al., 2014). On the other hand, concerns and limitations for the amount and quantity used of sage attributed to the predominant volatile neurotoxic components; camphor, α - and β -thujone, which are most present phytochemicals in *Salvia officinalis* leaves. A daily intake of 5.0 mg/ person is acceptable for a maximum duration of use of 2 weeks, and the overdose is more than 15 g of sage leaves per day (Iatrou et al., 2013). It is noteworthy that sage use during pregnancy and lactation is not recommended (Wharf, 2009). Our findings encourage further studies of sage components as promising antitumor therapy against breast cancer.

4. Conclusion

In conclusion, *Salvia officinalis* possessed **abundant** antioxidant, antimicrobial and anti-breast cancer activities due to its rich combination of nutrients and constituents. Sage considered a good source of calcium, iron, zinc, vitamins B6 and B12, essential amino acids, lysine, phenylalanine and leucine and **unsaturated fatty acids (UFAs)** ω -3, ω -6 and ω -9, which all reported to have a role as anti-cancer in different mechanisms. Presence of a wide range of constituents include, volatile oil, flavonoids/ flavones, tannins, steroids, terpenoids combined with significant amounts of phenolic compounds, especially quercetin and cinnamic,

interpreted its high antioxidant powers. In addition, to marked antimicrobial potentials that showed against all nine tested pathogens, that could be beneficially used in food fortification for preservation and pharmacologically as anti-tumour alternative natural supplement. SAE performed anti-proliferative activity against MDA-MB-231 cell line with IC₅₀ of 300 µg/mL. These results encourage further research on sage components as promising antitumor agent. Sage leaves possessed variable nutrients and active compounds that reflected on its vast nutritional and pharmacological potentials and support it as a functional nutritional additive and a natural bio-preservative in food and dairy preparations.

Acknowledgment

Authors appreciate the support provided by, Food Technology Dept., Arid Lands Cultivation Research Institute (ALCRI) and Genetic Engineering and Biotechnology Research Institute (GEBRI), City of Scientific Research and Technological Applications (SRTA-City), Alexandria, Egypt, to achieve this work.

References

- Agilent_Technologies (2008). Agilent Application Note, Publication number 5989-9313EN.
- de Aguiar Pastore Silva, J., Emilia de Souza Fabre, M., and Waitzberg, D.L. (2015). Omega-3 supplements for patients in chemotherapy and/or radiotherapy: A systematic review. Clin. Nutr. 34, 359–366.
- Al-terehi, M., Al-saadi, A.H., Al-musawi, H. turki, Zaida, H. k, Behjet, R.H., and Haleem, Z. (2013). Synergism Effect of Antibacterial Activity of Some Medicinal Plants. J. Biol. Med. Sci. 1, 46–53.
- AOAC (1990). AOAC: Official Methods of Analysis (Association of Official Analytical Chemists. INC., Arligton, Virginia, USA.).
- AOAC (2005). AOAC: Official Methods of Analysis (AOAC INTERNATIONAL, Gaithersburg, Maryland 20877-2417, USA).

Babich, H., and Borenfreund, E. (1987). Structure-activity relationship (SAR) models established in vitro with the neutral red cytotoxicity assay. *Toxicol. Vitro.* 1, 3–9.

Bahadori, M.B., Asghari, B., Dinparast, L., Zengin, G., Sarikurkcu, C., Abbas-Mohammadi, M., and Bahadori, S. (2017). *Salvia nemorosa*: A novel source of bioactive agents with functional connections. *LWT - Food Sci. Technol.* 75, 42–50.

Beaty, R.D., and Kerber, J.D. (1993). *Concepts, Instrumentation and Techniques in Atomic Absorption Spectrophotometry* (The Perkin-Elmer Corporation, Norwalk, CT, U.S.A.).

Borenfreund, E., and Puerner, J.A. (1985). Toxicity determined in vitro by morphological alterations and neutral red absorption. *Toxicol. Lett.* 24, 119–124.

Brand-Williams, Cuvelier, M.E., and Berset, C. (1995). Use of a Free Radical Method to Evaluate Antioxidant Activity. *Food Sci. Technol.* 28, 25–30.

Braun, L., and Cohen, M. (2014). *Herbs & natural supplements, Volume 2. An evidence-based guide.* (Churchill Livingstone Australia, Elsevier).

Croci, A.N., Cioroiu, B., and Lazar, D. (2009). HPLC evaluation of phenolic and polyphenolic acids from propolis. *Farmacia LVII*, 52–57.

Csomos, E., and Simon-Sarkadi, L. (2002). DETERMINATION OF BIOLOGICALLY ACTIVE COMPOUNDS. 46, 73–81.

D’Eliseo, D., and Velotti, F. (2016). *Omega-3 Fatty Acids and Cancer Cell Cytotoxicity: Implications for Multi-Targeted Cancer Therapy.*

Eltweri, A.M., Thomas, A.L., Metcalfe, M., Calder, P.C., Dennison, A.R., and Bowrey, D.J. (2017). Potential applications of fish oils rich in omega-3 polyunsaturated fatty acids in the management of gastrointestinal cancer. *Clin. Nutr.* 36, 65–78.

Emam, S.S. (2011). Bioactive constituents of *Atriplex halimus* plant Shalabia. *J. Nat. Prod. - India* 4, 25–41.

FAO (1985). Food and Agriculture Organization (FAO) (Food and Agriculture Organization

of the United States/ World Health Organization/ United Nations University, Energy and Protein requirements. Report of a joint FAO/WHO/UNU meeting. World Health Organization, Geneva).

Al Faris, N.A., and Ahmad, D. (2011). Distribution of trace elements like calcium, copper, iron and zinc in serum samples of colon cancer - A case control study. J. King Saud Univ. - Sci. 23, 337–340.

Ferreira, I.C.F.R., Baptista, P., Vilas-Boas, M., and Barros, L. (2007). Free-radical scavenging capacity and reducing power of wild edible mushrooms from northeast Portugal: Individual cap and stipe activity. Food Chem. 100, 1511–1516.

Garcia, C.S.C., Menti, C., Lambert, A.P.F., Barcellos, T., Moura, S., Calloni, C., Branco, C.S., Salvador, M., Roesch-Ely, M., and Henriques, J.A.P. (2016). Pharmacological perspectives from brazilian *Salvia officinalis* (Lamiaceae): Antioxidant, and antitumor in mammalian cells. An. Acad. Bras. Cienc. 88, 281–292.

Ghorbani, A., and Esmaeilizadeh, M. (2017). Pharmacological properties of *Salvia officinalis* and its components. J. Tradit. Complement. Med. 8–15.

Gîrd, C.E., Nencu, I., Costea, T., Duțu, L.E., Popescu, M.L., and Ciupitu, N. (2014). Quantitative analysis of phenolic compounds from *Salvia officinalis* L. leaves. Farmacia 62, 649–657.

Hall, C.J., Soares da Costa, T.P., and Soares da Costa, T (2018). Lysine: biosynthesis, catabolism and roles. WikiJournal Sci. 1, 4.

Hamad, G.M., Darwish, A.M.G., Abu-Serie, M.M., and El Sohaimy, S.A. (2017). Antimicrobial, Antioxidant and Anti-inflammatory Characteristics of Combination (*Cassia fistula* and *Ocimum basilicum*) Extract as Natural Preservative to Control & Prevent Food Contamination. J. Food Nutr. Res. 5, 771–780.

Hamidpour, M., Hamidpour, R., Hamidpour, S., and Shahlari, M. (2014). Chemistry,

Pharmacology, and Medicinal Property of Sage (*Salvia*) to Prevent and Cure Illnesses such as Obesity, Diabetes, Depression, Dementia, Lupus, Autism, Heart Disease, and Cancer. *J. Tradit. Complement. Med.* 4, 82–88.

Iatrou, G., Lamari, F., Dimitrellos, G., and Tsakiri, M. (2013). Catalogue of aromatic and medicinal plants (European Territorial Cooperation Programme Greece - Italy. Eptalofos, S A, Athens, Greece.).

Itani, W.S., El-Banna, S.H., Hassan, S.B., Larsson, R.L., Bazarbachi, A., and Gali-Muhtasib, H.U. (2008). Anti colon cancer components from lebanese sage (*Salvia libanotica*) essential oil: Mechanistic basis. *Cancer Biol. Ther.* 7, 1765–1773.

Jantová, S., Hudec, R., Sekretár, S., Kučerák, J., and Melušová, M. (2014). *Salvia officinalis* L. extract and its new food antioxidant formulations induce apoptosis through mitochondrial/caspase pathway in leukemia L1210 cells. *Interdiscip. Toxicol.* 7, 146–153.

Kaefer, C.M., and Milner, J.A. (2011). Herbs and Spices in Cancer Prevention and Treatment. *Herb. Med. Biomol. Clin. Asp.* 19, chapter 16.

Lashinger, L.M., O’Flanagan, C.H., Dunlap, S.M., Rasmussen, A.J., Sweeney, S., Guo, J.Y., Lodi, A., Tiziani, S., White, E., and Hursting, S.D. (2016). Starving cancer from the outside and inside: separate and combined effects of calorie restriction and autophagy inhibition on Ras-driven tumors. *Cancer Metab.* 4, 18.

Lovik, M., Marchelli, R., Martin, A., and Moseley, B. (2009). Scientific Opinion of the Panel on Dietetic Products Nutrition and Allergies on a request from the European Commission on the safety of ‘Chia seed (*Salvia hispanica*) and ground whole Chia seed’ as a food ingredient. *EFSA J.* 996, 1–26.

Mensink, R.P. (2016). Effects of saturated fatty acids on serum lipids and lipoproteins: a systematic review and regression analysis.

De Natale, C., Annuzzi, G., Bozzetto, L., Mazzarella, R., Costabile, G., Ciano, O., Riccardi,

G., and Rivellesse, A.A. (2009). High-Carbohydrate / High-Fiber Diet Versus Carbohydrate Diet on Postprandial Lipids in Type 2 Diabetic Patients. *Diabetis Care* 32, 2168–2173.

Niero, E.L. de O., and Machado-Santelli, G. (2013). Cinnamic acid induces apoptotic cell death and cytoskeleton disruption in human melanoma cells. *J. Exp. Clin. Cancer Res.* 32, 31.

Olah, N.K., Hanganu, D., Oprean, R., Mogoşan, C., Dubei, N., and Gocan, S. (2004). Selective extraction of caffeic acid derivatives from *Orthosiphon stamineus* Benth. (Lamiaceae) leaves. *J. Planar Chromatogr. - Mod. TLC* 17, 18–21.

Olsen, S.R., and Sommers, L.E. (1982). Phosphorus. In: Page A.L., et al (eds), *Methods of Soil Analysis*. (Part 2, 2nd edn, Agron Monogr 9.ASA and ASSA, Madison WI), pp. 403–430.

Özcan, M.M., and Özkan, G. (2015). Determination of antioxidant activity and total phenol contents of two *Salvia* extracts. *Indian J. Tradit. Knowl.* 14, 226–230.

Pandey, S., Garg, P., Lim, K.T., Kim, J., Choung, Y.H., Choi, Y.J., Choung, P.H., Cho, C.S., and Chung, J.H. (2013). The efficiency of membrane transport of vitamin B6 coupled to poly(ester amine) gene transporter and transfection in cancer cells. *Biomaterials* 34, 3716–3728.

Pop, A. V, Tofana, M., Socaci, S., Varban, D., Nagy, M., Bors, M., and Sfechis, S. (2014). Evaluation of antioxidant activity and phenolic content in different *Salvia officinalis* L. extracts. *Bull. UASVM Food Sci. Technol.* 71, 195–199.

Rupasinghe, V., Jiang, Y., and Zhang, L. (2016). The anticancer properties of phytochemical extracts from *Salvia* plants. *Bot. Targets Ther.* Volume 6, 25.

Saghiri, M.A., Asatourian, A., Ershadifar, S., Moghadam, M.M., and Sheibani, N. (2017). Vitamins and regulation of angiogenesis: [A, B1, B2, B3, B6, B9, B12, C, D, E, K]. *J. Funct. Foods* 38, 180–196.

Saito, Y., Moriya, S., Kazama, H., Hirasawa, K., Miyahara, K., Kokuba, H., Hino, H.,

Kikuchi, H., Takano, N., Hiramoto, M., et al. (2018). Amino acid starvation culture condition sensitizes EGFR-expressing cancer cell lines to gefitinib-mediated cytotoxicity by inducing atypical necroptosis. *Int. J. Oncol.* 1165–1177.

Sakanaka, S., Tachibana, Y., and Okada, Y. (2005). Preparation and antioxidant properties of extracts of Japanese persimmon leaf tea (kakinoha-cha). *Food Chem.* 9, 569–575.

Scapin, G., Schmidt, M.M., Prestes, R.C., and Rosa, C.S. (2016). Phenolics compounds, flavonoids and antioxidant activity of chia seed extracts (*Salvia hispanica*) obtained by different extraction conditions. *Int. Food Res. J.* 23, 2341–2346.

Seca, A., and Pinto, D. (2018). Plant Secondary Metabolites as Anticancer Agents: Successes in Clinical Trials and Therapeutic Application. *Int. J. Mol. Sci.* 19, 263.

Singh, B.P., Dwivedi, S., Dhakad, U., Murthy, R.C., Choubey, V.K., Goel, A., and Sankhwar, S.N. (2016). Status and Interrelationship of Zinc, Copper, Iron, Calcium and Selenium in Prostate Cancer. *Indian J. Clin. Biochem.* 31, 50–56.

Singleton, V.L., Orthofer, R., and Lamuela-Raventos, R.M. (1999). Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin–Ciocalteu reagent. *Oxidants and antioxidants part A.* (Academic Press).

Sirirat, D., and Jelena, P. (2010). Bacterial inhibition and antioxidant activity of kefir produced from Thai jasmine rice milk. *Biotechnology* 9, 332–337.

Tajkarimi, M.M., Ibrahim, S.A., and Cliver, D.O. (2010). Antimicrobial herb and spice compounds in food. *Food Control* 21, 1199–1218.

USDA (2016). National Nutrient Database for Standard Reference Release 28 slightly revised May , 2016 Full Report (All Nutrients) 12006 , Seeds , chia seeds , dried.

Wadood, A., Ghufuran, M., Jamal, S.B., Naeem, M., Khan, A., and Ghaffar, R. (2013). Phytochemical Analysis of Medicinal Plants Occurring in Local Area of Mardan. *Biochem. Anal. Biochem.* 2, 2–5.

Wharf, C. (2009). Community Herbal Monograph on *Salvia officinalis* L., Folium. Reproduction 1–8.

WHO/FAO (2002). Food energy – methods of analysis and conversion. In Fao Food and Nutrition Paper, p. 93.

Xavier, C.P.R., and Pereira-Wilson, C. (2016). Medicinal plants of the genuses *Salvia* and *Hypericum* are sources of anticolon cancer compounds: Effects on PI3K/Akt and MAP kinases pathways. *PharmaNutrition* 4, 112–122.

Xavier, C.P.R., Lima, C.F., Fernandes-Ferreira, M., and Pereira-Wilson, C. (2009). *Salvia fruticosa*, *salvia officinalis*, and rosmarinic acid induce apoptosis and inhibit proliferation of human colorectal cell lines: The role in MAPK/ERK pathway. *Nutr. Cancer* 61, 564–571.

Yadav, A., Joshi, A., Kothari, S.L., Kachhwaha, S., and Purohit, S. (2017). Review Article Medicinal, Nutritional and Industrial Applications of *Salvia* species: A Revisit. *Int. J. Pharm. Sci. Rev. Res.* 43, 27–37.

Zomorodian, K., Ghadiri, P., Saharkhiz, M.J., Moein, M.R., Mehriar, P., Bahrani, F., Golzar, T., Pakshir, K., and Fani, M.M. (2015). Antimicrobial activity of seven essential oils from Iranian aromatic plants against common causes of oral infections. *Jundishapur J. Microbiol.* 8, e17766.

Table (1): Proximate composition of sage (*Salvia officinalis*)

Component	Unit	Content (per 100g)
Nutrients		
Protein	g	17.01±0.64
Crude fat	g	3.46±0.28
Carbohydrate, by difference	g	57.87
Crude fiber	g	35.96±0.46
Moisture	g	9.07±0.30
Ash	g	12.59±0.76
Electrolytes		

Sodium, Na	mg	19.893
Potassium, K	mg	34.605
Minerals		
Calcium, Ca	mg	894.273
Iron, Fe	mg	84.037
Magnesium, Mg	mg	19.601
Zinc, Zn	mg	5.472
Phosphorus, P	mg	Traces
B Vitamins		
Niacin, B3	mg	3.258
Pyridoxine, B6	mg	1.545
Cobalamin, B12	mg	0.329

Nutrients data represented are the mean $\pm$ standard deviation, n=3

Table (2): Amino acids content of sage (*Salvia officinalis*)

Amino Acid	Symbol	Content (g/100g)
<i>Essential amino acids</i>		
Histidine	His	0.17
Leucine	Leu	0.45
Isoleucine	Ile	0.13
Lysine	Lys	10.44
Methionine + cystine	Met+ Cys	0.03
Phenylalanine + tyrosin	Phe+ Tyr	0.74
Threonine	Thr	0.17
Valine	Val	0.28
CS	-	-
<i>Non-essential amino acids</i>		
Alanine	Ala	0.75
Aspartic acid	Asp	0.53
Glutamic acid	Glu	0.83
Glycine	Gly	1.02
Proline	Pro	0.01
Serine	Ser	0.25
Tyrosine	Tyr	0.15

Table (3): Fatty acid profile of sage (*Salvia officinalis*)

<i>SFAs</i>		
Capric	C10:0	1.28
Undecylic	C11:0	6.07
Lauric	C12:0	7.60
Tridecylic	C13:0	11.33
Myristic	C14:0	7.64
Pentadecylic	C15:0	6.95
Palmitic	C16:0	1.44

Stearic	C18:0	0.93
Arachidic	C20:0	0.54
%of total fat		43.78
USFAs		
MUSFAs		
Oleic	C18:1 ω-9	3.13
PUSFAs		
Linoleic	C18:2 ω-6	4.40
α-Linolenic	C18:3 ω-3	6.46
%of total fat		13.99
USFAs/ SFAs		0.32
ω6/ ω3 ratio		0.68

SFAs; Saturated fatty acids, USFAs; Unsaturated fatty acids, MUFAs; Monounsaturated fatty acids, PUFAs; Polyunsaturated fatty acids

Table (4): HPLC analysis of phenolic compounds concentrations of sage aqueous extract

Phenolic compound	μg/mL
Caffeic acid	0.6132
Syringic acid	0.5134
Rutin	0.0766
Coumaric acid	0.4653
Vanillin	0.0856
Quercetin	604.7575
Cinnamic acid	390.3971

Table (5): Antioxidant and scavenging potentials of sage aqueous extract

Test	Unit	Values
Total phenolic	*mg /g	88.57±3.14
Total flavonoids	**mg /g	37.13±1.83
DPPH	^a mg/mL	27.52±7.63
Reducing power	^b mg/mL	239.52±0.41

Data represented in means of duplicates ± standard deviation

*Total phenolic was expressed as gallic acid equivalents (GAE) mg/ g sample

**Total flavonoids was expressed as mg catechol/g sample

^a IC₅₀ (mg/mL): Inhibitory concentration at which 50% of DPPH radicals are scavenged.

^b EC₅₀ (mg/mL): Effective concentration at which the absorbance is 0.5.

Table (6): Inhibition zone diameter in mm of different concentrations of sage (*Salvia officinalis*) aqueous extract and MIC for each strain

Pathogenic strain	Inhibition zone diameter (mm)**					
	7	50*	25*	12.50*	6.25*	MIC
Gram-positive bacteria						
<i>Staphylococcus aureus</i> NCTC10788	2	20.0	18.0	12.0	6.0	6.25
<i>Staphylococcus pyogenes</i> EMCC1772	1	13.5	10.0	5.5	ND	12.50
<i>Streptococcus mutants</i> EMCC1815	2	18.0	11.0	7.0	ND	12.50
<i>Bacillus subtilis</i> DB100	2	19.0	14.5	5.5	ND	12.50
<i>Clostridium botulinum</i> ATCC3584	2	15.5	11.0	7.5	ND	12.50
Gram-negative bacteria						
<i>Escherichia coli</i> BA12296	2	18.0	17.0	10.0	5.0	6.25
<i>Escherichia coli</i> ATCC25922	2	19.0	13.0	5.0	ND	12.50
<i>Klebseilla pneumonia</i> ATCC12296	2	18.0	13.0	5.0	ND	12.50
Yeast						
<i>Candida albicans</i> ATCCMYA2876	2	22.0	18.5	8.0	5.0	6.25

Data represented are average of duplicates, *Concentrations of extract and MIC are in mg/mL

**Diameter included 5 mm well diameter, MIC; Minimum Inhibition Concentration





Figure 1. Anti-proliferative activity of sage (*Salvia officinalis*) aqueous extract (SAE) against MDA-MB-231 cell line via neutral red assay (IC₅₀ of 300µg/mL)

a



b




Figure 2. HPLC graph of phenolic compounds concentrations of sage aqueous extract comparing to control.




Figure 3. Morphological changes in MDA-MB-231 Triple negative breast cancer cell following treatment with sage aqueous extract (SAE) with concentration of 300 µg/ mL for 48 h.

Mentions: Light microscopy observation of SAE-treated MDA-MB-231 Triple negative breast cancer cells revealed morphological changes induced by the extract. As shown on figure 2b, MDA-MB-231 treated cells with concentration of 300 µg/ mL SAE, underwent visible morphological changes after 48 h of treatment, where treated cells appeared smaller and rounded, characteristics of dying cells (Fig.2b, arrows) in addition to decrease of neutral red uptake in a concentration-dependent manner.