

Phytochemical Screening and Anti-diarrheal Activity of the Pulp and Seed Extracts of *Ziziphus mauritiana* in Rats.

Martha Asugu Mbahi^{1*}, Mary Asugu Mbahi², and Pigweh Isa Amos³.

¹Department of Biochemistry, Ahmadu Bello University, Nigeria.

²Department of Biological Sciences, Federal University of Technology Kashere, Gombe State.

³Department of Chemistry, School of Physical Sciences, Modibbo Adama University of Technology, Yola, Adamawa State, Nigeria.

*Corresponding Author: Email:asugum@yahoo.com.com

Abstract

In the present study, attempt was made to evaluate the phytochemical composition and anti-diarrheal activity of the seed and pulp extract of *Ziziphus mauritania*. The anti-diarrheal activity of the crude seed and pulp were evaluated using castor oil induced diarrheal model, charcoal meal test and anti-fluid accumulation test in rats. The result of phytochemical test indicated that tannins, flavonoids, saponin, cyanogenic glycosides, and terpenoids were present in both seed and pulp. In the castor oil induced model both the seed and pulp extract significantly prolonged diarrheal onset was observed in treated rats compared to the negative control. Similarly, in the fluid accumulation test, the extract of the seed and pulp produced a significant decline in volume of intestinal contents. Results from the charcoal meal test revealed that all the extract produced a significant anti-motility effect. Based on the findings of this work, the pulp extract of this plant possess anti-diarrheal properties and validates its use in traditional medicine for the treatment of diarrhea.

Key words: *Ziziphus Mauritania*, Diarrheal, Charcoal meal, Castor oil.

1.0 Introduction

The use of traditional medicines and medicinal plants in most developing countries as therapeutic agents for the maintenance of good health has been widely observed by UNESCO [1]. Modern pharmacopoeia still contains at least 25% drugs derived from plants and many others, which are synthetic analogues, built on prototype compounds isolated from plants. Interest in medicinal plants as a re-emerging health aid has been fuelled by the rising costs of prescription drugs in the maintenance of personal health and wellbeing and the bioprospecting of new plant-derived drugs. The growing recognition for medicinal plants use is due to several reasons, including escalating faith in herbal medicine [2]. Furthermore, an increasing reliance on the use of medicinal plants in the industrialized societies has been traced to the extraction and development of drugs and chemotherapeutics from these plants as well as from traditionally used herbal remedies [1].

The medicinal properties of plants could be based on the antioxidant, antimicrobial, antipyretic effects of the phytochemicals in them [3]. According to World Health Organization, medicinal plants would be the best source to obtain a variety of drugs. Therefore, such plants should be investigated to better understand their properties, safety and efficacy [4].

Medicinal plants produce bioactive compounds used mainly for medicinal purposes. These compounds either act on different systems of animals including man, and/or act through interfering in the metabolism of microbes infecting them. The microbes may be pathogenic or symbiotic. In either way, the bioactive compounds from medicinal plants play a determining role in regulating host-microbe interaction in favour of the host. So the identification of bioactive compound in plants, their isolation, purification and characterization of active ingredients in crude extracts by various analytical methods is important. The instant rising demand of plant-

based drugs is unfortunately creating heavy pressure on some selected high-value medicinal plant populations in the wild due to over-harvesting. Several of these medicinal plant species have slow growth rates, low population densities, and narrow geographic ranges [5]; therefore they are more prone to extinction. Conversely, because information on the use of some plant species for therapeutic purpose has been passed from one generation to the next through oral tradition, this knowledge of therapeutic plants has started to decline and become obsolete through the lack of recognition by younger generations as a result of a shift in attitude and ongoing socioeconomic changes [5].

Phytochemical screening, evaluation of antimicrobial properties and antidiarrheal status of medicinal plants are today recognized as the most viable methods of identifying new medicinal plants or refocusing on those earlier reported for bioactive constituents [6]. Plants which are observed to be efficacious and frequently prescribed may contain compounds that are potential drug candidates and could rightly be recommended for further examination [7].

The growing resistance of pathogenic organisms against the antibiotics formerly recognized for their efficiency is today a real problem of public health [4]. In human pathology, microorganisms are responsible for many infections including respiratory tract diseases (pneumonia, bronchitis), skin, wound and mucous infections, sinusitis, endocarditis, osteomyelitis, syphilis, gonorrhea, tuberculosis, food poisoning and carbuncles to mention a few. They are also the germs frequently met during surgical wound infections which are often provoked by the use of intravascular catheters or by the spread of bacteria from another source of infection. *Staphylococcus aureus* for instance one of the most pathogenic among the species of *Staphylococcus* is responsible for almost 25 % of septicemias met in hospitals [8]. Generally, the treatment of infections caused by microorganisms is long and expensive. The antimicrobial properties of plants have been

investigated by a number of researchers worldwide though thorough biological evaluation of plants extracts is vital to ensure their efficacy and safety. These factors are of importance if plant extracts are to be accepted as valid medical agents for the treatment of infectious diseases especially in the light of the emergence of drug resistance microorganism [4].

Diarrhea is characterized by increased stool frequency and a change in stool consistency is one of the major health threats to populations in the tropical and subtropical poor countries [9]. The World Health Organization (WHO) 2013 estimated that 3-5 billion cases occur each year and that approximately 5 million deaths are due to diarrhea annually.

Diarrhea is a common symptom of gastrointestinal infections caused by a wide range of pathogens, including bacteria (*Escherichia coli*, *Shigella*, *Campylobacter*, *Vibrio cholerae*), viruses (rotavirus), and protozoa (*Cryptosporidium*). However, a handful of other organisms are also responsible for most acute cases of childhood diarrhea. Among these, rotavirus is the leading cause of acute diarrhea, and responsible for about 40% of all hospital admissions due to diarrhea among under five children worldwide. Globally, rotavirus is the most common cause of severe diarrhea in children [10].

Evidence from several experimental studies showed that plant material with antimicrobial activity also possesses significant antidiarrheal activity particularly in infectious diarrhea. In both cases, these activities have been attributed to the presence of bioactive agents such as tannins, alkaloids, saponins, flavonoids, steroids, and terpenoids [11-13].

Ziziphus mauritiana belongs to the family *Rhamnaceae*. It is widely grown in mild-temperate region and is adapted to warm climates. *Ziziphus mauritiana* can grow either as shrublets, shrubs or trees with thorny branches and are used as a hedge to form defensive fences for animals [14].

The plant is known with many local names including jujube, Chinese date, Indian plump [15]. It is called “Magarya” in Hausa or “Huhue” in bura among the northern people of Nigeria.

The plant finds various uses in traditional medicine for instance; the pulp is applied on cuts and ulcers; are employed in pulmonary ailments and fevers; the dried ripe pulp is a mild laxative.

The seeds are sedative and are taken sometimes with butter, to halt nausea, vomiting and abdominal pains in pregnancy. Mixed with oil, they are rubbed on rheumatic areas. The leaves are helpful in liver trouble, asthma and fever. The bitter, astringent bark decoction is taken to halt diarrhoea and dysentery and relieve gingivitis. A root decoction is given as a febrifuge, taenicide and emmenagogue, and the powdered root is dusted on wounds. Juice of the root bark is said to alleviate gout and rheumatism [15]. The root is also used in the treatment of epilepsy [16]. The dried root is also used to treat diarrhoea in Northern Parts of Nigeria (Personal communication). The leaves are applied as poultices and are helpful in liver troubles, asthma and fever [17]. The hepatoprotective activity of ethanol extract of *Ziziphus mauritiana* leaf against CCl₄ - induced liver damage in rats and the antidiarrheal activity of the methanol root extract were reported [18,19]. The antioxidant activity of the aqueous extract of *Ziziphus mauritiana* leaf has also been reported [20].

2.0 Materials and Method

2.1 Experimental Animals

A total of twenty (20) albino rats of either sex with 110-150g body weight were used for this study. The animals were obtained from the animal house of National Veterinary Research Institute, Vom, Plateau State, Nigeria. The rats were housed in clean and disinfected plastic cages and were allowed to acclimatize for one week in the laboratory unit of Biochemistry

Department, Ahmadu Bello University Zaria before the start of the experiment. The rats were fed with a standard rat chow and allowed to drink water *ad libitum*. All animals were handled properly and carefully to minimize the effects of experimental stress. All experiments were carried out in accordance with the guidelines of the Institutional Animal Ethics Committee.

2.2 Collection of Plant Material

Pulp and seeds of *Ziziphus mauritiana* were collected from Hyera village of Hawul local Government Borno State, Nigeria, identified and authenticated at the herbarium of department of Biological Sciences, Ahmadu Bello University with the voucher number 295.

2.3 Preparation of Plant Extracts.

The dry pulp and seed of *Z. mauritiana* were pulverized using pestle and mortar and sieved. About 200g of the coarse powdered fruit and seed were extracted with 1000 ml of methanol by cooled maceration. Methanol was evaporated in water bath at less than 40°C. Distilled water was used to reconstitute the solid extract to obtain a desired concentration for the studies.

2.4 Preliminary Phytochemical Screening

Preliminary phytochemical screening for the presence of phenols, tannins, flavonoids, alkaloids, terpenoids, anthraquinones, steroids and saponins were carried out using standard test protocols [21].

2.5 Castor Oil Induced Diarrhea

The effect of the pulp and seed extractsof *Z. mauritiana*on castor oil induced diarrhea was evaluated according to the method of Awouter *et al* [22]. Rats were weighed and grouped into 4 groups (n=5). Group 1 receive distilled water, group 2 and 3 were administered 50 and 100 mg/kg extract orally while group 4 received loperamide (2mg/kg) orally. Each animal was then given 0.5 ml of castor oil orally after 30 minutes of treatment and placed in transparent cages.

The consistency of faecal matter and frequency of defecation by the animals were recorded during an observation interval of 4 hours.

2.6 Castor Oil Induced Fluid Accumulation

Rats were weighed and grouped into 4 groups (n=5). Group 1 receive distilled water, group 2 and 3 were administered 50 and 100 mg/kg extract orally while group 4 received loperamide (2mg/kg) orally. 30 minutes later, each rat was administered 2ml castor oil. The rats were anaesthetized 30 minutes later by inhalation with chloroform. The small intestine from the pylorus to caecum was dissected out and its content expelled into a measuring cylinder and the volume of the fluid was measured [23].

2.7 Small Intestinal Propulsion

The effect of *Z. mauritiana* pulp and seeds extracts on Intestinal Propulsion in rats was tested using the charcoal meal method [24]. The rats were fasted for 24 hours but allowed free access to water. Rats were weighed and grouped into 4 groups (n=5). Group 1 receive distilled water, group 2 and 3 were administered 50 and 100 mg/kg extract orally while group 4 received loperamide (2mg/kg) orally. After 30 minutes each rat was administered 1ml of 5% activated charcoal suspended in 10% aqueous tragacanth orally. The rats were sacrificed 30 min later by inhalation with chloroform. The small intestine of each animal was carefully inspected and the distance traversed by the charcoal meal from the pylorus was measured. The length of the whole small intestine was also measured. The distance traversed by the charcoal meal from the pylorus was expressed as a percentage of the distance from the pylorus to the ileocaecal junction.

3.0 Statistical Analysis

The results of the experiment were expressed as means \pm SD. The data were analyzed using one-way ANOVA followed by Duncan Multiple's Range test. Results with $p < 0.05$ were considered significant.

4.0 Results

Result of phytochemical screening (Table 1.0) revealed the presence of tannins, flavonoids, saponins, terpenoids, and cyanogenic glycosides in both the pulp and seed. Phenol was only detected in the seed. Alkaloids and anthraquinones were not detected at all in both seed and pulp. In the castor oil-induced diarrheal model for pulp (Table 2.0), the pulp extracts of *Ziziphus mauritiana* significantly prolonged the time of diarrheal induction and the frequency of stooling. Data from the experiment revealed that the percentage of diarrheal inhibition compared to controls was 63.64%, and 61.36% at the doses of 50 and 100 respectively. The drug showed a higher percentage inhibition (80.64%) compared to the extract as well as the control. In intestinal fluid accumulation test for pulp (Table 3.0), the pulp extract reduced the volume of intestinal fluid. Maximum percentage inhibition of the volume of intestinal contents was observed at 50 mg/kg of the extract, being 27.71% followed by 26.51% at dose of 100mg/kg. However, the differences observed were not statistically significant. The standard drug produced a better result compared to the test doses. The pulp extract significantly inhibited gastrointestinal transit time of charcoal meal at 50 (8.67%) and 100 (20.33%) mg/kg as compared to the control (Table 4.0).

Table 1.0: Result of Preliminary Qualitative Phytochemical Screening

Phytochemical	Seed	pulp
Phenolics	+	-
Tannins	+	+
Flavonoids	+	+
Alkaloids	-	-
Saponin	+	+
Steroids	-	-
Cyanogenic glycoside	+	+
Anthraquinones	-	-
Terpenoids	+	+

+= Present - = Absent

Table 2.0: Effect of *Ziziphus mauritiana* pulp extract on castor oil induced diarrhoea in rats

Treatment	Dose	No. of Watery Diarrhoea	% Inhibition
Control	-	11.00±3.46 ^a	-
Extract	50 mg/kg	4.00±1.15 ^b	63.64
Extract	100 mg/kg	4.25±1.50 ^b	61.36
Loperamide	2 mg/kg	2.13±1.10 ^c	80.64

Values are mean±SD (n=5). Results with different superscript differ significantly at p<0.05 (One Way ANOVA followed by Duncan Multiple Range Test).

Table 3.0: Effect of methanolic pulp extract of *Ziziphus mauritiana* on castor oil induced intestinal fluid accumulation

Treatment	Dose	Fluid Volume (ml)	% Inhibition
Control	-	3.32±0.23 ^a	-
Extract	50 mg/kg	2.40±0.42 ^b	27.71
Extract	100 mg/kg	2.44±0.46 ^b	26.51
Loperamide	2 mg/kg	1.40±0.12 ^c	57.83

Values are mean±SD (n=5). Results with different superscript differ significantly at p<0.05 (One Way ANOVA followed by Duncan Multiple Range Test).

Table 4.0: Effect of methanolic pulp extract of *Ziziphus mauritiana* on gastrointestinal transit in rats.

Treatment	Dose	Length of Intestine (cm)	Distance Travelled by Charcoal meal (cm)	% Inhibition
Control	-	79.28±8.71	60.00±6.67 ^a	-
Extract	50 mg/kg	82.10±5.66	54.80±7.89 ^b	8.67
Extract	100 mg/kg	83.40±4.98	47.80±9.07 ^b	20.33
Loperamide	2 mg/kg	84.7±4.41	33.22±3.23 ^c	44.63

Values are mean±SD (n=5). Results with different superscript differ significantly at p<0.05 (One Way ANOVA followed by Duncan Multiple Range Test).

Results for the study of antidiarrheal activity of the seed are shown in (Table 5-7). The seed extracts of *Ziziphus mauritiana* significantly prolonged the time of diarrheal induction and the frequency of stooling in the castor oil-induced diarrheal model (Table 5.0). Our findings revealed that 100mg/kg of the seed extract had the maximum percentage diarrheal induction (80.70%) inhibition. The synthetic drug used produced a percentage inhibition of 91.23%. In intestinal fluid accumulation test (**Table 6.0**), there were no significant differences in percentage reduction of the volume of intestinal fluid of the seed extract. Table 7.0 shows the result of gastrointestinal transit time of charcoal meal. The seed extract significantly inhibited gastrointestinal transit time of charcoal meal by 7.63%, 19.87% and 42.27% at 50mg/kg of extract, 100mg/kg of extract and 2mg/kg of the standard drug respectively.

Table 5.0: Effect of Ziziphusmauritiana Seed extract on castor oil induced diarrhoea in rats

Treatment	Dose	No. of Watery Diarrhoea	% Inhibition
Control	-	11.40±2.07 ^a	-
Extract	50 mg/kg	5.40±1.34 ^b	52.63
Extract	100 mg/kg	2.20±1.10 ^b	80.70
Loperamide	2 mg/kg	1.00±0.04 ^c	91.23

Values are mean±SD (n=5). Results with different superscript differ significantly at p<0.05 (One Way ANOVA followed by Duncan Multiple Range Test).

Table 6.0: Effect of methanolic Seed extract of *Ziziphus mauritiana* on castor oil induced intestinal fluid accumulation

Treatment	Dose	Fluid Volume (ml)	% Inhibition
Control	-	3.44±0.26 ^a	-
Extract	50 mg/kg	2.18±0.15 ^b	36.63
Extract	100 mg/kg	2.08±0.13 ^b	39.53
Loperamide	2 mg/kg	1.52±0.36 ^c	55.81

Values are mean±SD (n=5). Results with different superscript differ significantly at p<0.05 (One Way ANOVA followed by Duncan Multiple Range Test).

Table 7.0: Effect of methanolic Seed extract of *Ziziphus mauritiana* on gastrointestinal transit in rats.

Treatment	Dose	Length of Intestine (cm)	Distance Travelled by Charcoal meal (cm)	% Inhibition
Control	-	93.4±4.72 ^a	63.88±12.87 ^a	-
Extract	50 mg/kg	101.180±4.19 ^b	59.00±9.67 ^b	7.63
Extract	100 mg/kg	103.94±6.07 ^b	51.19±9.07 ^b	19.87
Loperamide	2 mg/kg	96.34±9.11 ^c	36.88±3.21 ^c	42.27

Values are mean±SD (n=5). Results with different superscript differ significantly at p<0.05 (One Way ANOVA followed by Duncan Multiple Range Test)

5.0 Discussion and Conclusion

The use of castor oil to induce diarrhea has been reported in several studies [25,26]. The diarrhea inducing properties is as a result of ricinoleic acid; a castor oil active metabolite, which is liberated by the action of lipases in the upper part of the small intestine [27]. Ricinoleic acid exerts its effect by production of local irritation and inflammation of the intestinal mucosa, causing the release of prostaglandins that eventually increases gastrointestinal motility and net secretion of water and electrolytes [28]. Tunaruet *al.*, [29], opined that this effect could also occur due to the capability of ricinoleic acid to activate the G protein-coupled prostanoid receptor (EP3) on the smooth muscle cell of the intestine. In addition, it forms ricinoleate salts with sodium and potassium in the lumen of the intestine and these salts antagonize sodium-potassium ATPase and increase permeability of the intestinal epithelium, which in turn results in cytotoxic effect on intestinal absorptive cells [27].

From the result of castor oil induced diarrheal model, the seed and pulp extracts at all the tested doses significantly decreased the frequency of defecation. This study is in line with other studies in which the extract of different plants reduced the frequency of stooling [30-32]. Terpenoids such as abietic acid and steroids like phytosterols have been shown to inhibit production of prostaglandin E2 [33,34], which are known to play a crucial role in the stimulation of intestinal secretions [35]. Thus, the antidiarrheal effect of the seed and fruit could be attributed to inhibition of castor oil-induced prostaglandin synthesis. The anti-diarrheal activity might also be due to inhibition of active secretion of ricinoleic acid, resulting in the activation of Na⁺, K⁺ATPase activity that promotes absorption of Na⁺ and K⁺ in the intestinal mucosa. This effect could probably be linked to the presence of terpenoids, tannins and flavonoids in the seed and fruit extract, which are shown to promote colonic absorption of water and electrolytes [36].

342 In the castor oil induced intestinal fluid accumulation test, treatment of rats with graded doses of
343 the seed and fruit extract produced a significant reduction in the intestinal fluid accumulation.
344 Mascolo *et al.* (1994) reported that ricinoleic acid the active metabolite of castor oil might
345 activate the nitric oxide pathway and induce nitric oxide (NO) dependent gut secretion. A
346 number of studies have revealed that terpenoids (Jang *et al.*, 2004) and flavonoids (Kim *et al.*,
347 2004; Messaoudene *et al.*, 2011) are implicated in attenuation of NO synthesis. Hence, the
348 pronounced inhibition of castor oil induced intestinal fluid accumulation might possibly be due
349 to the presence of flavonoids and terpenoids that increase the reabsorption of electrolytes and
350 water by hindering castor oil mediated NO synthesis. The fact that intestinal fluid accumulation
351 and Na⁺ secretion induced by castor oil is attenuated by pretreatment of rats with NO synthesis
352 inhibitors [37] reinforces the notion that the anti-fluid accumulation effect of both the seed and
353 fruit extract could probably be by interfering with the NO pathway. Alkaloids which are detected
354 in the seed and fruit have also been demonstrated to inhibit NO synthesis [38].

355 Flavonoids exert their antidiarrheal effect by inhibiting intestinal motility and hydro-electrolytic
356 secretion [39,40]. Flavonoids are also able to inhibit the intestinal secretory response induced by
357 prostaglandins E2 [41]. Moreover, the enteric nervous system stimulates intestinal secretion
358 through neurotransmitters such as acetylcholine and vasoactive intestinal peptide. On the other
359 hand, intestinal absorption can be stimulated with alpha two adrenergic agents, enkephalins, and
360 somatostatin [35]. Secondary metabolites such as flavonoids from plant sources could stimulate
361 alpha two adrenergic receptors in the absorptive cells of the gastrointestinal tract [40]. Hence, the
362 significant anti-secretory activity of the seed and fruit extract could probably be related to the
363 presence of flavonoids that in turn stimulate alpha two adrenergic receptors in the enterocytes
364 and facilitate fluid and electrolyte absorption.

Results of evaluation of gastrointestinal transit in rats demonstrated that the seed and fruit significantly reduced the intestinal propulsive movement of charcoal meal at all the test doses as compared to the negative control. The findings are in line with other studies in which the plant extracts significantly inhibited the distance travelled by charcoal meal [42]. This observation could be ascribed to the synergistic effects of terpenoids and alkaloids present in the seed and fruit to prolong the time for absorption of water and electrolytes through hampering peristaltic movement of the intestine. Indeed, alkaloids and terpenoids have been demonstrated to have inhibitory effect on gastrointestinal motility [36,43]. Although the phytochemical constituents responsible for the antidiarrheal effect are yet to be identified, the amount of phytochemical constituents that are responsible for impeding gastrointestinal motility such as tannins [44-46] and alkaloids appear to increase with dose [36]. This could possibly be the reason why significant anti-motility effect was observed.

Plants that have tannins in their composition can precipitate proteins of the enterocytes, reducing the peristaltic movements and intestinal secretions [44-46]. The layers formed by the precipitate of proteins on the mucosal surface of the enterocytes also inhibit the development of microorganisms, thus explaining the antiseptic action of tannins [44].

The pulp and seed extracts of *Ziziphus mauritiana* have anti-diarrheal activity against castor oil induced diarrhea. The anti-diarrheal activity is thought to be due to the presence of phytochemicals in the plant.

Ethical Disclaimer:

As per international standard or university standard written ethical permission has been collected and preserved by the authors.

References

- [1] UNESCO, (1998). FIT/504-RAF-48 Terminal Report:Promotion of Ethnobotany and the Sustainable Use of PlantResources in Africa, pgs. 60, Paris, 1998
- [2] Kala, C.P.(2005). Ethnomedicinal botany of the Apatani in the Eastern Himalayan region of Indian.*Journal of Ethnobiology and Ethnomedicine*; **1**:1-8.
- [3] Ayodele, A. E. (2005). The Medicinally Important Leafy Vegetables of Southwestern Nigeria.*Ethnobotanical leaflets*, **1**: 16.
- [4] Nascimento, G. G., Locatelli, J., Freitas, P. C. and Silva, G. L. (2000).Antibacterial activity of plant extracts and phytochemicals on antibiotic-resistant bacteria.*Brazilian journal of microbiology*, *31*(4), 247-256.
- [5] Kala, C. P., Dhyani, P. P., and Sajwan, B. S. (2006). Developing the medicinal plants sector in northern India: challenges and opportunities.*Journal of Ethnobiology and Ethnomedicine*, **2**(1): 1.
- [6] Adjanahoun, E., Ahyi, M. R. A., Ake-Assi, L., Elewude, J. A., Dramane, K., Fadoju, S.O., Gbile, Z. O., Goudole, E.,Johnson, C. L .A., Keita, A., Morakinyo, O. , Ojewole, J. A. O. , Olatunji, A. O. and Sofowora, E. A. (2001). *Traditional medicine and pharmacopoeia. Contribution to ethnobotanical floristic studies in Western Nigeria*.Organization of African Unity Scientific Technical and Research Commission, Lagos, Nigeria, pp 420.
- [7] Tor-Anyiin, T. A., Sha'ato, R., and Oluma, H. O. A. (2003).Ethnobotanical survey of anti-malarial medicinal plants amongst the Tiv people of Nigeria.*Journal of herbs, spices & medicinal plants*, **10**(3): 61-74.

- 414 [8] Moyen, G., Nkoua, J. L., Mbemba, A. B., Fourcade-Pauty, V. and Nzingoula, S.
415 (2003).Septicémie à *Staphylococcus aureus* de l'enfant- à propos de 12 cas. Médecine
416 d'Afrique noire, **40** (6).
- 417 [9] Farthing, M. J. G. (2002). Novel targets for the control of secretory diarrhea. *Gut***50**: 15–
418 18.
- 419 [10] Centers for Disease Control and Prevention (CDC).(2008). Rotavirus surveillance--
420 worldwide, 2001-2008.*MMWR.Morbidity and mortality weekly report*, **57**(46), 1255.
- 421 [11] Longanga-Otshudi, A., Vercruysse A and Foriers A;(2000). Contribution to the
422 Ethnobotanical, Phytochemical and Pharmacological Studies of Traditionally used
423 Medicinal Plants in the Treatment of Dysentery and Diarrhea in Lomela area, Democratic
424 Republic of Congo (DRC), *Journal of Ethnopharmacology*, **73**(3): 411-423.
- 425 [12] Umar, Y. A., Maikaje, D. B., Garba, U. M. and Alhassan, M. A. F (2013).Prevalence of
426 gastro-intestinal parasites in horses used for Cadets training in Nigeria.*Journal of*
427 *Veterinary Advances*, **3**(2): 43-48.
- 428 [13] Getnet, M. (2015).*Evaluation of the anti-diarrheal activity of 80% methanol extract and*
429 *solvent fractions of the leaves of Lantana camara Linn (Verbenaceae) in mice* (Doctoral
430 dissertation, AAU).
- 431 [14] Cherry, M. (1985). The needs of the people. In Wickens, G. E., Goodin, J.R., Field
432 [15] Morton, J. (1987). Indian Jujube. In: Pulp of Warm Climates. Morton JF (ed),
433 Miami,Florida. 272-275. Last updated: 23/4/2004.
- 434 [16] Msonthi, J. D., andMagombo, D. (1983).Medicinal herbs in Malawi and their
435 uses.*Hamdard*, **26**(2), 94-100.
- 436 [17] Michel, A. (2002). *Tree, Shrub and Liana of West African Zone*.MargrafPublishers
437 GMBH, Paris, p. 440.
- 438 [18] Dahiru, D., William, E. T. and Nadro, M. S. (2005). Protective effect of Ziziphus
439 mauritiana leaf extract on carbon tetrachloride-induced liver injury.*African Journal of*
440 *Biotechnology***4**(10): 1177-1179.
- 441 [19] Dahiru, D., Sini, J. M., and John-Africa, L. (2006).Antidiarrhoeal activity of
442 Ziziphusmauritiana root extract in rodents. *African Journal of Biotechnology*, **5**(10).
443
- 444 [20] Dahiru, D. and Obida, O. (2008). Evaluation of the antioxidant effects of
445 Zizyphusmauritiana lam. Leaf extracts against chronic ethanolinduced hepatotoxicity in
446 rat liver.*African Journal of Traditional and complimentary medicine*.**5**(1): 39-45.
- 447 [21] Trease, G.E. and, Evans W.C. (1996): *Pharmacognosy*.BailliereTindall, London; 89-
448 122.

- 449 [22] Awouters, F., Niemegeers, C. J. E., Lenaerts, F. M., and Janssen, P. A. J. (1978). Delay
450 of castor oil diarrhoea in rats: a new way to evaluate inhibitors of prostaglandin
451 biosynthesis. *Journal of Pharmacy and Pharmacology*, **30**(1): 41-45.
- 452 [23] Adzu, B., Amos, S., Amizan, M. B., and Gamaniel, K. (2003). Evaluation of the
453 antidiarrhoeal effects of *Zizyphus pinnatifida* stem bark in rats. *Actatropica*, **87**(2): 245-
454 250.
- 455 [24] Capasso, F., Pinto, A., Mascolo, N., Autore, G., and Franco, M.P. (1988). Effects of
456 flavonoids on PGE₂ and LTD induced contractions on guinea pig isolated ileum,
457 *Pharmacological Research Communications*, **20**: 201-202
- 458 [25] Rahman, A., Hasan, S. N., Sampad, K. S. and Das, A. K. (2011). Antinociceptive, anti
459 diarrhoeal and cytotoxic activities of *Rhizophoramide* Lamk. *Pharmacologyonline* **1**:
460 921-929.
- 461 [26] Akter S, Sarker A, Hossain S (2013). Antidiarrhoeal activity of rind of
462 *Punica granatum*. *International current Pharmaceutical journal* **2**(5): 101-104.
- 463 [27] Komal, Kumar S, Rana AC (2013). Herbal approaches for diarrhea: A review.
464 *International Research Journal of Pharmacy* **4**(1):31-38.
- 465 [28] Robert, A., Nezamis, J. E., Lancaster, C., Hanchar, A. J. and Klepper, M. S. (1976).
466 Enteropooling assay: a test for diarrhea produced by prostaglandins. *Prostaglandins*,
467 **11**(5), 809-828.
- 468 [29] Tunaru, S., Althoff, T. F., Nusing, R. M., Diener, M. and Offermanns S (2012). Castor oil
469 induces laxation and uterus contraction via ricinoleic acid activating prostaglandin EP₃
470 receptors. *Proceedings of National Academy of Science, USA* **109**(23):9179-9184.
- 471 [30] Mazumder, R., Bhattacharya, S., Mazumder, A., Pattnaik, A. K., Tiwary, P. M.
472 and Chaudhary S (2006). Antidiarrhoeal evaluation of *Aegle marmelos* (Correa) Linn. root
473 extract. *Phytotherapy Research*, **20**(1): 82-84.
- 474 [31] Karthik, P., Kumar, R. N. and Amudha, P. (2011). Antidiarrheal activity of the chloroform
475 extract of *Cayratia pedata* Lam in albino wistar rats. *Pharmacologyonline* **2**: 69-75.
- 476 [32] Billah, M.M., Khatun, H., Parvin, S., Islam, E., Islam, S.M., and Mia, A.A. *et al* (2013).
477 Antibacterial, antidiarrhoeal, and cytotoxic activities of methanol extract and its fractions
478 of *Caesalpinia bonducella* (L.) Roxb leaves. *BMC Complementary Alternative*
479 *Medicine*, **13**(1):101.
- 480 [33] Awad, A.B., Toczek, J. and Fink, C.S. (2004). Phytosterols decrease prostaglandin release
481 in cultured P388D1/MAB macrophages. *Prostaglandins Leukot Essent Fatty Acids*
482 **70**(6):511- 520.

- 483 [34] Fernandez MA, Tornos MP, Garcia MD, De lasHeras B, Villar AM, Saenz MT (2001).
 484 Antiinflammatory activity of abietic acid, a diterpene isolated from *Pimentaracemosa*
 485 *var. grissea*. *Journal of Pharmacy and Pharmacology***53**(6):867-872.
- 486 [35] Bern, M.J., Sturbaum, C.W., Karayalcin, S.S., Berschneider, H.M., Wachsman, J.T. and
 487 Powell, D. (1989). Immune system control of rat and rabbit colonic electrolyte transport.
 488 Role of prostaglandins and enteric nervous system. *Journal of Clinical*
 489 *Investigation***83**(6):1810-1820.
- 490 [36] Palombo, E.A and Semple, S.J. (2001). Anti-bacterial activity of traditional medicinal
 491 plants. *Journal of Ethnopharmacology*, **77**: 151-157.
- 492 [37] Mascolo N, Izzo AA, Barbato F, Capasso F (1993). Inhibitors of nitric oxide synthetase
 493 prevent castor-oil-induced diarrhoea in the rat. *British Journal of*
 494 *Pharmacology*.**108**(4):861-864.
- 495 [38] Kondo Y, Takano F, Hojo H (1993). Inhibitory effect of bisbenzylisoquinoline alkaloids
 496 on nitric oxide production in activated macrophages. *Biochemical*
 497 *Pharmacology***46**(11):1887- 1892.
- 498 [39] Rao, V. S., Santos, F. A., Sobreira, T. T., Souza, M. F., Melo, C. L. and Silveira, E. R.
 499 (1996). Investigations on the gastroprotective and antidiarrhoeal properties of ternatin, a
 500 tetramethoxyflavone from *Egletes viscosa*. *Planta Medica***63**(2):146-149.
- 501 [40] DiCarlo, G., Autore, G. and Izzo, A. (1993). Inhibition of intestinal motility and the
 502 secretion by flavonoids in mice and in rats: structural activity relationship, *Journal of*
 503 *Pharmacy and Pharmacology*, **45**: 1054–1059.
- 504 [41] Hamalainen M, Nieminen R, Asmawi MZ, Vuorela P, Vapaatalo H, Moilanen E (2011).
 505 Effects of flavonoids on prostaglandin E2 production and on COX-2 and mPGES-1
 506 expressions in activated macrophages. *Planta Medica***77**(13):1504-1511.
- 507 [42] Zavala-Mendoza, D., Alarcon-Aguilar FJ, Perez-Gutierrez S, Escobar-Villanueva MC,
 508 Zavala Sanchez MA (2013). Composition and Antidiarrheal Activity of *Bidens odorata*
 509 Cav. *Evidence Based Complement Alternative Medicine* **1**:1-7.
- 510 [43] Maciel MA, Pinto AC, Arruda AC, Pamplona SG, Vanderlinde FA, Lapa AJ, *et al*
 511 (2000). Ethnopharmacology, phytochemistry and pharmacology: a successful
 512 combination in the study of *Croton cajucara*. *Journal Ethnopharmacology***70**(1):41-55.
- 513 [44] Almeida CE, Karnikowski MG, Foletto R, Baldisserotto B (1995). Analysis of
 514 antidiarrhoeic effect of plants used in popular medicine. *Revista*
 515 *de Saude Publica***29**(6):428-33.
- 516 [45] Tripathi, K. (2008). *Essentials of Medical Pharmacology*, 8 ed. New Delhi: Jaypee
 517 Brothers Medicals Publishers (P) Ltd, New Delhi.

- 518 [46] Yadav, A. K. andTangpu, V. (2007). Antidiarrheal Activity of Lithocarpusdealbata and
519 Urenalobata Extracts: Therapeutic Implications. *Pharmaceutical Biology*,**45**(3): 223–229.

520

UNDER PEER REVIEW