

Case study

METASTATIC SIGMOID COLON CANCER PRESENTED AS INCARCERATED INGUINAL HERNIA – CASE REPORT

ABSTRACT

Inguinal hernia containing metastases of intestinal adenocarcinoma is a rare finding. Metastases of sigmoid colon adenocarcinoma are most commonly found inside the hernia sac. Older males are more often affected. A 84-years old male patient presented with pain in the right groin, highly suspicious to be an incarcerated right inguinal hernia. During emergency operation we found mesenteric metastases of the small intestine, that was incarcerated inside the inguinal hernia sac. Histopathological result of biopsy has shown metastasis of intestinal type adenocarcinoma. Additional diagnostics has shown sigmoid colon adenocarcinoma, peritoneal carcinosis and liver metastases. Primary tumor was locally advanced and unresectable, groin hernia was repaired using Bassini technique and diverting colostomy was performed. Due to advanced metastatic disease, symptomatic treatment was advised.

In patients presenting with groin hernia, when suspected malignant lesion is found within hernia sac, histopathologic verification of the lesion is needed. Further diagnostics is also indicated for the definition of the primary tumor.

KEYWORDS: Adenocarcinoma, Carcinoma, Sigmoid Colon , Groin, Inguinal hernia.

INTRODUCTION

Malignant lesion inside the inguinal hernia sac is a rare finding [1]. Metastases of colorectal cancer are localized within the inguinal hernia sac in less than 1 in 200 patients and they are usually asymptomatic [2]. Data suggest that older male patients are most often affected. The most common origin is from the sigmoid colon and metastases are usually found in the left groin hernia sac [1, 2]. The first case of a tumor inside the inguinal hernia sac was reported in 1749 [3].

The following report presents a case of an older male patient with a right inguinal hernia containing metastases of a well differentiated adenocarcinoma of the intestinal type. The origin was an adenocarcinoma of the sigmoid colon.

CASE REPORT

84-years old male patient was referred to the emergency department (ED) with pain and bulge in the right groin, suspected to be an incarcerated right inguinal hernia. He had pain in the right groin for a longer time, which has become worse two days prior to the examination at the ED. He reported nausea, without vomiting. In the past, he has been conservatively treated at our department due to acute cholecystitis. Patient had some difficulties with defecation for a few months and he lost some weight.

Irreducible right inguinoscrotal hernia was found on clinical examination. The lower abdominal quadrants were painful on palpation. Abdominal x-ray has shown signs of small intestinal obstruction (Figure 1). Abdominal ultrasound (US) was also performed, which showed aperistaltic, edematous and poorly vascularized segment of small intestine. Bilateral hydroceles were also described. The right testicle was hyperemic, without focal lesions and the left testicle was normal.

58 Patient was operated on the day of admission to the hospital. Right parainguinal
59 skin incision was performed. Inside the hernia sac, small intestine was found. The
60 intestine was vital, without signs of obstruction. In the small intestinal mesentery a
61 large macroscopically malignant lesion was found (Figure 2). We took a sample for
62 histopathological examination. At that time we did not decide for further abdominal
63 exploration, because we wanted to perform further diagnostics and to wait for
64 histopathological results. We performed only a hernioplasty of the right inguinal
65 hernia according to Bassini technique.

66 During hospitalization, after the first operation, further diagnostics was performed
67 and we acquired the result of histopathological examination, which has shown a well
68 differentiated adenocarcinoma of the intestinal type - mesenteric carcinosis.
69 Colonoscopy showed an obstructive tumor at 15 cm proximally from anal verge.
70 Biopsy was taken and sent for histopathologic verification, the result was colonic
71 adenocarcinoma. Irigography showed colonic obstruction 13 cm proximally from anal
72 verge (Figure 3). Abdominal computed tomography (CT) has shown a
73 heterogenous tumor formation of the distal sigmoid colon, measuring 12x7.5 cm.
74 Enlarged intraabdominal lymph nodes were described, up to 1.2 cm in diameter.
75 Colon was elongated, filled with intestinal content, without signs of intestinal
76 obstruction. Liver metastases were also seen on abdominal CT (Figure 4).

77 According to the results of extensive diagnostics, we were aware, that the primary
78 tumor was a sigmoid colon adenocarcinoma with peritoneal carcinosis and liver
79 metastases. The primary tumor was causing a bowel obstruction, so we decided for
80 an exploratory laparotomy. Patient was once again operated 21 days after the first
81 operation. We performed a median laparotomy and a thorough abdominal
82 exploration. During abdominal exploration, we found a large tumor in the upper third

83 of the rectum and distal sigmoid colon, also peritoneal carcinosis and liver
84 metastases were found. Primary tumor was large and fixed in to the retroperitoneum.
85 According to the advanced metastatic disease, we decided to perform a diverting
86 colostomy to avoid further colonic obstruction. A loop bipolar sigmoidostomy was
87 performed.

88 Further recovery was uneventful and patient was discharged from hospital 10 days
89 after the second operation.

90 Patient was presented to the multidisciplinary oncological team - symptomatic
91 treatment was advised.

92

93 **DISCUSSION**

94

95 Malignant neoplasm is found inside the hernia sac in less than 0.5 % of all inguinal

96 hernias [1]. Hernia sac tumors are classified into three groups regarding the
97 relationship of the tumor to the hernia sac [3]. Intrascacular tumors include primary
98 tumors of organs, that are incarcerated in the hernia (e.g. bladder cancer, colon
99 cancer, appendix cancer, metastatic neoplasms involving omentum) [1, 3, 4].

100 Saccular tumors are primary or secondary malignant lesions, that involve peritoneum
101 (e.g. primary mesothelioma, peritoneal metastases from prostate, ovary, colon,
102 pancreas) [1, 3, 5]. Extrasaccular malignant lesion is any tumor protruding through
103 the hernia defect, but outside the hernia sac (e.g. metastatic inguinal lymph node) [3,
104 6, 7]. Carcinomas are the most frequent tumors found in a hernia sac.

105 However, these malignant epithelial tumors are rare [8]. Inguinal hernia sac
106 containing malignant lesion is usually asymptomatic [3]. There are some hypotheses,
107 that a longstanding hernia, that becomes acutely incarcerated, has a higher
108 likelihood of containing tumor [3]. Some authors believe, that any irreducible inguinal

mass, that lacks a tactile impulse, should raise suspicion of cancer [3]. Data from literature suggest, that about a fifth of all male patients with colorectal cancer have concurrent inguinal hernia or have had a repair of inguinal hernia 1-2 years prior to the diagnosis of cancer. Every malignant lesion found within hernia sac should be examined histologically [3, 4, 6]. Hernias are the most frequent structural abnormalities of the groin area. Differential diagnosis includes groin masses, as they may simulate inguinal or femoral hernias. Therefore, malignant disease should be considered as a possible diagnosis in patients, who present with an unexplained groin mass. The surgeon should be alert to the possibility of encountering malignant disease at hernia sites [8, 9, 10]. A prospective study has shown no association between inguinal hernia and colorectal cancer [4, 5]. Symptoms such as abdominal pain and weight loss should raise suspicion of cancer [3]. Our patient had some bowel symptoms prior to the first operation. He lost his appetite and had difficulties with defecation. He also lost some weight. Even though the sigmoid colon tumor was causing colonic obstruction, patient was passing a liquid stool and he did not have any major abdominal pain. This was probably because he was on a liquid diet for some time because of reduced appetite.

Routine microscopic study of all hernia sacs is inadequate for the high cost (insufficient cost-benefit ratio). This practice should be reserved for cases in which a significant lesion is clinically or macroscopically suspected. Therefore, all hernia sacs should be examined grossly. Microscopic evaluation should be done for abnormal tissue discovered at surgery or at the pathology ward, which is suggestive of an underlying disease process. The decision to submit a hernia sac for histology may be left to the discretion of the surgeon [8, 11].

There are no clear guidelines, which surgical approach is the most appropriate [1, 4]. It usually depends on local anatomy, surgical findings and surgeon's experience [1, 4, 7]. In most reported cases, colonic resection at laparotomy is followed by conventional inguinal hernia repair through separate incision [1].

In our case, we decided for a two stage operation, because at the time of the first operation, we did not have any diagnostics, except of abdominal x-ray and we wanted to wait for a histopathological result of biopsy. However, in our case, according to the primary finding of macroscopically malignant tissue within hernia sac, further exploratory laparotomy or laparoscopy would be meaningful.

The main limitation in our case was, that at the time of the first operation, there was no clinical suspicion of a possible malignant lesion inside the incarcerated hernia. We did not have any other diagnostics, except the abdominal x-ray. If there would be a clinical suspicion of a possible malignancy, we would perform more extensive diagnostics and we would decide for an exploratory laparotomy at the time of the first operation.

CONCLUSION

Inguinal hernia sac containing colon cancer metastases is a rare finding. In such patients additional diagnostics is indicated to find the origin and to evaluate the stage of the disease. Systematic histologic study of all hernia sacs is impractical because of the high cost. The decision to examine the hernia sac microscopically may be left to the discretion of the surgeon. The finding of a malignant epithelial tumor in a hernia sac usually suggests advanced disease and a short life expectancy of the patient. Surgical approach depends mostly on patient's habitus and surgeon's experience and preferences.

159

160 **CONSENT**

161 As per international standard of university standard written patient consent has been
162 collected and preserved by the author(s).

163

164 **ETHICAL APPROVAL**

165 It is not applicable.

166

167

168 **REFERENCES**

169

- 170 1. Ruiz Tovar J, Ripalda E, Beni R, Nistal J, Monroy C et al. Carcinoma of the sigmoid
171 colon in an incarcerated inguinal hernia. *Can J Surg.* 2009; 52: 2.
172
- 173 2. P. B. Salemans, G. F. Vles, S. A. F. Fransen, and R. M. Smeenk, "Sigmoid
174 Carcinoma in an Inguinal Hernia: A Blessing in Disguise?," *Case Reports in*
175 *Surgery.* 2013; Article ID 314394, 3 pages. doi:10.1155/2013/314394.
176
- 177 3. Phifer Nicholson C, Donohue IH, Thompson GB, Lewis IE. A study of metastatic
178 cancer found during inguinal hernia repair. *Cancer.* 1992; 69: 3008-3011.
179
- 180 4. Marsden M, Curtis N, McGee S, Bracey E, Branagan G et al. Intrascapular caecal
181 adenocarcinoma presenting as enlarging right inguinoscrotal hernia. *International*
182 *Journal of Surgery Case Reports.* 2014; 5: 643-645.
- 183 5. Rong Q, Qiaoyu Z, Jianfeng W, Yongdong P. Incidental finding of a malignant
184 tumour in an inguinal hernia sac. *Contemp Oncol.* 2014; 18(2): 130–133. doi:
185 10.5114/wo.2014.42728.
186


- 187 6. Chen KT. Metastatic carcinoma in inguinal hernia sac. J Surg Oncol. 1984; 25(4):
188 248-249.
- 189
190
- 191 7. Ping-Hung L, Wen-Ching K, Yu-Chiuan W, Shang-Tao C, Wen-Yen C, Chin-Wen H.
192 Metastatic malignant gastrointestinal stromal tumor mimicking a right incarcerated
193 inguinal hernia. Formosan Journal of Surgery. 2014; 47(5): 189-192.
- 194 8. Val-Bernal JF, Mayorga M, Fernández FA, Val D, Sánchez R. Malignant epithelial
195 tumors observed in hernia sacs. Hernia. 2014; 18(6): 831-835. doi: 10.1007/s10029-
196 014-1283-z.
- 197 9. Qin R, Zhang Q, Weng J, Pu Y. Incidental finding of a malignant tumour in an
198 inguinal hernia sac. Contemp Oncol. 2014; 18(2): 130-133. doi:
199 10.5114/wo.2014.42728.
- 200 10. Burke TP, Waters P, Khan W, Barry K. Bilateral saccular inguinal hernias in an
201 elderly woman presenting with advanced ovarian cancer. BMJ Case Rep. 2014; 27.
202 doi: 10.1136/bcr-2013-202337.
- 203 11. Li S, Li Y, Tang L, Zhao P. Obstructed small bowel ruptured toward the inguinal
204 canal resulting from metastatic colon carcinoma in an irreducible, recurrent inguinal
205 hernia with mesh-plug repair: Report of a case. Eur J Surg Oncol. 2010; 36(10): 1012-
206 1014. doi: 10.1016/j.ejso.2010.01.008.
- 207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224

FIGURES

Figure 1: Abdominal x-ray showing signs of small intestinal obstruction.



Figure 2: Small intestine with large metastatic lesion in the mesentery (arrow).



236
237

Figure 3: Irigography showing a colonic obstruction at 13 cm from anal verge.



238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262

263 **Figure 4:** Abdominal CT showing a large sigmoid colon tumor (arrow).
264



265