

Case Study**Intra Bony Epidermoid Cyst Mimicking Odontogenic Cyst – A Case Report****Abstract:**

Epidermoid and Dermoid cysts are benign lesions encountered throughout the body. These cysts are cystic malformations lined with squamous epithelium. Development of these cysts in the oral cavity is extremely rare. Epidermoid cysts can arise by a development of entrapped ectodermal tissue of the first and second branchial arches or can also arise due to surgical or accidental implantation of epithelial cells into deeper tissues. Clinically, the lesion presents as a slow-growing asymptomatic mass, usually located in the midline, above or below the mylohyoid muscle. Histologically, they can be further classified as epidermoid, dermoid or teratoid. Originally the implantation cysts are developed from congenital inclusion of ectodermal tissue during embryological development. But sometimes they may originate through implantation of epithelium by either surgical or accidental trauma into deeper mesenchymal tissues. This article presents a rare case of inclusion type of epidermoid cyst in the body of the mandible.

Keywords:

Cysts, Epidermoid cysts, jaw cysts, mandible, inclusion cysts.

Introduction:

Epidermoid and Dermoid cysts are benign lesions encountered throughout the body, particularly in areas where embryonic elements fuse together. Most (80%) are located in the ovaries and sacral region. Some (about 7%) can be found in head and neck region.¹ These cysts are cystic malformations lined with squamous epithelium. Development of these cysts in the oral cavity is extremely rare. They constitute 1.6% to 6.9% of all cysts in the head and neck area.² Depending on the pathogenesis; epidermoid cysts can be divided into the congenital and acquired types. The former are called epithelial and the latter are known as implantation or posttraumatic cysts.³ in this article a unique case of intra bony epidermoid cyst involving the body of the mandible is presented.

Case Report:

A 20 year old man presented with good health with a chief complaint of a swelling and facial asymmetry in the left side body of the mandible for past 6 months. The swelling was painless. His past medical and family history was noncontributory. On clinical examination the lesion presented as a firm bony hard painless swelling on the left body of the mandible. Teeth at the region of the swelling were normal. An intra oral periapical and occlusal radiographs showed a well-defined radiolucent lesion with cloudy appearance and sclerotic margin which displaced the roots of teeth in that region. A panoramic radiograph showed a well- defined radiolucent lesion with a sclerotic margin displaying the teeth in that

39 region (figure 1). A CT scan revealed the lesion to be 4.5 x 6.5 cm in size with expansion of
40 both the buccal and lingual plates (figure 2&3). When the swelling was aspirated, it revealed
41 white colored keratin material. A differential diagnosis of collateral - type odontogenic
42 keratocyst, early stage of cemento-ossifying fibroma, unicystic ameloblastoma, aneurysmal
43 bone cyst and epidermoid implantation cyst was made. The lesion was completely enucleated
44 and histopathological examination showed cystic lining and stromal wall. The lining was
45 made up of 4-5 cells thick epithelium with a well pronounced granular cell layer and thick
46 keratin layer resembling epidermis. The stroma was made up of fibrous connective tissue
47 with areas of hyalinization and focal areas of inflammation. There was no evidence of any
48 dermal appendages.(figure 4) The final diagnosis was made as implantation type of
49 epidermoid cyst.

50 **Discussion:**

51 Epidermoid cysts can arise by a development of entrapped ectodermal tissue of the
52 first and second branchial arches or can also arise due to surgical or accidental implantation
53 of epithelial cells into deeper tissues.¹ According to McCallum, the congenital variety can
54 originate as a late displacement of the ectoderm or may develop, as the teratomatous or
55 branchial cyst does, from the residual tissue separated from the branchial opening. Brosch,
56 Axhausen, Hendricks, and Ward, cited by Schuchardt, believed that these cysts are formed in
57 much the same way as the epithelial germ that remained displaced within the maxillary bone
58 during embryonic development.⁴ Sutton believed that implantation cysts originated through
59 implantation of epithelium by either surgical or accidental trauma into deeper mesenchymal
60 tissues. There is usually a latent period after injury before the cyst is noticed clinically.
61 Sometimes the injury is so slight, as in an insect bite that the trauma escapes unnoticed or is
62 forgotten by the patient. When healing takes place, the implanted epithelial cells multiply,
63 producing a central mass of keratin and lipid-rich debris.^{3,5}

64 The best available explanation of the existence of these cysts is the epithelium implant
65 theory. According to this theory, such a cyst originates as an implant of covering epithelium
66 in deeper structures, thereby becoming independent of parent structures but still carrying on
67 its normal keratin-forming function of which the end product is a true epidermal cyst.
68 Conceivably the implant could be caused by trauma or by embryonal inclusion.⁶

69 These cysts occur most often in patients in their second or third decade of life. It is
70 usually solitary and connects with the surface by keratin-filled pores. Clinically, the lesion
71 presents as a slow-growing asymptomatic mass, usually located in the midline, above or
72 below the mylohyoid muscle. When located above the muscle, the cyst manifests itself as a
73 sublingual swelling; when below the muscle, the clinical aspect will be a submental
74 swelling.⁷ Archer says that these cysts are frequently located in the ovaries and testicles, but
75 he adds that they also can be found occasionally, they are found in the floor of the mouth, the
76 palate, and the tongue.⁴

77 Histologically, they can be further classified as epidermoid cysts, dermoid cysts and
78 teratoid cysts.⁸ Most reported cases have involved the floor of the mouth (sublingual
79 dermoids), usually in the midline.⁹⁻¹⁴ Rare cases have been reported in the tongue, lips,¹⁵
80 uvula,¹⁶ temporomandibular joint dermal grafts,¹⁷ intradiploic,¹⁸ buccal mucosa,
81 intracrania^{15, 18} and only three case reports of intraosseous epidermoid cysts were reported.^{3, 4}

82 The radiological differential diagnosis of unicystic lesions of the body of the mandible
83 was made which includes odontogenic keratocyst, unicystic ameloblastoma, early stage of
84 cemento-ossifying fibroma and aneurysmal bone cyst.¹⁹ Odontogenic keratocyst is
85 characterized by a distinct sclerotic margin which may be smooth or scalloped. These tend to
86 spread along the medullary space causing only slight expansion until a considerable size is
87 reached.²⁰ Our case was in contrast to this which showed bicortical expansion. Unicystic
88 ameloblastoma occurs commonly in the body of the mandible. These are well-corticated
89 without a sclerotic margin but in our case a well-defined sclerotic margin was seen. Early
90 stage of cemento-ossifying fibroma presents as well-defined radiolucency with sclerotic
91 margins mostly found at the premolar and molar regions of the mandible.¹⁹ Aneurysmal bone
92 cyst usually produces a radiolucent ovoid or fusiform unilocular expansion of bone and may
93 balloon the cortex.²⁰ These two entities were excluded with the aspiration of white colored
94 keratin material from the cystic space of our case.


95

96 **References:**

- 97 1) Koca H, Seckin T, Sipahi A, Kazanc A. Epidermoid cyst in the floor of the mouth:
98 Report of a case. *Quintessence Int* 2007; 38:473-477.
- 99 2) Louis PJ, Hudson C, Reddi S. Lesion of Floor of the Mouth. *J Oral Maxillofac Surg*
100 2002; 60:804-807.
- 101 3) Noffke CEE. Implantation-type epidermoid cyst of the mandible. *Dentomaxillofac*
102 *Radiol* 1999; 28:383-385.
- 103 4) Torres JS, Higa TT, Epidermoidal cysts in the oral cavity. *J Oral Surg* 1970; 30: 592 -
104 600.
- 105 5) Golden BA, Zide MF. Cutaneous Cysts of the Head and Neck. *J Oral Maxillofac Surg*
106 2005; 63:1613-1619.
- 107 6) Kellen EE. Oral epidermal cysts and probable histogenesis. *Oral Surg Oral Med Oral*
108 *Pathol* 1965; 19:359-364.
- 109 7) Jham BC, Duraes GV, Jham AC, Santos CR. Epidermoid Cyst of the Floor of the
110 Mouth: A Case Report. *J Can Dent Assoc* 2007; 73:525-528.
- 111 8) Ozan F, Polat HB, Ay S, Goze F. Epidermoid Cyst of the Buccal Mucosa:A Case
112 Report . *J Contemp Dent Pract* 2007; 8:90-96.
- 113 9) Worley CM, Laskin DM. Coincidental sublingual and submental epidermoid cysts. *J*
114 *Oral Maxillofac Surg* 1993; 51:787-790.

- 115 10) Zachariades N, Skoura-Kafoussia C. A life-threatening epidermoid cyst of the floor
116 of the mouth: report of a Case. *J Oral Maxillofac Surg* 1990; 48:400-403.
- 117 11) Calderon S, Kaplan I. Concomitant sublingual and submental epidermoid cyst: a case
118 report. *J Oral Maxillofac Surg* 1993; 51:790-792.
- 119 12) De Ponte FS, Brunelli A, Marchetti E, Bottini DJ. Sublingual epidermoid cyst. *J*
120 *Craniofac Surg* 2002; 13:308-310.
- 121 13) Walstad WR, Solomon JM, Schow SR, Ochs MW. Midline cystic lesion of the floor
122 of the mouth. *J Oral Maxillofac Surg* 1998; 56:70-74.
- 123 14) Cortezzi W, De Albuquerque EB. Secondarily infected epidermoid cyst in the floor of
124 the mouth causing a life-threatening situation: report of a case. *J Oral Maxillofac Surg* 1994;
125 52:762-4.
- 126 15) Rajayogeswaran V, Eveson JW. Epidermoid cyst of the buccal mucosa. *Oral Surg*
127 *Oral Med Oral Pathol* 1989;67:181-184.
- 128 16) Yoshinari M, Nagayama M. Epidermoid cyst of the uvula: report of a case. *J Oral*
129 *Maxillofac Surg* 1986;44:828-829.
- 130 17) Weinberg S, Kryshchalskyj B. Epidermoid cyst in a temporomandibular joint dermal
131 graft: report of a case and review of the literature. *J Oral Maxillofac Surg* 1995;53:330-332.
- 132 18) Sudhakar N, Stephenson GC. Swelling on the head- a forgotten lesson: a case report
133 of an intradiploic epidermal cyst with an iatrogenic complication. *Br J Oral Maxillofac Surg*
134 2004;42:155-157.
- 135 19) N.K. Wood and P.W. Goaz, Editors, *Differential diagnosis of oral and maxillofacial*
136 *lesions* (5th edn), St. Louis: Mosby, 1997.
- 137 20) Shear M, Speight P. *Cysts of the Oral and Maxillofacial Regions*(4th edn), Singapore:
138 Blackwell Munksgaard, 2007.


139
140



141

142 **Figure 1: OPG showing well- defined radiolucent lesion with a sclerotic margin**

143




144

145 **Figure 2: CT scan showing Buccal and Lingual Plate expansion**

146

147




148

149 **Figure 3: 3d Reconstruction showing Buccal and Lingual Plate expansion**

150

151

152



153

154 **Figure 4: Histopathology showing thick keratin layer resembling epidermis**