

## Case Study

# *Hydrangea arborescence* and *Berberis vulgaris* Ease Ureteric Calculi Emission: Single Case Report

### Abstract:

Renal calculi or nephrolithiasis refers stone in the kidney-ureter-bladder (KUB) region. Calcium oxalate is the chief ingredient to form most of the calculi and kidney stone is one of the most common health problems in Bangladesh as well as in the world. Multiple risk factors play important role in the formation of stone including fatty foods, increased intake of NSAID, sedentary life and few chronic clinical conditions. Recurrent infection in the urinary tract and dull aching pain in the flank is a most common symptom along with difficulties in urine output and may lead to sudden obstruction of urine flow if not treated properly. Conventionally surgery is the most prescribed solution however, there are few tangles of this including fear to surgery, expenses, and recurrence. Therefore, finding an alternative, more beneficial and less expensive for patients is essential. The current case study involves a single case of ureteric calculi complicated with enlarged prostate, increased creatinine and previously diagnosed diabetes mellitus, where *Hydrangea arborescence* and *Berberis vulgaris* used with a positive outcome. The detailed mechanism of these two could be determined with extensive study along with new verge of research get opened.

**Keyword:** Renal calculi, ureteric calculi, hydrangea, berbaris.

### 1. INTRODUCTION:

The term nephrolithiasis represents calculi or stone in KUB and ureteric calculi defines calculi in the ureter, the tube carries urine from kidney to the urinary bladder (1, 2). The abnormal concretion of various mineral salts including calcium oxalate, calcium phosphate, uric acid and struvite, where about 80% of kidney stone formed chiefly by calcium oxalate and grossly all originate attached to the interstitial plaque over renal papilla (3, 4). In major health problem index, it is in 3<sup>rd</sup> position due to its high prevalence and recurrence (5, 6) though prevalence in Bangladesh is not known, yet it's a very common health complaint (7). Multiple risk factors are involved in renal stone diseases including excessive use of NSAID, increased intake of fatty foods, Chrones's disease, adenine phosphoribosyltransferase deficiency, cystinuria, Dent disease, familial hypo-magnesemia with hyper-calciuria, reduced physical activity and nephron-calcinosis (8, 9). Although kidney stone disease may remain asymptomatic for years, features of recurrent infection in urinary tract along with frequent pain in lower abdomen and flank, nausea-vomiting and obstruction in renal flow are common (10, 11). Untreated patients may suffer sudden obstruction in renal output, however, surgery is widely prescribed treatment which is less eventful yet fever, hematuria, ileus, and urine leak are common complications (7, 12). Homeopathic medicine could be well accepted alternative treatment for renal calculi patients. Homeopathic medicine is based on symptom similarity, mostly depending on patient's mental and physical makeup, not only depending on clinical symptoms. Usually, it takes 12 weeks to 24 or more weeks to get a positive outcome in stone disease, depending on size-shape and position of stone. Sometimes patients become bored to take medicine due to this long period to respond and sometimes patient give up before the treatment has completed. So, it will be beneficial to the patients gets a positive result by consuming both less time and expenses. In this current study, we applied a combination of two different medicine, *Hydrangea arborescens*, and *Berberis vulgaris*, in a single ureteric calculi patient. The patient observed complete recovery of the condition with 4 weeks of consecutive treatment.

46

## 47 2. PRESENTATION OF CASE:

48 Male, 52 years old, having known history of diabetes mellitus (DM) and hypertension (HTN) presented  
 49 with low back pain, more on the left side which was radiating towards the groin, suffering from these for  
 50 last 3-4 days. He had periods of moderate to severe spasms in the renal passages, which refers it as  
 51 renal colic. The Patient was advised for some laboratory tests and with ultrasonogram diagnosed as a  
 52 case of left ureteric calculi in the distal part measuring about 0.66 mm, along with mild left  
 53 hydroureteronephrosis (Fig.1A), along with enlarged prostate with post void residue (PVR) calculated  
 54 20.64 ml (Fig.3), fatty liver change in grade-I. From the biochemical analysis of blood revealed that his  
 55 prostate-specific antigen (PSA) was 0.74 ng/dl which suggest enlarged prostate is benign in nature;  
 56 serum creatinine was 2.22 mg/dl (Fig. 4) and fasting blood glucose (FBS) was 10.4 mmol/L (Fig. 5). Due  
 57 to fear to go to under surgery, the patient was inclined to take homeopathic medicine as a popular  
 58 alternative choice in this locality.

## 59 3. MEDICINES AND DOSE:

60 After being diagnosed the patient was advised to take *Hydrangea arborescence*, orally in tincture form, 60  
 61 ml distributed in 30 equal doses (2ml each time, twice a day, before meal) and *Berberis vulgaris*, orally in  
 62 3X potency, 0.5 ml mixed with 60 ml distilled water, distributed in 30 equal doses (2ml each time, twice a  
 63 day, after meal) for 15 days. Evaluating the latest sonographic image found (Fig.1B) and comparing  
 64 biochemical report with previous reports (Fig.3, Fig.4, Fig.5) same medicines were prescribed for next 15  
 65 days in given order. Close image monitoring is done with computed tomography (Fig.1C), and a final  
 66 sonographic image was taken at the end of four weeks of treatment (Fig.1D).

67 During this 4 weeks of a continuous period of treatment, use of other homeopathic or conventional  
 68 medicine, topical or internal, were prohibited.

69 The patient was asked to take less quantity of water as he was taking earlier and advised to drink 2 Lt/24  
 70 hours along with maintaining personal hygiene and diabetic food chart strictly.

71

## 72 4. RESULTS

73 We observed that a large stone flowed out through urethra within 4 weeks (Fig: 2); PVR, elevated serum  
 74 creatinine and FBS level started to reduce (Fig.3, Fig.4 , and Fig.5). A small ureterocele observed in the  
 75 left bladder causing no significant complaints (Fig.1D). The frequency of renal colic reduced within first 2  
 76 (two) weeks and lastly felt few hours prior to stone expulsion. No change found in the prostate status and  
 77 remaining hydronephrosis condition needed no further treatment as it may resolve with the given advice.



78

79 **Figure 1: Follow-up images of calculi, marked in the yellow ring. A) Before starting treatment, calculi found in the distal part**  
 80 **of the left ureter, B) Two weeks later, calculi in the left uretero-vesicular junction C) Three weeks later, D) Four weeks later,**  
 81 **calculi free sonographic image.**

82



Figure 2: Flowed out calculi, length-18 mm, and width- 9 mm.




Figure 3: Post void residue (PVR) during the course of treatment. Initially, it was 20.64 ml, reduced to 11.87 ml after 2 weeks and 12 ml after 3 weeks of treatment.




Figure 4: Serum creatinine level during the treatment period. Before medication, it was 2.22 mg/dl, which moved to 2.53 mg/dl after 2 weeks of treatment. After 3 weeks it reduced to 1.93 mg/dl and after 4 weeks it was found at 1.62 mg/dl.




Figure 5: Fasting blood glucose level during the treatment period. Initially, it was 10.4 mmol/L, which 2 weeks later dropped to 9.6 mmol/L ; after 3 weeks it was at 8.4 mmol/L and at the end of 4 weeks of treatment it moved to 8.0 mmol/L.

## 5. DISCUSSION

Among various available treatment options, homeopathy is one with non-invasive, cost-effective and without any known side effects. A number of homeopathic drugs including Benzoicum acidum, *Lycopodium clavatum*, *Calcarea carbonicum*, *Berberis vulgaris*, *Sarsaparilla officinalis* etc. are prescribed in treating nephrolithiasis (13). According to homeopathic principles, the application of more than one drug is prohibited (14). Now a day, in some particular clinical condition the application of two medicines at a time is practiced to reduce the duration of time to cure (15). In our study, we treated with *Hydrangea arborescens* for its various properties including cracking stone in small pieces or leaching stone to softer and smaller, increases urinary outflow and reduces the prostate size (16-18). Along with this, *Berberis vulgaris* was given for its known antibacterial activity, delays or prevent stone formation and has the affinity to help urinary tract health including stone expulsion (16, 19, 20). After two weeks, the stone found in the left uretic-vesicular-junction, measuring 15 mm, which may miss interpreted that an increase of stone size, though it is the same stone in soft, flattened and elongated shape which justify the use of *Hydrangea arborescence* (Fig.1B). We noticed that flowed out stone is almost 18 mm in length and 9 mm in width (Fig.2), and the recovery time took significantly shorter than we usually find in using the single drug. Noticeably gradual reduction of PVR (Fig.3) and FBS (Fig.5) found; though after two weeks of treatment serum creatinine raised a little, which gradually reduced after three and four weeks respectively (Fig.4). Recommended dilution in homeopathy is not fixed, could be given frequently according to individuals severity of illness, using various potencies including both higher and lower potency is common (21). In our study, we used the given potencies throughout the study with the significant outcome.

## 6. CONCLUSION:

Findings of this case study may help to practice a new treatment option to reduce the curing time and expenses in the treatment of ureteric calculi and it's related complicacy. Further study is needed to determine the possible correction of using both of these drugs and justification of this application.

## CONCENT:

Author declares that written informed consent was obtained from the patient for publication of this article and accompanying images. He was not against in taking homeopathic treatment as it is already famous in Bangladesh.

## ETHICAL CLEARANCE

Not applicable in this case.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Venes D. Taber's Cyclopedic Medical Dictionary, Illustrated in full color. FA Davis Company; 2009.
2. Standring S, editor. Gray's anatomy e-book: the anatomical basis of clinical practice. Elsevier Health Sciences; 2015 Aug 7.
3. Serter S, Sahin E, Yildirim UM, Arslan M, Karakose A. Unenhanced computed tomography findings of renal papillae in patients with a ureteral stone. Clinical & Investigative Medicine. 2016 Dec 1;39(6):141-6.
4. Boutwell E, Stine R, Tucker K. Effect of prosthetic gel liner thickness on gait biomechanics and pressure distribution within the transtibial socket. Journal of Rehabilitation Research and Development. 2012 Feb 10;49(2):227.
5. Papadoukakis S, Stolzenburg JU, Truss MC. Treatment strategies of ureteral stones. EAU-EBU update series. 2006 Oct 1;4(5):184-90.

6. Tanagho E, McAninch J. Smith's general urology. McGraw-Hill Prof Med/Tech; 2007 Nov 21.
7. Rahman MM, Shamim IA, Kormokar U, Chowdhury SA, Gupta SD. Outcome of extracorporeal shock wave lithotripsy in the treatment of mid ureteric stones. Bangladesh Medical Journal. 2014 Dec 30;43(2):72-8.
8. Edvardsson VO, Goldfarb DS, Lieske JC, Beara-Lasic L, Anglani F, Milliner DS, Palsson R. Hereditary causes of kidney stones and chronic kidney disease. Pediatric nephrology. 2013 Oct 1;28(10):1923-42.
9. Fagagnini S, Heinrich H, Rossel JB, Biedermann L, Frei P, Zeitz J, Spalinger M, Battegay E, Zimmerli L, Vavricka SR, Rogler G. Risk factors for gallstones and kidney stones in a cohort of patients with inflammatory bowel diseases. PloS one. 2017 Oct 12;12(10):e0185193.
10. Boyce CJ, Pickhardt PJ, Lawrence EM, Kim DH, Bruce RJ. Prevalence of urolithiasis in asymptomatic adults: objective determination using low dose noncontrast computerized tomography. The Journal of urology. 2010 Mar 1;183(3):1017-21.
11. Nabi G, Downey P, Keeley F, Watson G, McClinton S. Extra-corporeal shock wave lithotripsy (ESWL) versus ureteroscopic management for ureteric calculi. Cochrane Database Syst Rev. 2007 Jan 24;1.
12. Dongol UM, Limbu Y. Safety and Efficacy of Percutaneous Nephrolithotomy in Children. Journal of Nepal Health Research Council. 2017 Sep 15;15(2):130-4.
13. Schroyens F. *Synthesis 10.0 General Repertory Millenium*, version 2010
14. Hahnemann HC, Boericke W. Organon of Medicine. 6th ed. New Delhi: B. Jain Publishers; 2012.
15. Mercaldo M. How to prescribe Medorrhinum: the frequency of symptoms and signs in homeopathic patients. British Homoeopathic Journal. 1999 Apr 1;88(2):69-77.
16. Vermueulen F. Concordant Materia Medica. Dev Nagar, New Delhi: Indian Books & Periodicals Publishers; 2001
17. Horne S. Hydrangea (Hydrangea arborescens). Tree of Light. 2010. Accessed 4 February 2018. Available: [https://www.treeelite.com/articles/articles/hydrangea-\(hydrangea-arborescens\).html](https://www.treeelite.com/articles/articles/hydrangea-(hydrangea-arborescens).html)
18. Hughes L. Hydrangea: An Herbal Treatment for Kidney Stones. Epoch times. 2013. Accessed 4 February 2018. Available: [https://www.theepochtimes.com/hydrangea-an-herbal-treatment-for-kidney-stones\\_185520.html](https://www.theepochtimes.com/hydrangea-an-herbal-treatment-for-kidney-stones_185520.html)
19. Peng L, Kang S, Yin Z, Jia R, Song X, Li L, Li Z, Zou Y, Liang X, Li L, He C. Antibacterial activity and mechanism of berberine against Streptococcus agalactiae. International journal of clinical and experimental pathology. 2015;8(5):5217.
20. Jyothilakshmi V, Thellamudhu G, Kumar A, Khurana A, Nayak D, Kalaiselvi P. Preliminary investigation on ultra high diluted B. vulgaris in experimental urolithiasis. Homeopathy. 2013 Jul;102(03):172-8.
21. Rawat PS, Select Your Dose and Potency, 11<sup>th</sup> impression, New Delhi: B. Jain Publishers; 2011