

**Original Research Article****The water infiltration, hydraulic conductivity and water retention effects of ground *Saponaria officinalis* (L.) root as a soil surfactant**1  
2  
3  
4  
5  
6  
7  
8  
9**ABSTRACT**

Climate change is affecting precipitation patterns and intensity; increasing regional drought conditions and reducing precipitation infiltration times, respectively. The use of soil surfactants presents an opportunity to improve soil water content and infiltration in soils. As aridity and drought vulnerability increase globally, improving water infiltration and retention is becoming increasingly important for agriculture as water resources are scarce and climate change shifts precipitation patterns. While surfactants are widely available for agricultural use, most or all are unapproved in sustainable and organic production. Ground *Saponaria officinalis* (L.) root produces saponins, natural surfactants used in several industrial applications, including soap and soil contaminant recovery. To determine its potential to improve soil water interactions its effects on soil hydraulic conductivity, water content, infiltration and drainage rates were tested in washed sand, heavy clay soil and clay loam soil. When compared to untreated soils, drainage and infiltration was slowed ( $P < 0.05$ ) in sand and loam with soapwort applications while no significant differences in any variable were present in clay soil compared to any treatment. Soil water content was not significantly different in any treatment. While soapwort did not increase infiltration rates it did markedly slow drainage rates in sand and loam. The benefit of this may be realised as longer opportunity for plant available water in the root zone.

10  
11  
12  
13  
14  
15  
16  
17  
18  
19

*Keywords: drought vulnerability, ground soapwort, soil surfactant, soil water content*

**1. INTRODUCTION**

Global cropland is estimated at 1.82 billion hectares, 455 million hectares are considered dryland [1]. Further, more than 30% of the worlds crop supply is produced on irrigated lands [2]. Water is fast becoming a resource restrain in crop production with increased groundwater depletion and climate change.

20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30

Most major arid region aquifers are being over-drafted with depletion leaving residual low quality water [3, 4]. The wider impacts of global climate change on water availability are the increases of variability in seasonal precipitation [5, 6]. There are indications the variability not the overall amount of rainfall has and will continue to change [6]. This includes a reduction in the duration and increase in the intensity of precipitation events [7]. This has led to increased runoff and reduced infiltration globally, with North America experiencing greater runoff than most other continents [8]. Regionally, the southwestern U.S. droughts are strongly linked to El Nino events; whether these represent increasingly common occurrences, long-term or cyclic events, El Nino cycling appears to be responsible for the ongoing severe to exceptional drought in the arid U.S. southwest (2009 to 2015) [9, 10]. The IPCC projects continued drought over the coming century [7].

31 In arid regions soil properties are greatly influenced by development in a dry climate.  
32 They tend to lack organic matter because of low productivity [11]. Lack of organic matter  
33 inhibits the development of aggregates, reduces porosity and water retention. These soils  
34 also contain higher concentrations of soluble ions, such as  $\text{Ca}^{2+}$  and  $\text{Na}^+$ , because of lack of  
35 precipitation driven leaching. Arid developed soils also tend to have clays with high shrink-  
36 swell capacity, which increase the tendency for sealing [12]. Sealing reduces the  
37 opportunity for water to infiltrate and increases runoff losses. Plant waxes are also more  
38 common in these soils and they coat soil particles with a hydrophobic film [13]. All soil types  
39 can have hydrophobic conditions present, but some soils are more prone and hydrophobic  
40 soils are now seen as more common than previously thought [14, 15]. This hydrophobicity  
41 causes soils to repel water rather than infiltrate readily. Thus arid soils have multiple factors  
42 reducing water infiltration.

43 Soil surfactants offer several opportunities to improve soil water management. One  
44 is reducing the infiltration time by attaching to the hydrophobic tails of the repellent coatings  
45 of the soil surface and aggregates, leaving the hydrophilic head exposed to infiltrating water.  
46 They also reduce surface tension of water and allow for freeing movement of the water into  
47 soil pores. Lastly, surfactants behave as an adsorbent, holding water in the soil pores,  
48 reducing the soil water drainage time, thus increasing the water volume and contact time for  
49 plant roots.

50 By reducing surface tension of water they increase the rate of infiltration. Typically  
51 agricultural use of soil surfactants has been exclusively in turf management for athletic fields  
52 and golf courses [15]. However, more recently there has been an increase in the use of soil  
53 surfactants to improve water use in agriculture [16, 17]. They have been used to increase  
54 infiltration, increase soil water content, and therefore plant available water (PAW), and  
55 generally increase water conservation [15].

56 A soil order with particularly problematic water infiltration issues in arid regions is  
57 Vertisols. They are characterized by a high percentage of montmorillonitic clay, which in dry  
58 periods causes large vertical cracks to appear with spans up to 50cm. These cracks  
59 disappear with precipitation during which time the surface seals. Thus, these are  
60 problematic soils in that they swell so rapidly that infiltration pores quickly close. The high  
61 clay content also reduces PAW by encapsulating a large portion of the soil water in the clay  
62 micropores and the dense clays reduce root penetration. Thus plants struggle to acquire  
63 enough water to maintain metabolic and transpirative needs in arid Vertisols. While not a  
64 dominant soil order, they are very productive with proper management and have a high  
65 cation exchange capacity (CEC). The largest expanses of Vertisols are in arid regions of  
66 south central India, southern Sudan and South Sudan, and eastern Australia. Water  
67 infiltration studies of Vertisols indicate tilled soils have slower infiltration, probably caused by  
68 reduced macropore continuity that results from pulverized soil caused by tillage [18].  
69 However, they are still considered prime agricultural land because the high CEC makes  
70 them very fertile.

71 Soil surfactant application in agricultural soils has shown promise. The use of a soil  
72 surfactant improved infiltration in a poorly drained loamy Crosby soil (Alfisols) with clayey B-  
73 horizons. The result was a 19.4% reduction in runoff [19]. Sandy loams have less shrink  
74 and swell related to clay content, and tend to drain rapidly. The rapid drainage is a result of  
75 larger pore space as sand and silt per cent is higher. However, the organic matter of loamy  
76 soils also contributes to hydrophobic conditions by developing an organic waxy coat on soil  
77 particles. Loamy sands with soil surfactants took more than twice the time to begin runoff  
78 [20].

79 Soil surfactant efficacy is still undergoing debate, depending on which condition is to  
80 be improved. Surfactants can improve soil infiltration [21], change preferential flows [22] or  
81 increase soil water content [23]. Nearly all of the products in use possess similar properties  
82 of a hydrophilic head and hydrophobic tail. Most are short chain organic compounds.  
83 Typically they are marketed to turf and golf management operations, but some are marketed  
84 for application in high value crops like vegetables [24]. All of the products are synthetic in  
85 origin and thus far none appear certified for organic operations.

86 Natural plant derivatives lend themselves to organic certification but there are little to  
87 no studies quantifying the effects of plant-based surfactants on soil water properties.  
88 Surfactant properties can be found in several plant derived products, specifically saponins.  
89 Saponins are present in plants of the family Sapindaceae as well as a few others. Saponins  
90 derived from plant materials have been used for soil contaminant remediation in the past.  
91 Specifically, 10% solutions of *Sapindus mukorossi* (Geartn.) has been tested for use in soil  
92 contaminant remediation with promising results [25]. Commercially saponins are extracted or  
93 derived from, *S. mukorossi*, *Saponaria officinalis* (L.) and *Quillaja saponaria* (Molina). The  
94 compound is amphipathic, thus possessing the hydrophilic head and hydrophobic tail.  
95 Though larger in molecular weight the structure of saponin is similar to synthetic surfactants  
96 with hydroxyl groups at one end and lipophilic carbon rings at the other.

97 The objective of this study was to explore the effects of raw ground *S. officinalis* root  
98 on the rates of infiltration and hydraulic conductivity, and the water holding capacity of two  
99 arid and drought vulnerable local soils series from central Texas. The raw product is  
100 untested but has known surfactants in the form of tripterpenoid saponins [26] and is used as  
101 a surfactant in soap production [27]. The application and results of soils surfactant vary  
102 according to soil types and ground *S. officinalis* surfactant capacity is largely untested. The  
103 purpose of this study was to determine significant differences ( $P < 0.05$ ) in infiltration rate,  
104 hydraulic conductivity or water retention as result of the application of ground *S. officinalis* to  
105 soils or sand when compared to untreated conditions (water only application).

106

## 107 2. MATERIAL AND METHODS

108

109 Two local soils series and washed sand were used in the study. One soil was Houston  
110 Black; a blackland prairie Vertisol defined as fine, montmorillonitic, thermic, Udic, Pellusterts.  
111 Blackland prairie soils are a highly productive agricultural row crop soil in Texas, but are  
112 problematic due to the high shrink swell character and low infiltration rates caused by a high  
113 percentage of clay. The other soil was Tarpley; a gravelly Mollisol defined as clayey,  
114 montmorillonitic, thermic, lithic, vertic, Argiustolls. Tarpley is a Texas Hill Country upland  
115 Mollisols with lesser amounts of clay than Houston Black and typically used for pasture as it  
116 is not very productive, but has potential for rocky soil adapted orchard crops, such as olive  
117 (*Olea eurpoeaea*). Washed sand was used as a control.

118 Soils and sand were preparations were similar. Sand was sieved through a 2 mm  
119 mesh screen and washed in a 0.05 mm mesh screen to remove clays and silts, then dried  
120 for 24 hours at 100° C. Each soil was dried for 24 hours at 100° C, then screened through 2  
121 mm mesh sieve to remove any rock fragments and large organic matter. Bulk density was  
122 taken from the Comal-Hays County WSS Soil Survey: Houston Black  $P_b = 1.35$  and Tarpley  
123  $P_b = 1.27$ . A 40 cm long x 5 cm diameter cylinders were used to hold the soil column. The  
124 bottom was covered with a wire mesh and then covered with filter paper to prevent soil loss.  
125 Dried and sifted soil or sand was placed in the tube. Cylinders were filled to 30 cm with soil  
126 or sand and lightly packed [28]; [16]. Cylinders were suspended above a basin to catch  
127 drainage water.

128 The treatment, powdered *S. officinalis* root is an untested soil surfactant for this  
 129 application. There is little information regarding solution concentrations, therefore we used  
 130 10 g/100 ml (g/g) saponin solution concentration for soil remediation as a baseline [25]. The  
 131 equivalent soil application of dry *S. officinalis* using the 10 g/100 ml concentration would be  
 132 prohibitive in field agriculture; therefore an soil application of 1.0 g powdered *S. officinalis*  
 133 was chosen as a soil treatment. This was chosen as the baseline application with 0.5 g and  
 134 1.5 g as alternative soil applications, with no soil application of soapwort as the control. The  
 135 treatments were knifed into the surface 2-3 cm of soil to mimic a superficial application of a  
 136 dry flowable product followed by soil/turf scarification. Water was released on the surface of  
 137 the soil from a 1 L Marriotte reservoir and a constant head of 3 cm of water above the soil  
 138 surface was maintained [16, 28]. The water level of the reservoir was recorded in cm every  
 139 minute; the start time of drainage was recorded, as was the last drainage time. Drainage  
 140 water volume was recorded after 1 minute after drainage ceased. Each treatment was  
 141 repeated 3 times on fresh, untreated soil or sand. The variables measured were infiltration  
 142 rates, drainage time and water retained. Hydraulic conductivity (K) was calculated from  
 143 these data. A general linear model (GLM) was used to statistically analyse the data. SAS  
 144 software was used to perform the analyses.

145

### 146 3. RESULTS AND DISCUSSION

147

148 The GLM analysis of the sand data for all variables indicated an interaction between the  
 149 treatments: drainage time -  $F = 7.96$ ,  $P = 0.003$ ; infiltration rate -  $F = 8.76$ ,  $P = 0.002$ ; water  
 150 retention -  $F = 6.71$ ,  $P = 0.006$ . Contrasts were then performed between each treatment  
 151 within each variable. For the Tarpley soil the GLM resulted in only the water retention with  
 152 no interaction within the treatments but for clarity contrasts were performed on all: drainage  
 153 time -  $F = 4.0$ ,  $P = 0.034$ ; infiltration rate -  $F = 6.92$ ,  $P = 0.006$ ; water retention -  $F = 0.38$ ,  $P$   
 154  $= 0.815$ . Interactions were only present in water retention for Houston Black, but similar to  
 155 the Tarpley soil, contrast were preformed on all treatments and variables for clarity: drainage  
 156 time -  $F = 1.39$ ,  $P = 0.307$ ; infiltration rate -  $F = 1.31$ ,  $P = 0.330$ ; water retention -  $F = 3.78$ ,  $P$   
 157  $= 0.040$ .

158 Drainage time for sand treated with any surfactant application was significantly  
 159 longer when compared to untreated sand drainage rates (Table 1). Infiltration rates were  
 160 more variable, with all soapwort applications taking longer to infiltrate compared to no  
 161 application water. The 1 g application had the longest infiltration time compared to all  
 162 others. There were no significant differences in water retention based on treatment.

163 Table 1. Contrast analysis of drainage, infiltration and water retention of powdered soapwort  
 164 in washed sand, ( $n=3$ ). P-values are presented for each application contrast in the right  
 165 three columns.

Treatment	Mean drainage rate (min)	StdDev	0.5 g SW	1.0 g SW	1.5 g SW
0.5 g SW	48.29	16.26			
1.0 g SW	63.24	3.61	0.069		
1.5 g SW	53.81	10.51	0.470	0.228	
H <sub>2</sub> O	24.62	0.52	0.009	0.000	0.002

	Mean infiltration rate (min)				
0.5 g SW	10.03	0.93			
1.0 g SW	8.63	0.55	0.198		
1.5 g SW	12.41	2.38	0.040	0.003	
H <sub>2</sub> O	7.74	0.23	0.047	0.400	0.001

	Mean H <sub>2</sub> O retention (ml)				
0.5 g SW	265.00	5.00			
1.0 g SW	241.67	28.87	0.104		
1.5 g SW	246.67	2.89	0.191	0.710	
H <sub>2</sub> O	268.33	2.89	0.804	0.068	0.128

166

167 In contrast to sand, the analysis of the drainage time and infiltration rate for Houston  
 168 Black soil indicated no significant difference for any treatment (Table 2). The analysis of  
 169 water retention though, indicated significant differences based on treatment applications with  
 170 water retention of Houston Black treated with 1 g application of soapwort retained  
 171 significantly more water than the 1.5 g application.

172 Table 2. Contrast analysis of drainage, infiltration and water retention of powdered soapwort  
 173 in Houston Black. (*n*=3). P-values are presented for each application contrast in the right  
 174 three columns.

175

Treatment	Mean drainage rate (min)	StdDev	0.5 g SW	1.0 g SW	1.5 g SW
0.5 g SW	102.81	25.91			
1.0 g SW	113.35	29.94	0.666		

1.5 g SW	114.31	15.97	0.638	0.968	
H <sub>2</sub> O	151.84	41.13	0.065	0.135	0.144

Mean  
infiltration  
rate (min)

---

0.5 g SW	93.62	18.60			
1.0 g SW	111.15	77.91	0.691		
1.5 g SW	89.39	7.21	0.923	0.622	
H <sub>2</sub> O	170.19	84.61	0.104	0.198	0.088

Mean H<sub>2</sub>O  
retention  
(ml)

---

0.5 g SW	416.67	20.82			
1.0 g SW	455.00	30.41	0.116		
1.5 g SW	403.33	37.53	0.563	0.043	
H <sub>2</sub> O	420.00	30.00	0.884	0.148	0.472

176

177 Tarpley soils are loams, with more sand and silt than Houston Black soils.  
 178 Treatment effects were prevalent for drainage time and infiltration rate, but not water  
 179 retention (Table 3). The soapwort treatments all had significantly longer drainage times  
 180 compared to water only applications. Infiltration rates were up to 3 times slower with all  
 181 soapwort applications taking significantly longer than water. There were no differences in  
 182 water retention for any application.

183 Table 3. Contrast analysis of drainage, infiltration and water retention of powdered soapwort  
 184 in Tarpley. (*n*=3). P-values are presented for each application contrast in the right three  
 185 columns.

186

Treatment	Mean drainage rate (min)	StdDev	0.5 g SW	1.0 g SW	1.5 g SW
-----------	--------------------------------	--------	----------	----------	----------

---

0.5 g SW	160.02	11.30			
1.0 g SW	192.56	67.86	0.274		
1.5 g SW	178.10	33.08	0.535	0.618	
H <sub>2</sub> O	92.82	7.53	0.038	0.005	0.012

Mean  
infiltration  
rate (min)

---

0.5 g SW	183.16	44.89			
1.0 g SW	203.22	60.98	0.536		
1.5 g SW	193.72	28.43	0.743	0.767	
H <sub>2</sub> O	61.75	17.84	0.003	0.001	0.001

Mean H<sub>2</sub>O  
retention  
(ml)

---

0.5 g SW	403.33	41.63			
1.0 g SW	371.67	54.85	0.410		
1.5 g SW	376.67	60.28	0.485	0.894	
H <sub>2</sub> O	380.00	27.84	0.540	0.825	0.929

187

188 Hydraulic conductivity for each treatment and soil was calculated using Darcy's Law:  
 189  $k=[Q/(A*t)]*(H/L)$ . As would be expected based on the analysis of time for complete  
 190 drainage mean hydraulic conductivity in sand was significantly slower for all soapwort  
 191 applications compared to water only (Table 4a). There was no apparent difference in  
 192 hydraulic conductivity between soapwort treatments. Similarly, the effects of soapwort on  
 193 hydraulic conductivity in Houston Black clay soils mirrored the drainage times with no  
 194 difference between treatments and control (Table 4b). The effects of soapwort on K in  
 195 Tarpley loam soil were similar to sand significantly slower K in soapwort treatments; slowing  
 196 by about half with any application of soapwort (Table 4c).

197 Table 4. GLM contrast analysis of mean hydraulic conductivity in sand (a), Houston Black  
 198 (b) and Tarpley (c), ( $n = 3$ ). P-values are presented for each application contrast in the right  
 199 three columns.

200 a. Mixed effects  $F = 18.87$ ,  $P = 0.0005$

Treatment	Mean hydraulic conductivity (K) $\text{cm sec}^{-1}$	StdDev	0.5 g SW	1.0 g SW	1.5 g SW
0.5 g SW	0.0012	0.00040			
1.0 g SW	0.0009	0.00000	0.131		
1.5 g SW	0.0011	0.00026	0.520	0.343	
H <sub>2</sub> O	0.0023	0.00006	0.000	0.000	0.000

201

202 b. Mixed effects  $F = 1.21$ ,  $P = 0.3680$

Treatment	Mean hydraulic conductivity (K) $\text{cm sec}^{-1}$				
0.5 g SW	0.00045	0.00014			
1.0 g SW	0.00037	0.00007	0.356		
1.5 g SW	0.00040	0.00003	0.515	0.773	
H <sub>2</sub> O	0.00030	0.00010	0.098	0.397	0.267

203

204 c. Mixed effects  $F = 14.77$ ,  $P = 0.0013$

Treatment	Mean hydraulic conductivity (K) $\text{cm sec}^{-1}$				
0.5 g SW	0.00028	0.000025			
1.0 g SW	0.00027	0.000087	0.764		
1.5 g SW	0.00026	0.000025	0.708	0.940	
H <sub>2</sub> O	0.00050	0.000047	0.000	0.000	0.000

205



#### 206 4. CONCLUSION

207

208

209

210

211

212

213

214

215

This study offers only a glimpse of possibilities of soapwort as a surfactant. Soil surfactants have been proposed as treatments to increase water retention in soils, slow drainage and improve water use efficiency [29]. In compliance with our current social demands to find sustainable approaches to agriculture the use of natural plant based products appears to be a good alternative to synthetics, especially considering USDA Organic Standards do not provide for synthetic surfactant use (USDA). The soapwort application is similar to commercial soil surfactant applications in which no soil surfactant increased water content in loams or sands significantly compared to no treatment [16].

216

217

218

219

220

221

222

223

224

225

226

In the washed sand the drainage time was significantly longer, as was K with any application of soapwort, which could mean more opportunity for PAW during that period, however the applications of soapwort increased the infiltration time in sand. Similar effects on K were found with an anionic (Sulphonic) surfactant in a very sandy loam [28]. Total water retention based on the soapwort applications did not differ in the sand, thus even though drainage time was longer, post drainage PAW may not differ. Again this is similar to others where applications of surfactants did not increase soil water retention compared to water control [28] [30]. In the sand treated with 1.5 g soapwort the water infiltration times were significantly longer compared to the other soapwort applications and water; and only the 1.0 g application infiltration rate no different from water. These results conflict with those where no significant difference in infiltration rates occurred [16].

227

228

229

230

231

232

233

234

235

236

237

238

239

Sandy soils are particularly difficult to wet evenly [31] and drain very quickly, reducing PAW and opportunity for crops to maintain turgor. In sandy soils, prevalent in south Florida field vegetable production, there may be an application for soapwort, specifically for even seedling emergence [32]. The use of soils surfactants have been shown to increase soil water content in sandy soils by up to 3 times [15], however none of the soapwort applications increase the soil water content in sand. Soil surfactants have also proven useful in turf greens management by decreasing infiltration time in sandy soils [21]. While soapwort actually appears to increase the infiltration time in sand, this study indicates drainage times could be extended by up to two times or more compared to untreated soils increasing opportunity for plants to uptake water (Table 1 and 4a). Though hydrogels are not surfactants per se, they do help soils retain water and in a sandy loam they increased the soil water content and number of days to the permanent wilting point in barley, wheat and chickpea fields [33].

240

241

242

243

244

245

246

247

248

The Tarpley soils in this study is a clay loam and compared to the heavy clay Houston Black soil drainage times for water are very fast, but slower than washed sand. All soapwort applications increased the drainage time and infiltration rate significantly when compared to water. Others have found no differences in infiltration in their loam soil type based on the surfactants in their study [16]. Also differing from the soapwort applications are results with no significant differences in the drainage time with the use of surfactants [34]. K was significantly slower for all soapwort applications in the loam soil. Others have found all surfactant applications in their study resulted in slower K [28]. Soapwort did not increase the retained water, which is similar to many [16, 28, 30, 34].

249

250

251

252

253

Heavy clay soils have very high porosity, but very low pore sizes, thus they tend to drain very slowly. Though the actual times for drainage were much faster (40 minutes) with the soapwort treatments the analysis did not indicate a significant difference in drainage times for any application. Similarly the results for infiltration rates, K and water retention in the clay soil indicated no significant differences based on soapwort applications. The results

254 of soapwort applications concur with the results of [34] and the surfactants applied to clay  
255 soil in their study.

256 Soapwort shows some promise slowing infiltration, reducing drainage rates and  
257 slowing K in loams or sands. It shows no apparent ability to improve water retention in any  
258 soil type used here. In clay soil soapwort had no impact on any soil water interactions  
259 compared to water. There may be application in fast draining soil types to help reduce  
260 drainage times, but only in mesic regions as drought prone regions may require the slow  
261 drainage to prevent plant water stress.

262

## 263 REFERENCES

264

265 1. Hassan R, Scholes R, Ash N. Ecosystems and human well-being: Current state and  
266 trends: Findings of the Condition and Trends Working Group. Washington: Island Press  
267 2005.

268 2. FAO. 2002; Accessed 1 May 2015. Available:

269 [http://www.fao.org/docrep/005/y3918e/y3918e10htm#P0\\_0](http://www.fao.org/docrep/005/y3918e/y3918e10htm#P0_0)

270 3. Gleeson T, Wada Y, Bierkens MF, van Beek LP, Water balance of global aquifers  
271 revealed by groundwater footprint. *Nature*. 2012;488:197-200.

272 4. Konikow LF and Kendy E. Groundwater depletion: A global problem. *Hydrogeol J*.  
273 2005;13:317-320.

274 5. Easterling DR, Evans JL, Groisman PY, Karl TR, Kunkel KE, Ambenje P. Observed  
275 Variability and Trends in Extreme Climate Events: A Brief Review. *B Am Meteorol Soc*.  
276 2000;81:417-425.

277 6. Trenberth KE, Dai A, Rasmussen RM, Parsons DB. The changing character of  
278 precipitation *B Am Meteorol Soc*. 2003;84:1205-1217.

279 7. Meehl GA, Stocker TF, Collins WD, Friedlingstein P, Gaye AT, Gregory JM, et al. Global  
280 climate projections. In: Solomon S, Qin D, Manning M, Chen Z, Marquis M, Averyt KB,  
281 Tignor M, Miller HL, editors. *Climate Change 2007: The Physical Science Basis Contribution*  
282 *of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on*  
283 *Climate Change Cambridge: Cambridge University Press. 2007.*

284 8. Labat D, Godd ris Y, Probst JL, Guyot JL. Evidence for global runoff increase related to  
285 climate warming. *Adv Water Res*. 2004;27:631-642.

286 9. Cook ER, Seager R, Heim RR, Vose RS, Herweijer C, Woodhouse C. Megadroughts in  
287 North America: Placing IPCC projections of hydroclimatic change in a long-term  
288 palaeoclimate context. *J Quat Sci*. 2010;25:48-61.

289 10. Trenberth KE. Framing the way to relate climate extremes to climate change *Climatic*  
290 *Change*. 2012;115:283-290.

291 11. Schillinger, WF, Papendick, RI, Guy, SO, Rasmussen, PE, Van Kessel, C. Dryland  
292 Cropping in the Western United States In: Peterson GA, Unger PW, Payne WA editors  
293 *Dryland agriculture Madison WI, American Society of Agronomy; 2006.*

- 294 12. Morin J Soil crusting and sealing In: Soil tillage in Africa: Needs and challenges FAO  
295 Soils Bulletin 69 1993; Accessed 1 January 2016. Available:  
296 [http://www.fao.org/docrep/t1696e/t1696e06htm#P232\\_14979](http://www.fao.org/docrep/t1696e/t1696e06htm#P232_14979) ISBN 92-5-103442-7
- 297 13. DeBano LF. Water repellent soils: A state-of-the-art Pacific Southwest Forest and Range  
298 Experiment Station. Volume 46 USDA, Forest Service 1981; Accessed 1 January 2016.  
299 Available: [http://svinet2fsfedus/psw/publications/documents/psw\\_gtr046/psw\\_gtr046pdf](http://svinet2fsfedus/psw/publications/documents/psw_gtr046/psw_gtr046pdf)
- 300 14. Dekker LW, Oostindie K, Ritsema CJ. Exponential increase of publications related to soil  
301 water repellency. Soil Res. 2005;43:403-441.
- 302 15. Moore D, Kostka S, Boerth T, Franklin M, Ritsema C, Dekker L, et al. The effect of soil  
303 surfactants on soil hydrological behavior, the plant growth environment, irrigation efficiency  
304 and water conservation. J Hydrol and Hydromechanics. 2010;58:142-148.
- 305 16. Mobbs TL, Peters RT, Davenport JR, Evans MA, Wu JQ. Effects of four soil surfactants  
306 on four soil-water properties in sand and silt loam J of Soil and Water Conserv. 2012;67:275-  
307 283.
- 308 17. Oostindie K, Dekker LW, Wesseling JG, Ritsema CJ, Moore D. Influence of a single soil  
309 surfactant application on potato ridge moisture dynamics and crop yield in a water repellent  
310 Sandy soil. Acta Hortic. 2012;938:341-346.
- 311 18. Potter KN, Torbert HA, Morrison JE. Tillage and residue effects on infiltration and  
312 sediment losses on vertisols. T ASAE. 1995;38:1413-1444.
- 313 19. Sepúlveda NB. Wetting agents and irrigation water conservation: efficacy for golf course  
314 fairways and identification of management practices [MS Thesis] Cranfield University.  
315 Cranfield, England; 2004.
- 316 20. Mitra S, Vis E, Kumar R, Plumb R, Fam M. Wetting agent and cultural practices increase  
317 infiltration and reduce runoff losses of irrigation water Biologia, 2006;61:S353-S357.
- 318 21. Kostka SJ. Amelioration of water repellency in highly managed soils and the  
319 enhancement of turfgrass performance through the systematic application of surfactants. J  
320 Hydrol. 2000;231:359-368.
- 321 22. Oostindie K, Dekker LW, Wesseling JG, Ritsema CJ. Soil surfactant stops water  
322 repellency and preferential flow paths. Soil Use and Manage. 2008;24:409-415.
- 323 23. Kostka S, Dekker K, Oostindie CJ, Ritsema CM, Miller D, Karcher E. Advances in  
324 understanding and managing water repellent soils Aquatrols Corporation, Cherry Hill, NJ  
325 2011; Accessed 1 January 2016. Available:  
326 [https://www.researchgatenet/profile/Coen\\_Ritsema/publication/40147559\\_Advances\\_in\\_und](https://www.researchgatenet/profile/Coen_Ritsema/publication/40147559_Advances_in_understanding_and_managing_water_repellent_soils/links/0046353bbdd5fe0e28000000pdf)  
327 [erstanding\\_and\\_managing\\_water\\_repellent\\_soils/links/0046353bbdd5fe0e28000000pdf](https://www.researchgatenet/profile/Coen_Ritsema/publication/40147559_Advances_in_understanding_and_managing_water_repellent_soils/links/0046353bbdd5fe0e28000000pdf).
- 328 24. Madsen MD, Coronel EG, Hopkins BG. Soil surfactant products for improving  
329 hydrologic function in post-fire water-repellent soil. Soil Sci Soc Am J. 2013;77:1825-  
330 1830.
- 331 25. Roy D, Kommalapati RR, Mandava SS, Valsaraj KT, Constant WD. Soil washing  
332 potential of a natural surfactant. Envir Sci Tech. 1997;31:670-675.

- 333 26. Jia Z, Koike K, Nikaido T. Major triterpenoid saponins from *Saponaria officinalis*. J  
334 Nat Prod. 1998;61:1368-1373.
- 335 27. Oleszek W, Hamed A. Saponin-based surfactants. In: Kjellin M, Johansson I editors;  
336 Surfactants from Renewable Resources, West Sussex: John Wiley and Sons Ltd; 2010.
- 337 28. Abu-Zreig M, Rudra RP, Dickinson WT. Effect of application of surfactants on hydraulic  
338 properties of soils. Biosyst Eng. 2003;84:363-372.
- 339 29. Stroosnijder L, Moore D, Alharbi A, Argaman E, Biazin B, van den Elsen E. Improving  
340 water use efficiency in drylands. Curr Opin Env Sustain. 2012;4:497-506.
- 341 30. Lehrs GA, Sojka RE, Reed JL, Henderson RA, Kostka SJ. Surfactant and irrigation  
342 effects on wettable soils: Runoff, erosion, and water retention responses. Hydrol Process.  
343 2011;25:766-777.
- 344 31. Selker JS, Steenhuis TS, Parlange JY. Wetting front instability in homogeneous sandy  
345 soils under continuous infiltration. Soil Sci Soc Am J. 1992;56:1346-1350.
- 346 32. Wallis MG, Horne DJ, McAuliffe KW. A study of water repellency and its amelioration in a  
347 yellow-brown sand: 2. Use of wetting agents and their interaction with some aspects of  
348 irrigation. New Zeal J Agr Res. 1990;33:145-150.
- 349 33. Akhter J, Mahmood K, Malik KA, Mardan A, Ahmad M, Iqbal MM. Effects of hydrogel  
350 amendment on water storage of sandy loam and loam soils and seedling growth of barley,  
351 wheat and chickpea. Plant Soil and Environ. 2004;50:463-469.
- 352 34. Mingorance MD, Gálvez JF, Pena A, Barahona E. Laboratory methodology to approach  
353 soil water transport in the presence of surfactants. Colloids and Surface A. 2007;306:75-82.