

Original Research Article**Intercharacter correlation between budding success
in *Hevea brasiliensis* Muell. Arg. and seven weather
characters****ABSTRACT**

Aim: To evaluate the relationship between budding success and weather factors

Place and Duration of Study: Nursery site of the Rubber Research Institute of Nigeria, Benin City, Nigeria for three years.

Methodology: Seven weather characters were evaluated for correlation with budding success in Rubber Research Institute of Nigeria, Benin City, Nigeria. The seven weather characters were rainfall, relative humidity (RH) at 0900hrs and 1500hrs, minimum and maximum temperature, evaporation and radiation. Budding was carried out in the rootstock nursery and data on budding success was recorded over a period of six months for three consecutive years. Data for the corresponding six months was collated and for three years. Intercharacter correlation was calculated for budding success and the seven weather characters. The t-test was applied to test significance of the correlation coefficients.

Results: There was positive correlation between budding success and relative humidity at 0.74 to 0.82, while correlation between budding success was negatively correlated with evaporation at -0.83 to -0.88. Correlation between budding success and radiation was also negative at -0.77. Each significant weather character was significant in correlation with temperature. In this case, relative humidity was negatively correlated with temperature at -0.79 to -0.99. Correlation between evaporation and temperature was positive at 0.81 to 0.98 and radiation had significant positive correlation with temperature at 0.81 to 0.98.

Conclusion: The significant correlation coefficients between budding success and three weather characters suggest influence of climate change on budding success for production of planting materials of *Hevea brasiliensis*. This is an indication of appropriate location considerations for nursery facilities in order to enhance budding success. Path analysis to detect direct and indirect effects of the

significant weather factors on budding success will be evaluated in further study.

Keywords: Hevea, weather, budding, correlation, nursery, planting materials

1. INTRODUCTION

Hevea brasiliensis is valued for the natural rubber and it is the major source of natural elastomer world wide. It is an economic crop in many tropical countries such as Nigeria, Cote d'Ivoire, Cameroun and Liberia in West Africa, India, Thailand, Malaysia and China in Asia. *H. brasiliensis* produces viable seeds, yet it is propagated as budded material because of the heterozygosity [1]. The rubber tree is an outcrossing plant and hence undergoes segregation to produce seeds. The implication is that the seeds produced will be different from the mother tree genetically. In order to preserve the genetic integrity of the improved genetic materials, budding is practiced.

Climate change is a common phenomenon all over the world and there is noticeable change in weather pattern affecting a number of agricultural practices and hence models such as ecosystem based management and climate smart agriculture have been recommended [2, 3]. Budding success is liable to effect of climate change and this affects the production and availability of planting materials for farmers. This is more relevant as the bulk of farm activities in developing countries is nature dependent. The challenge caused by influence of climate change on budding success is therefore the bases of this study. The objective is to evaluate the response of budding success to six weather characters.

2. MATERIAL AND METHODS

2.1 Study site and data collection

The study was conducted at the nursery of the Rubber Research Institute of Nigeria, Iyanomo, Benin City. Budding was carried out and record of budding success was taken as described by Oghide [4]. Budding was in seven months from April to October each year. There was weather data collection for corresponding months as recorded by the Federal Department of Meteorological Services, Lagos, Nigeria. The weather characters were rainfall, minimum and maximum temperature, relative humidity at 0900hr and 1500hr, evaporation and radiation. The study was carried out in three consecutive years.

2.2 Analysis

Correlation between budding success and each weather character was calculated in each year and for combined data. The T-test of significant correlation was applied.

3. RESULTS AND DISCUSSION

Intercharacter correlation coefficients among budding success and the six weather characters are presented in Tables 1 - 4 for the various years and combined data. There was significant variation for correlation between budding success and relative humidity, evaporation and radiation (Tables 1 and 4). In a study conducted by Omokhame and Emuedo [5], relative humidity was a critical factor affecting latex yield in *Hevea brasiliensis*. There was positive correlation between budding success and relative humidity at 0.74 to 0.82 (Table 1) and this is an indication of high budding success with increasing relative humidity. This justifies concentration of budding activities during the period of high humidity from April to September in many rubber producing countries. Budding during the period of low humidity is often with supplementary water such as irrigation [6].

Table 1. Correlation coefficient of % budding success with weather characters in year I

Weather factor	% Budding Success	Rainfall	Min. Temp.	Max. Temp.	Rh 0900hrs	Rh 1500hrs	Evaporation
Rainfall	0.39						
Min. Temp	-0.63	-0.41					
Max. Temp	-0.69	-0.58	0.69				
Rh 0900 hrs	0.74*	0.49	-0.90*	-0.91*			
Rh 1500 hrs	0.82*	0.66	-0.85*	-0.91*	0.96*		
Evaporation	-0.88*	-0.59	0.87*	0.84*	0.09	-0.96*	
Radiation	-0.77 *	-0.64	0.85*	0.94*	-0.95*	-0.98*	0.93*

*: Correlation coefficients are significant at $P = .05$

Budding success was negatively correlated with evaporation at -0.83 to -0.88 (Tables 1 and 4). There was significant relationship between evaporation and relative humidity at correlation of -0.91 to -0.98 (Tables 1, 2 and 4). This suggests high evaporation will be accompanied by low relative humidity which can lead to desiccation/dryness of the budding union of scion and rootstock and hence low

budding success. This negative impact of evaporation on budding success was reported by Singh et al [7].

Table 2. Correlation coefficient of % budding success with weather characters in year II

Weather factor	% Budding Success	Rainfall	Min. Temp.	Max. Temp.	Rh 0900hrs	Rh 1500hrs	Evaporation
Rainfall	0.02						
Min. Temp	-0.55	0.07					
Max. Temp	-0.23	-0.1	0.91*				
Rh 0900hrs	0.35	0.26	-0.84*	-0.94*			
Rh 1500hrs	0.16	0.31	-0.83*	-0.97*	0.90*		
Evaporation	-0.29	-0.25	0.87*	0.98*	-0.98*	-0.97*	
Radiation	-0.01	-0.1	0.81*	0.98*	-0.90*	-0.95*	0.94*

*: Correlation coefficients are significant at $P = 0.05$

Radiation is a critical primary factor resulting in climate change [8]. According to Hartl-Meier et al [9], increased radiation induces warmer and dryer ecosystem and this can adversely affect plants especially a grafted system. This is evident in this study with significant negative correlation between budding success and radiation at -0.77 (Table 1). In addition, radiation was negatively correlated with relative humidity at -0.90 to -0.99 (Tables 1 - 4). The relationship between radiation and temperature was positive at 0.81 to 0.98 (Tables 1 - 4). The indirect effects of temperature and relative humidity operating through radiation will be further investigated using path analysis as proposed by Omokhame and Emuedo [10].

Table 3. Correlation coefficient of % budding success with weather characters in year III

Weather factor	% Budding Success	Rainfall	Min. Temp.	Max. Temp.	Rh 0900hrs	Rh 1500hrs	Evaporation
Rainfall	-0.12						
Min. Temp	-0.35	-0.25					
Max. Temp	-0.57	0.61	0.78*				
Rh 0900hrs	0.27	-0.65	-0.52	-0.93*			
Rh 1500hrs	0.22	-0.22	-0.79*	-0.98*	0.86*		
Evaporation	-0.71	0.42	0.60	0.81*	-0.76	-0.72	

Radiation	-0.35	0.48	0.59	0.95*	-0.92*	-0.13	0.82*
-----------	-------	------	------	-------	--------	-------	-------

*: Correlation coefficients are significant at $P = 0.05$

Among other weather factors, relative humidity was negatively correlated with temperature at -0.79 to -0.99 (Tables 1 - 4). The combined effects of temperature and relative humidity have been reported to affect tolerance of the rubber plant to climate change [11, 12]. This was further expressed as Temperature-Relative humidity index (THI) described by Mu et al. [13] and Meiyappan et al. [14]. The THI will be applied in subsequent study. Correlation between evaporation and temperature was positive at 0.81 to 0.98 (Tables 1 - 4). Radiation had significant positive correlation with temperature at 0.81 to 0.98 (Tables 1 - 4). The consistent relationship between temperature and the first three weather factors (evaporation, relative humidity and radiation) is a reflection of the position of temperature in climate change. The first concern of climate change is global warming and it is measured as change in temperature [15].

Table 4. Correlation coefficient of % budding success with weather characters across the three years

Weather factor	% Budding Success	Rainfall	Min. Temp.	Max. Temp.	Rh 0900hrs	Rh 1500hrs	Evaporation
Rainfall	0.42						
Min. Temp	-0.69	-0.76*					
Max. Temp	-0.63	-0.69	0.91*				
Rh 0900hrs	0.64	0.64	-0.90*	-0.99*			
Rh 1500hrs	0.59	0.81*	-0.94*	-0.98*	0.96*		
Evaporation	-0.83*	-0.77*	0.96*	0.93*	-0.92*	-0.91*	
Radiation	-0.64	-0.65	0.93*	0.97*	-0.95*	-0.99*	0.89*

*: Correlation coefficients are significant at $P = 0.05$

4. CONCLUSION

There was a significant relationship between budding success and each of evaporation, relative humidity and radiation. These three weather characters had significant correlation with temperature. These results suggest appropriate cultural practices to avoid or minimise the hazards of climate

change on production of planting materials of *Hevea brasiliensis*. This will be enhanced by better understanding of the indirect effects of temperature either through THI or path analysis.

REFERENCES

1. Soman TA, Mydin KK, Jacob, J. Root trainer planting technique for *Hevea* – A review. Rubber Sci., 2013; 26: 175 – 187.
2. Barrow E, Moiseev A, Raza A. Learning framework for IUCN's work on EbA (Ecosystem Based Adaptation). International Union for Conservation of Nature, 2013; 2p.
3. FAO Climate smart agriculture source book. Food and Agricultural Organisation of the United Nations, 2013; 570p.
4. Oghide A. Rubber nursery establishment. Rubber value chain training on nursery establishment, budding, rubber based intercropping, mixed farming, tapping, nursery operators and processor. Rubber Research Institute of Nigeria, Benin City, Nigeria. 2016; 10p.
5. Omokhaye KO, Emuedo OA. Evaluation of influence of five weather characters on latex yield in *Hevea brasiliensis*. International Journal of Agricultural Research, 2006a; 1: 234 – 239.
6. Omokhaye KO, Oghide A, Imoren EA. Investment opportunities in the upstream sector of the natural rubber industry. Third Palm Oil, Rubber and Cocoa Conference, Labadi Beach Hotel, Accra, Ghana. Organised by Centre for Management Technology, Singapore, 2015; 16p.
7. Singh M, Jacob J, Chandrashekar TR, Annamalaiathan K. Performance of rubber plantation raised from bud grafted clones and polycross seeds under different moisture regimes in north Konkan region of Maharashtra. In: RRII (ed.) Climate change and rubber cultivation: R & D Priorities. Rubber Research Institute of India, Kottayam, Kerala, India, 2010; 111p.
8. Stephens GL, Li J, Wild M, Clayson CA, Loeb N, Kato S, L'Ecuyer T, Stackhouse Jr PW, Lebsock M, Andrews T. An update on Earth's energy balance in light of the latest global observations. Nature Geoscience, 2012; 5: 691 – 696.
9. Hartl-Meier C, Dittmar C, Zang C, Rothe A. Mountain forest growth response to climate change in the Northern Limestone Alps. Trees, 2014; 28:819–829.

- 138 10. Omokhafa KO, Emuedo OA. The use of path analysis to facilitate breeding procedure in *Hevea*
139 *brasiliensis*. Thirteenth Annual Conference of Genetics Society of Nigeria, National Centre for
140 Genetic Resources and Biotechnology, Ibadan, Nigeria, 2006; pp. 1 – 3.
- 141 11. Mathew J, Abraham T, Jose VT, Mondal, GC, Raj S, Sailajadevi T. Prevalence of pests and
142 diseases of *Hevea brasiliensis* in India – Past and present. In: RRII (ed.) Climate change and
143 rubber cultivation: R & D Priorities. Rubber Research Institute of India, Kottayam, Kerala, India,
144 2010; 111p.
- 146 12. Roy CB, Sailajadevi T, Raj S, Gogoi NK, Mathew J. Fishing for *Phytophthora* from rubber
147 plantations in Tripura (North East India) – Is climate inimical to the development of
148 *Phytophthora* in NE India? In: RRII (ed.) Climate change and rubber cultivation: R & D
149 Priorities. Rubber Research Institute of India, Kottayam, Kerala, India, 2010; 111p.
- 151 13. Mu JE, McCarl, BA, Wein, AM. Adaptation to climate change: changes in farmland use and
152 stocking rate in the U.S. Mitigation Adapt. Strat. Global Change, 2013; 18: 6
153 <http://dx.doi.org/10.1007/s11027-012-9384-4>
- 155 14. Meiyappan P, Dalton M, O'Neill BC, Jain AK. Spatial modeling of agricultural land use change
156 at global scale. Ecological Modelling, 2014; 291: 152–174.
- 158 15. Hartmann DL., Klein-Tank AMG, Rusticucci M, Alexander LV, Brönnimann S, Charabi, Y,
159 Dentener FJ, Dlugokencky EJ, Easterling DR, Kaplan A, Soden, BJ, Thorne PW, Wild M, Zhai
160 PM. Observations: Atmosphere and surface, pp. 161 - 254. In: Stocker TF, Qin D, Plattner GK,
161 Tignor M, Allen SK, Boschung J, Nauels A, Xia Y, Bex V, Midgley PM. (eds.) Climate Change:
162 The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of
163 the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge,
164 United Kingdom and New York, NY, USA, 2013.