

**Effects of different storage methods and fertilizer rates on quality of sweetpotato (*Ipomoea batatas* L.) storage roots**

Abstract

Storage of fresh roots of two sweet potato varieties (*Apomuden* and *Okumkom*) for up to 3 months was conducted from January to March, 2011 (minor season) and from August to October, 2012 (major season) using three different storage methods: pit, ash, and grass. Roots were harvested 4 months after planting. Sweet potato was previously amended by treatments of chicken manure (CM) and inorganic fertilizer (NPK). The experimental design was designed in a randomized complete block. The result shows that *Okumkom* applied with 15-30-30 kg/ha NPK + 5t/ha CM and stored in grass gave lowest pest infested roots in both seasons. (Please check again this result, because, *Okumkom* applied with 10 t/ha CM and stored in pit/grass also produce low percentage pest infested roots in 2011 (Fig. 1 b)). *Okumkom* and *Apomuden* stored in ash or grass had lower root weight loss and root sprout during the major season (Please check this result, because, treatments stored in pit also gave to lower weight loss percentage of roots in major season than minor season). *Okumkom* applied with 15-15-15 kg/ha NPK +5t/ha CM and 30 – 45 – 45 kg/ha NPK and stored in grass did not sprout at 12 weeks after storage in both seasons. *Okumkom* applied with 15-30-30 kg/ha NPK + 5t/ha CM and 15-15-15 kg/ha NPK +5t/ha CM and stored in grass produced the least pest infestation, lower root weight and root sprout than *Apomuden*. However, for lower rotten roots, the amended roots should be stored in pit than in grass or ash.

**Keywords:** Sweet potato, pit storage, grass storage, ash storage, *Apomuden*, *Okumkom*

Please consider a alternated abstract as follows:

**[One of the problems in utilization of sweet potato roots in Ghana is how to ensure the quality of post-harvest product. The 3-month experiments was conducted after two harvests to study effects of three different of storage methods on quality of sweet potato (*Ipomoea batatas* L.). Roots were obtained from previous culture experiments with two sweet potato varieties (*Okumkom* and *Apomuden*) and eight fertilizer combinations.**

**The results show that:** (In this part, you should clarify 5 problems)

1. Effect of season on roots quality (is there any difference between major crop and minor crop, please explain base on your weather data?)
2. Effect of varieties (is there any different effects between *Apomuden* and *Okumkom*?)
3. Effect of different fertilizer combinations
4. Effect of 3 methods of storage.
5. Is there any interaction effects between above factors (please choose 1 or 2 suitable storage methods together with fertilizer combinations for each variety which gave to significant results)]

Please note that your experiment can be seen as a 3 factor design which are repeated for 2 seasons. To answer above questions, it would be better to include statistic results in the Section 3 (Results and Discussion), including ANOVA table or multiple tests for each factor, and interaction combination, for example:

| Factors        | df<br>(degree of freedom) | F | Prob |
|----------------|---------------------------|---|------|
| Year           | 1                         |   |      |
| Variety (V)    | 1                         |   |      |
| Fertilizer (F) | 7                         |   |      |
| Storage (S)    | 2                         |   |      |
| V x F          | ...                       |   |      |
| F x S          | ...                       |   |      |
| V x S          | ...                       |   |      |
| V x F x S      | ...                       |   |      |

## INTRODUCTION

Sweet potato (*Ipomoea batatas* L.) is a perennial plant. It is a member of the Convolvulaceae family. Flowers can be white or purple, and leaves can be green or purple. The flesh can be white, cream, yellow, orange, or purple [1] with orange, white, and cream being the most commonly grown and eaten [2]. Sweet potatoes grow well in tropical, subtropical, and temperate areas. In all, sub-Saharan Africa produces only 6% of the world's sweet potato crop [3]. Past and current production trends shows that sweet potato output in developing countries are increasing. Sweet potato output in Africa is estimated to be growing at about five percent per year [4]. Sweet potato is one of the most important root crops in tropical Africa with domestic and industrial uses. In many parts of Africa, including Ghana, the crop is grown for its staple properties. The root is eaten boiled or as fried chips. It is also prepared into flour for various domestic uses and drinking juice can also be extracted from the root [5]. Despite nutritional importance of sweet potato, the roots have a short storage life, generally less than four weeks in the tropics. In Ghana, the production of sweet potato is concentrated almost entirely in the Northern, Upper East and Upper West regions where it is sometimes included in traditional food crop rotations. The crop is also frequently cultivated in the northern part of the Volta region, Ada-Sege areas in Greater Accra region and Komenda –Pepeoso areas in the central region [3]. As a fast growing root crop that can be grown in all regions of the country, giving yields of 20 to 25 tons per hectare within four months, sweet potato has become an important food crop in Ghana. It is also gaining importance as an export crop in the Bawku East District of the Upper East Region. Farmers in the district export the crop to Burkina Faso, where good prices are obtained [5]. Sweet potatoes are a nutritious food, low in fat and protein, but rich in carbohydrate. Roots and leaves are good sources of antioxidants [6] fiber, zinc, potassium, sodium, manganese, calcium, magnesium, iron, and vitamin C [7]. Orange-fleshed sweet potatoes (OFSP) are also very good sources of vitamin A [6, 8]. Because of their high carotenoid content and good yields, OFSP have also been used in several small-scale studies to increase vitamin A status [9, 10, 11, 12]. One of the serious constraints in large scale production and utilization of sweet potato in Ghana is the short life of the harvested root. The crop is highly perishable and as such requires good storage technique. Lack of suitable storage facilities among small holder farmers continues to expose farmers to intermittent food shortages even after harvesting of the crop. Studies indicate that post- harvest losses due to pests and diseases attack can account for as much as 40-60% of crop output. Their

skin is easily damaged during harvest and post-harvest handling leaving the crop highly perishable [13]. Some of the pre and post-harvest problems associated with sweet potatoes cultivation are insect damage of leaves and roots on the field, dehydration and rotting of the roots, high moisture content, lack of storage skills, sprouting and chilling injury during low temperature storage [14]. Cured sweet potato can be stored for 4–7 months at 12.8–15.6°C and 85–90% relative humidity. Non-cured sweet potatoes do not store well and have a much shorter shelf life. The post-harvest physiological processes that may affect storability include; respiration, evaporation of water from the product, sprouting, changes in chemical composition, diseases and damage by extreme temperatures. The relative importance of these processes differs with the storage environment. Farmers rely on preservation methods derived from indigenous knowledge systems for storing the harvested sweet potato roots. Important techniques of preserving crops derived from indigenous knowledge have rarely been subjected to scientific enquiry. There is knowledge gap in the science of these important indigenous techniques of preserving crops. This study is premised on the observation that local small holder farmers in Ghana use most preservation methods informally but not much is known about their efficacy from a scientific perspective. This study will also examine the different sweet potato storage technique to compare their efficacies and comparative effects on root quality. Information gleaned from these studies may help to fine-tune the indigenous technologies and popularize their use for improvement in sweet potato production and preservation.

## MATERIALS AND METHODS

### Sample Source

A previous culture experiment was conducted in randomized complete block design (RCBD) with two sweet potato varieties (*Okumkom* and *Apomuden*) ~~were grown under~~ and eight fertilizer combinations treatments [(i) 10 t/ha chicken manure (CM), (ii) 30-30-30 kg/ha NPK, (iii) 15-15-15 kg/ha NPK + 5 t/ha CM, (iv) 30-45-45 kg/ha NPK, (v) 15-23-23 kg/ha NPK + 5 t/ha CM, (vi) 30-60-60 kg/ha NPK, (vii) 15-30-30 kg/ha NPK + 5 t/ha CM and (viii) No fertilizer (control)]. This experiment was carried out at the Multipurpose nursery research fields of the College of Agriculture Education, University of Education, Winneba, Mampong-Ashanti campus from September, 2011 to January, 2012 (minor season) and from April to July, 2012 (major season).

The experimental design was a 2 x 8 factorial arranged in randomized complete block (RCBD) with four replicates made up of eight organic manure and fertilizer rates and two sweetpotato varieties. The sweet potato roots were harvested at month 4 months after planting. After harvest, the sweet potato roots without pest and disease damage that weigh from 200-250 gram were randomly selected for the next storage experiment. sorted to physically remove pest or disease damaged roots. The sorted roots were then cured naturally in dried under natural sunlight for three days at temperature of by spreading them on the ground ( 25-30 °C, and relative humidity of RH 80-95%.

For storage experiment, the sweet potato roots obtained from culture experiment were stored by three different methods: pit, ash, and grass. There are total of 336 roots (2 varieties x 8 fertilizer combinations x 3 storage methods x 7 roots) for each replicate. The sweet potatoes from each treatment were divided into three portions. For each storage condition or method (pit, ash and grass) the roots were divided into sixteen treatments, each containing seven roots. The experiment was arranged in a randomized complete block with three replications where treatment factors are two varieties, three storage methods and fertilization combinations. A simple random sampling technique was used to select roots from each treatment and replicate. The random sampling approach was to ensure that the parameters under observation are taken from representative universe of units from each replicate and treatment. In both cropping seasons sweet potato roots were stored for three months after each harvest during the The storage duration for minor season and major rainy season was from January to March, 2012 and from August to October, 2012, respectively. Climate data were sourced from Ghana Meteorological Agency Ashanti-Mampong.

## Storage Methods

The various storage methods were set up as follows:

### Pit Storage Method

Roots in a pit 50 x 50 x 50 cm (diameter at top x depth x wide at base) were arranged alternately with grass layers and an upper layer with soil. Pit store was constructed (under a shade to prevent rain water from entering the storage pit) by digging circular pit of 0.5 m diameter at the top, 0.5

~~m depth and 0.5 m wide at the base.~~ The size of the pit was chosen to suit local climatic conditions and is modification of traditional pits. Pit was lined with dry plantain leaves before the sweet potato roots were stored in them. Layers of roots as per treatment were separated with about 1.0 kg dry spear grass (*Imperata cylindrica*). The sweet potatoes roots were finally covered with dry spear grass before covering them with approximately 2.0 kg of soil. The grass acted as an insulating material and ensured cool condition in the pit (17 °C, RH 95-100%).

### Grass Storage Method

Roots in a basket 50 x 50 x 70 cm (base??? x height x wide) were arranged alternately in with grass layers and an upper layer with grass too. ~~A basket measuring 0.5 m at the base, 0.5 m high and 0.7 m wide at the top.~~ The basket was lined with plantain leaves before roots were placed in them. Layers of roots as per treatment were separated with about 1.0 kg dry spear grass (*Imperata cylindrica*). The sweet potatoes were then finally covered with dry spear grass at the top.

### Ash Storage Method

Roots were arranged as the same way with grass storage method, but coated with wood ash. A basket measuring 0.5 m at the base, 0.5 m high and 0.7 m wide at the top was lined with dry plantain leaves before roots were stored in them. Roots as per treatment were thoroughly coated with wood ash by mixing them with 2.0 kg of wood ash and roots were then alternated with 1.0 kg of dry spear grass. The ash acted as an absorbent to moisture and has a repelling effect on pests. Wood ash has alkaline properties, which are not conducive to development of diseases.

In each basket and pit 112 roots were placed and subjected to the experimental conditions. These quantities of roots were obtained from the field experiments after harvest. After setting up the sweet potato under the various storage methods it was monitored for three months and data were collected every two weeks on weight loss in sweet potato (as a % of the total weight stored), sprouting of sweet potatoes (%), spoilage of sweet potatoes (%), root shrinkage and pests and disease infested roots.

## RESULTS AND DISCUSSION

The weather conditions during the experimental periods show that differences in climatic factors (rainfall, temperature and relative humidity) were observed between both cropping seasons (Tables 1 and 2). The total monthly rainfall in the minor rainy season was 429.8 mm and it occurred from September, 2011 to January, 2012 with the peak in September and October (Table 1). The mean monthly temperature of the area for the minor season ranged between 22.2 °C to 32.9 °C with the highest daily of 33.7 °C occurring in January, 2012. The mean monthly relative humidity ranged from 35 to 98 % with the peak occurring between September and November. The bimodal rainfall pattern of Mampong-Ashanti gave the area two seasons; the major season occurred between March and July and the minor from September to November with one month drought spell in August [15]. In the major rainy season (2012), the total monthly rainfall was 948.5 mm and it occurred from April to July, 2012 with the peak in May and June (Table 2). The mean monthly temperature of the site for the major season ranged between 21.9° C to 31.4° C, with the highest daily of 33.3° C occurring in April. The mean monthly relative humidity ranged from 61 to 97 % with the peak occurring between April to July [16].

Table 1: Climate data in 2011 minor season

| Month         | Total rainfall (mm) | Mean relative humidity (%) |       | Mean Temperature (°C) |      |
|---------------|---------------------|----------------------------|-------|-----------------------|------|
|               |                     | 06.00                      | 15.00 | Min.                  | Max. |
| September     | 155.6               | 98                         | 73    | 22.3                  | 29.4 |
| October       | 188.1               | 97                         | 67    | 22.3                  | 30.9 |
| November      | 38.0                | 96                         | 54    | 28.0                  | 32.7 |
| December      | 0.0                 | 92                         | 41    | 22.2                  | 32.9 |
| January, 2012 | 48.1                | 84                         | 35    | 20.8                  | 33.7 |
| <b>Total</b>  | <b>429.8</b>        |                            |       |                       |      |

Source: Ghana Meteorological Agency – Mampong Ashanti, 2011

Table 2: Climate data in 2012 major season

| Month | Total Rainfall (mm) | Mean Relative Humidity (%) |       | Mean Temperature (°C) |      |
|-------|---------------------|----------------------------|-------|-----------------------|------|
|       |                     | 06.00                      | 15.00 | Min.                  | Max. |
| April | 119.3               | 96                         | 61    | 23.0                  | 33.3 |
| May   | 270.8               | 96                         | 63    | 23.0                  | 31.4 |
| June  | 379.8               | 97                         | 71    | 22.6                  | 29.5 |
| July  | 178.6               | 96                         | 71    | 21.9                  | 28.2 |
| Total | 948.5               |                            |       |                       |      |

Source: Ghana Meteorological Agency – Mampong Ashanti, 2012

### Percentage pest infested roots under grass, ash and pit storage

*Apomuden* grown under 15-15-15 kg/ha NPK +5t/ha CM and stored under pit and ash had the highest percentage pest infested root at 12 weeks of storage during the minor growing season. Similarly, *Apomuden* grown under 15-23-23 kg/ha NPK +5t/ha CM and stored under pit, ash and grass gave the highest percentage pest infested root at 12 weeks of storage during the major growing season (Fig. 1). The high pest infestation in *Apomuden* grown under amended plots and stored under pit and ash during both growing seasons might be due to varietal differences in terms of high susceptibility of *Apomuden* to weevil attack and their response to different storage conditions. This agrees with Capinera, [17] that *Cylas puncticollis* is a serious pest for susceptible varieties of sweet potato. *Apomuden* and *Okumkom* grown under 30-30-30 kg/ha NPK and 15-15-15 kg/ha NPK +5t/ha CM, respectively and stored under ash did not store well at 12 weeks of storage during the minor growing season. *Apomuden* and *Okumkom* grown under amended and the unamended plots and stored in pit, ash and grass during the major season had higher percent pest infested roots compared with the minor season (Fig. 1). This might be due to



the low total monthly rainfall combined with high relative humidity during the day and relatively low maximum temperature during the storage period during the major season compared with the minor season storage period. CIP [4] indicated that wet and warm conditions increased the likelihood of serious pest infestations.

*Okumkom* grown under 15-30-30 kg/ha NPK + 5t/ha CM plot and stored in grass had the lowest pest infested roots at 12 weeks after storage (WAS) in both seasons (Fig. 1). The good keeping quality in terms of least pest infestation in *Okumkom* might be attributed to differences in cultivars and their response to manure treatment and storage conditions. *Apomuden* grown under amended plots and stored in pit at 12 WAS recorded the highest pest infested roots followed by *Apomuden* under ash storage with the least recorded by *Apomuden* under grass storage at 12 WAS in both seasons. The high pest infestation of *Apomuden* grown under amended plots and stored in pit might be due to the manure treatments on the roots and their response to different storage conditions. Ahn [18] indicated that the effectiveness of different storage treatments is closely linked with the conditions under which the crop was grown. In the course of the study it was observed that under the pit storage condition, apart from sweet potato weevils which were the common pest identified among all the three storage conditions cricket and millipede were also identified, especially during the major season which produced the result obtained.








Figure 1: Percentage weevil infested roots of sweet potato varieties as influenced by storage methods and organic and inorganic fertilizers during (a, b) minor season, 2011 and (c, d) major season, 2012.

### **Percentage rotten roots under grass, ash and pit storage**

*Apomuden* grown under 30-45-45 kg/ha NPK plot and stored in ash and pit had the highest number of rotten roots at 12 WAS during both storage periods and differed significantly from those grown under amended and control plots (Fig. 2). The highest severity of decay in *Apomuden* grown under 30- 45- 45 kg/ha NPK plot and stored in ash and pit compared with other treatments might be due to high weevil damage, especially in pit storage at 12 WAS. However, *Okumkom* grown under amended plots and stored in grass at 12 WAS during the minor season had zero rotten roots (Fig. 2). *Apomuden* and *Okumkom* grown under amended and control plots and stored in grass had higher number of rotten roots followed by storage in ash with the least number recorded in pit storage during the major season than during the minor season. The severity of decay under grass storage condition might be due to the high temperatures that existed during the long-term storage period coupled with high weevil infestation. This result is similar to those found by Mutandwa and Tafara [19] that if quality of the stored crop and weight variation of roots is considered, then use of soil banks is the most effective. According to Tortoe *et al.*, [20] grass based technique involves the use of dry grass to create dry and cool conditions within the storage area which help to avoid the development of fungal diseases that normally thrive under humid and warm conditions.



**Figure 2:** Percentage rotten roots of sweet potato varieties as influenced by storage methods and organic and inorganic fertilizers during (a, b) minor season, 2011 and (c, d) major season, 2012.

### Percentage weight loss of roots under grass, pit and ash storage

There was a significant difference between *Apomuden* and *Okumkom* grown under amended and control plots and stored in ash, grass and pit in weight loss of roots in both seasons (Fig. 3). The significant effect on root weight loss might be due to cultivar differences as influenced by mineral fertilizer and organic manure application during growth combined with the differences in storage conditions. Weight loss of roots for both *Apomuden* and *Okumkom* grown under amended and control plots and stored in ash, grass and pit increased linearly from the beginning of storage in both season. This result is similar to those found by Amoah [21] that weight loss of sweet potato roots stored in an evaporative cooling barn with three different pre-storage treatments (ash, brine and *Lantana camara* extract) increased linearly from the inception of storage. *Okumkom* applied with 15-15-15 kg/ha NPK + 5t/ha CM and stored in ash had the highest root weight loss at 12 WAS in both season. This observation might be due to combined effect of organic and inorganic fertilizers applied during growth coupled with application of ash during storage. This result agrees with those found by Biswas *et al.*, [22], Mutandwa and Tafara [19] that application of ash to sweet potato roots act as an absorbent to moisture resulting in low relative humidity in the storage condition with resultant high water loss through the root skin surface. Generally, amended roots of *Apomuden* stored in ash, grass and pit during the minor season storage at 12 WAS gave higher root weight loss than during the major season storage (Fig. 3). This might be attributed to differences in cultivar, low relative humidity coupled with relatively high temperature during the storage period in the minor season than during the major season storage period. Generally, *Apomuden* had lower dry matter content and therefore more weight loss compared with *Okumkom*. Metabolic activity of sweet potato is temperature dependent for the formation of starch based on the cultivar. Degras [23] reported that there is a significant role of starch content for determination of fresh and dry matter with the range dependent on cultivar for its classification.

High temperature under storage condition are likely to result in high rates of respiration, increased rates of metabolic breakdown which could result in increased levels of root weight loss. Woolfe [24] and Degras [23] reported that during the storage period of sweet potato roots, roots loss weight owing to respiration and transpiration.




Figure 3: Weight loss percentage of sweet potato roots as influenced by two varieties, three storage methods and fertilizer combinations during minor season, 2011 (a, b) and major season, 2012 (c, d)

### Percentage root sprouting under grass, ash and pit storage

*Apomuden* under amended plots except 15-15-15 kg/ha NPK +5t/ha CM and the control and stored in pit gave higher root sprouting than *Okumkom* grown under the same plots and under similar storage condition at 12 WAS in both seasons (Fig. 4). This might be due to differences in cultivar and its response to fertilizer before storage and resultant physiological changes during storage. Sweet potato cultivars differ very markedly in the tendency to sprout during storage. *Apomuden* and *Okumkom* grown under amended and control plots and stored in pit during the major season storage period differed significantly from ash and grass storage at 12 WAS in root sprout (Fig. 4). The relatively low temperature, high relative humidity and relatively high rainfall during the major season storage period compared with the minor season storage period might have resulted in humid and cool storage condition in pit. According to NSPRI [14] low temperature storage can result in sprouting which is one of the post-harvest problems associated with sweet potato cultivated and stored in Nigeria. Slight accumulation of water in the pit due to relatively high rainfall observed during the latter stage of the major season storage period might have also contributed to high root sprout in pit storage. This result is similar to those found by Dandago and Gungula [25] that accumulation of moisture in soil based technique which involves digging of pits at a certain level of inclination could lead to sprouting of roots.

*Okumkom* grown under amendment treatments, especially 15-15-15 kg/ha NPK +5t/ha CM and 30 – 45 – 45 kg/ha NPK and stored in grass did not record root sprout at 12 WAS in both seasons (Fig. 4). This might be due to differences in cultivar and their response to amended treatment during growth combined with the dry grass used during storage of root. Biswas *et al.*, [22] and Setiawati [26] reported that the effectiveness of different storage treatments is closely linked with the conditions under which the crop was grown and that storage conditions are more effective when the roots have matured in dry soils. Similarly, root damage is lower if the roots are covered with humid rice straw mulch, although the roots tend to sprout. Probably the dry grass used to store root reduced or caused no root sprout in grass storage due to the dry storage condition. *Apomuden* and *Okumkom* grown under amended and control plots and stored in pit gave higher root sprout than under ash and grass storage at 12 WAS in both seasons (Fig. 4). Sprouting was initiated in pit storage at two weeks of storage. This might be due to humid grass straw mulch as a result of high relative humidity in pit storage condition substantiating the

finding of Setiawati [26] that root damage is lower if the roots are covered with humid rice straw mulch, although the roots tend to sprout. Amoah *et al.*, [21] also reported that sprouting of sweet potato roots was initiated after 4 weeks of storage.




Figure 4: Percentage sprouted root of sweet potato varieties as influenced by storage methods and organic and inorganic fertilizers during minor season, 2011 (a, b) and major season, 2012 (c, d)



## CONCLUSION

For minimum pest infestation of sweet potato roots, farmers should grow *Okumkom* and apply 15-30-30 kg/ha NPK + 5t/ha CM and store the roots in grass during the minor and the major seasons. Farmers are to store *Okumkom* and *Apomuden* in pit, ash or grass for decreased root weight loss during the major season. For minimum root sprout farmers should store sweet potato roots in ash and grass than in pit during the major season. The high rainfall experienced during the major season storage period combined with low temperature might have resulted in lower root weight loss with high dry matter content. This might have led to the production of high root sprouts in pit during the major season. Farmers are encourage to store *Okumkom* grown under amendment treatments, especially 15-15-15 kg/ha NPK +5t/ha CM and 30 – 45 – 45 kg/ha NPK in grass as roots did not sprout at 12 weeks after storage in both seasons. *Apomuden* grown under amendments should be stored in ash or grass than in pit since the latter produced higher root sprout at 12 WAS in both seasons.

## REFERENCES

1. Bovell-Benjamin AC. Sweet potato: A review of its past, present, and future role in human nutrition. *Adv Food Nutr Res.* 2007; 52:1–59.
2. Sand hill Preservation Center, Sweet potato order.  
[http://www.sandhillpreservation.com/pages/sweet\\_potato\\_catalog.html](http://www.sandhillpreservation.com/pages/sweet_potato_catalog.html); 2010
3. Tweneboah CK. *Vegetables and Spices in West Africa.* Published by Co-Wood Ltd., Accra. 2000; pp 8- 65.
4. CIP. Sweet potato Facts. Leaflet. Lima, Peru, 1996; 2 pp.
5. Crop Research Institute of Council for Scientific and Industrial Research. Ghana. Annual Report. 2003
6. Teow CC, Truong VD, McFeeters RF, Thompson RL, Pecota KV. and Yencho GC. Antioxidant activities, phenolic and beta-carotene contents of sweet potato genotypes with varying flesh colours. *Food Chem.* 2007; 103:829–38.
7. Antia BS, Akpan EJ, Okon PA. and Umoren IU. Nutritive and anti-nutritive evaluation of sweet potatoes (*Ipomoea batatas*) leaves. *Pakistan J Nutr.* 2006; 2:166–8.

8. Wu X, Sun CJ, Yang LH, Zeng G, Liu ZY. and Li YM. Beta-carotene content in sweet potato varieties from China and the effect of preparation on beta-carotene retention in the Yanshu. *Innov Food Sci Emerg Techn.* 2008; 9:581–6.
9. Haskel LMJ, Pandey P, Graham JM, Peerson JM, Shreshtha KA. and Brown KH. Recovery from impaired dark adaptation in night blind pregnant Nepali women who receive small daily doses of vitamin A as amaranth leaves, carrots, goat liver, vitamin A-fortified rice, or retinyl palmitate. *Am J Clin Nutr.* 2005; 81:461–71.
10. Van-Jaarsveld PJ, Faber M, Tanumihardjo SA, Nestel P, Lombard CJ. and Benade AJ. Beta-carotene-rich orange-fleshed sweet potato improves the vitamin A status of primary school children assessed with the modified-relative-dose-response test. *Am J Clin Nutr.* 2005; 81:1080–1087.
11. Low JW, Arimond M, Osman N, Cunguara B, Zano F, Tschirley DA. Ensuring the supply of and creating demand for a biofortified crop with a visible trait: lessons learned from the introduction of orange-fleshed sweet potato in drought-prone areas of Mozambique. *Food Nutrition Bulletin.* 2007a; 28:S258–70.
12. Low JW, Arimond M, Osman N, Cunguara B, Zano F, Tschirley DA. A food-based approach introducing orange-fleshed sweet potatoes increased vitamin A intake and serum retinol concentrations in young children in rural Mozambique. *J Nutr.* 2007b; 137:1320–7
13. IITA. Sweet potato. In sustainable food production in sub-Saharan Africa 1; International Institute of Tropical Agriculture contribution. 1996; pp. 79-83.
14. NSPRI. Utilization of sweet potato. In proceedings of Nigerian Stored Products Research Institute (NSPRI) workshop in participatory technology development methods. Community analysis report on postharvest practices on cassava and sweet potato, Udo-Ekong CR. 2002; pp. 11-12.
15. Ghana Meteorological Agency-Mampong –Ashanti (2011)
16. Ghana Meteorological Agency, Mampong –Ashanti (2012)
17. Capinera LJ. Sweetpotato weevil (*Cylas formicarius (Fabricius)*). Publication No. EENY-27. University of Florida, 1998; pp 5.

18. Ahn PM. Tropical soils and fertilizer use, Intermediate. Trop. Agric Series, Longman Sci. and Tech. Ltd. UK. 1993.
19. Mutandwa E. and Tafara GC. Comparative assessment of indigenous methods of sweet potato preservation among smallholder farmers: case of grass, ash and soil based approaches in zimbabwe." african studies quarterly. 2007; 9, no. 3
20. Tortoe C, Obodai M. and Amoa-Awua W. Microbial deterioration of white variety sweet potato (*Ipomoea batatas*) under different storage structures. International Journal of Plant Biology. CSIR- Food Research Institute, Accra, Ghana. 2010
21. Amoah RS, Teye E, Abano EE. and Tetteh JP. Effect of pre-storage treatment on the shelf-life of TIS 2 sweetpotato variety. Journal of Agriculture and Biological science vo.2011; 6 (4), 9-12.
22. Biswas J, Sen H. and Mukhopadya Y. Effect of time of planting on root development of sweet potato (*Ipomoea batatas* L. Lam.) Journal of root crops 1988; 14 (1) 11-15
23. Degras L. Sweet potato: The tropical Agriculturalist. Macmillan publishers Ltd. Lima Peru. 2003
24. Woolfe JA. Sweet potato: an untapped food resource. Cambridge University Press, Cambridge, UK, 1992; 643 pp.
25. Dandago MA. and Gungula DT. Effects of various storage methods on the quality and nutritional composition of sweet potato (*Ipomoea batatas* L.) in Yola Nigeria. International Food Research Journal, 2011; 18: 271-278
26. Setiawati Y, Sudaryono A. and Setyono P. Studipenyim pananu bijalarsegar. Dalam: Seminar penerapante knologiproduksidan pascapanenu bijalaruntukmen dukung Agro industriubijalar. Edisikhusus Balitkabi. 1994; Pp. 100-109.