

**Lead induced oxidative stress and development change on
*Coriandrum Sativum***

Summary

Lead (Pb) is ubiquitous pollutant in environment which causes many toxic effects, at high and low dose, Lead makes disruptions at all plants by reduction its growth and development of aerial parts and roots, however in *Coriandrum* the mechanism it not well known. In this study, assessing the capacity of lead to induce a dysfunctioning in development of roots and aerial parts at different concentrations after 21 day and 40 days, biochemical parameters of oxidative stress and morphological change on *Coriandrum sativum* was studied in the present investigation. The seeds were sprayed with a solution containing lead to different concentrations for 40 days with six lead levels (0 as control, 500, 1000, 1500, 2000, 2500 and 3000 mg.l⁻¹). The soil used was characterized and parameters such as growth, oxidative damage markers (lipid peroxidation, proline, chlorophyll and hydrogen peroxide contents) were investigated. The results show that there is no change in properties of roots, sheets and stalks indicate that chlorophyll and carotenoid concentration were significantly decreased at 3000 mg.l⁻¹ Pb than control. Lipid peroxidation and H₂O₂ levels were increased at the dose of 2500 and 3000 mg l⁻¹ compared to control treatment; no difference was noted between 500mg.l⁻¹ and control in all part in plant. Morphological studies show that the group exposed to 3000 mg.l⁻¹ of lead shows a very important development of the tissues of roots and stems compared to control and to the group exposed to 500 mg.l⁻¹ of Pb. changes in morphological structure and non-enzymatic antioxidants have shown that lead exposure causes a significant perturbation on *Coriandrum sativum* Linn seedlings affecting biochemical and physiological processes.

Key words: *Coriandrum sativum*, chlorophyll, Proline, lead, lipid peroxidation.

Introduction

The contamination of soil by heavy metals is one of the most serious environmental problems and has significant implications for human health. Some industrial activities and agricultural practices increase their level in the substrate, and the possible introduction of the elements in the food chain is an increasing human health concern (Cakmak et al., 2000). The accumulation of heavy metals in plants presents a toxic hazard to man, because the cultivated plants are the point of entry into the food chain. Coriander (*Coriandrum sativum* L.), which belongs to the family Apiaceae (Umbelliferae) is mainly cultivated from its seeds throughout the year (Mhemdi et al., 2011), is widely used in the human diet, changing their composition or properties would cause deleterious effects to their consumers.

Engineering industrial techniques used to clean up contaminated soils are expensive and suitable only for small polluted areas (Lutts et al., 2004). Furthermore, these technologies are not only costly, but they also cause soil disturbances and they are not readily accepted by the general public (Perchet, 2008; Saifullah et al., 2009). Phytoremediation has been highlighted as an alternative technique to traditional methodologies, for the removal of heavy metals from soil. Two approaches have been generally proposed for the phytoremediation of heavy metals. The first one is using of natural hyperaccumulator plants with high metal accumulating capacity. However, the research must be pursued at the level of the increase of the assimilation of heavy metals by the vegetation we taking as example the lead. While the second is utilizing of high-biomass plants whereas, relatively high amounts of the metal accumulated by plants is often translocated from the root to the more easily harvestable shoots (Chen et al., 2004 and Manousaki et al., 2007).

Quantities of lead absorbed by the roots depend on the concentration of metal in the soil, but also of its migration capabilities from the ground to the roots and the amount of lead present in the various part of plants depends on the transport from outside of roots toward the inside, and then its translocation to leaves (Patra et al., 2004). Lead contamination in the plant environment is known to cause highly toxic effects on processes such as depression on seed germination (Wierzbicka and Obidzinska, 1998), toxicity of nucleoli (Liu et al., 1994), inhibition of root and shoot growth (Liu et al., 1994), reduction in photosynthesis (Poskuta and Wacławczyk-Lach, 1995), DNA synthesis (Gabara et

al., 1992) and inhibition or activation of enzymatic activities (Van Assche and Cliisters, 1990). Lead not only affects plant growth and productivity but also enters into the food chain causing health hazards to man and animals (Seaward and Richardson, 1990).

Lead was recognized as causing oxidative stress in plants; coriander may have a strong resistance to lead, can have an antioxidant activity and inhibit unwanted oxidant processes (Wangensteen et al., 2004; Melo et al., 2005). But little is known about the effects of lead on the physiological processes and the biochemical changes at short and long term of exposed to lead of coriander seedlings. The objective of the present work is to study the effects of different concentrations of lead on leaf, roots and stems of coriander including growth, physiological and biochemical processes such as the different pigments, soluble proteins, lipid peroxidation and hydrogen peroxide contents; and description of morphological changes. The possible mechanisms of Coriander seedlings tolerance of lead stress are briefly discussed in the present study.

27 **Materials and Methods**

28 *Plant material*

29 Coriander (*Coriandrum sativum* L.) seed were used in our experimentation and surface was sterilized
30 with 0.1% HgCl₂ for the prevention of fungal and bacterial contamination (Young, 1926). In the first,
31 we have tempered seeds overnight for accelerate germination. We conducted the experiment in 14
32 clean plastic pots which has a depth of 20 cm and a diameter of less than 10 cm perforated to allow
33 the water to drain out. Each pot was filled with 500 g sandy soil, containing compost from vegetable
34 origin added to the sterilized sand (2V of sand / 1V of compost), soil sample was characterized by pH
35 = 7.8, E_{Ce} = 2.5 dS/m, CaCO₃ = 3.2 %, organic matter = 0.02 %, Clay = 2 %, Silt = 3 % and Sand =
36 95 %. We have sowed our seeds (70 seeds) to approximately 2.5 cm in depth and pots were watered
37 to keep moisture content approximately at 60% of water holding capacity during 40 days with solutions
38 of lead acetate at different concentrations (500,1000,1500 , 2000, 2500, 3000 mg.l⁻¹) and a witness
39 sprayed with distilled water, with 3 replicates. Our seedling was placed in a greenhouse in Oran 1
40 University in controlled conditions as it is shown on Table 1. After 40 days of lead exposure, the fresh
41 sample weights was determined and were kept at -80°C for further analyses.

42 *Chlorophyll and carotenoid determination*

43 Fresh biomass (leaves) was homogenized in 80% icecold acetone in the dark and then centrifuged at
44 10000g for 10 min at 4°C and the supernatant was used for the immediate determination of pigments.
45 Absorbance of the solution was determined spectrophotometrically at 663, 645 and 480 nm the
46 contents of chlorophyll a, b, and carotenoid, respectively; with the following equations help of Arnon's
47 formulae (Arnon, 1949), for quantification of the total chlorophyll, chlorophyll a and chlorophyll b
48 content in an 80% acetone extract:

49 Total chlorophyll = 20.2 (A₆₄₅) + 8.02 (A₆₆₃)

50 Chlorophyll a = 12.7 (A₆₆₃) - 2.69 (A₆₄₅)

51 Chlorophyll b = 22.9 (A₆₄₅) - 4.68 (A₆₆₃)

52 And Carotenoids= (1000A₄₈₀ - 3.27[chl a] - 104[chl b])/227

53 Chlorophyll and carotenoid concentrations were expressed as mg g⁻¹fresh weight.

54 *Estimation of lipid peroxidation*

55 The level of peroxidation was measured in terms of malondialdehyde (MDA) (a product of lipid
56 peroxidation) content determined by the thiobarbituric acid (TBA) reaction as described by Heath and
57 Packer (1968). Frozen shoot was homogenized in 5 ml of 0.1% trichloroacetic acid (TCA). The
58 homogenate was centrifuged at 8000g for 20 min and 4.0 ml of 20% TCA containing 0.5% TBA was
59 added. The mixture was heated at 95°C for 30 min and then quickly cooled on ice bath. The contents
60 were centrifuged at 8000 g for 20 min and the absorbance of the supernatant was measured at 532
61 nm and the value for the non-specific absorption at 600 nm was subtracted. The concentration of MDA
62 was calculated using coefficient of absorbance of 155 mM⁻¹ cm⁻¹. MDA content expressed as nM g⁻¹
63 fresh weight.

64 *Determination of hydrogen peroxide*

65 The H₂O₂ concentration was determined according to Loreto and Velikova (2001). Approximately 0.1g
66 of shoots was homogenized at 4°C in 2 ml of 0.1% trichloroacetic acid (TCA) (w:v). The homogenate
67 was centrifuged at 12000 g for 15 min at 4°C. Then, 0.5 ml of the supernatant was added to 0.5 ml of
68 10 mM K-phosphate buffer (pH 7.0) and 1 ml of 1M KI. The H₂O₂ concentration of the supernatant was
69 evaluated by comparing its absorbance at 390 nm with a standard calibration curve. Hydrogen
70 peroxide concentration was expressed as $\mu\text{M.g}^{-1}$ fresh weight.

71 *Extraction and assays of soluble proteins*

72 The proteins have been doses according to the method (Bradford, 1976), briefly 1 g of fresh tissue,
73 was crushed in the presence of sand, in 10 ml of medium of extraction with following composition:
74 phosphate buffer (0.1M, pH 7); K₂HPO₄ to 0.1M; Triton x 100 to 0.1 % ; EDTA and centrifuged at 3000
75 rpm for 10 min, the supernatant (protein extract) is recovered for the assay of protein and reaction was
76 read to 595 nm and express by $\mu\text{g MF}$ (fresh material).

77 *Proline assay*

78 400 Mg of plant material is put into a mortar and chopped with 5 ml of ethanol at 95% followed by
79 three flushing and washes with 5ml of ethanol at 70 %. The final solution is collected in a test tube so
80 that it is decanted during 60 min. two phase are distinguished (a higher phase of light green color and
81 a lower phase of dark green color) (Nguyen and Paquin , 1971). 5 mL of the upper phase are
82 collected, to which are added 2 ml of chloroform and 3 ml of distilled water. After agitation, the solution
83 is maintained at rest for 24 hours in the cold for a good separation. After assay the optical density is
84 read using a colorimeter, the wavelength is 515 nm.

85 86 *Data analysis*

87 All data were analyzed in three replications and the obtained data were evaluated statistically using
88 Student's test, and least significant difference (LSD) was calculated at $p < 0.05$.

89 90 **Results**

91 92 *Growth and fresh weight*

93 After exposure of seeds to different concentration of lead (from 500 to 3000 mg.l^{-1}) for 20 days period
94 (see table 2 & 3); there was a significant reduction ($p < 0.05$) in leaves numbers, by 40%, 58%, 70%;
95 74%, 84% and 92% respectively, compared to control, and the weight of total sheets, rods and roots
96 decrease significantly after poisoned in all groups vs. control by 90%, 87%, 81%, respectively
97 ($p < 0.05$). After 40th days of exposure to different concentration of lead a net increase in weight of rods
98 and sheets was noted than control. The numbers of sheets was significantly decreasing with
99 increasing the concentration from 500 to 3000 mg.l^{-1} , respectively than control.

100 *Chlorophyll and carotenoid*

101 After 20th days, our results shown a significant decrease ($p < 0.05$) in the rate of chlorophyll (a) in all
102 groups exposed to lead (from 500 mg.l^{-1} to 3000 mg.l^{-1}) compared to control group by 10.41%,
103 24.94%, 43.27%, 47.50% ,65.5% and 76.46%. Chlorophyll (b) have significantly lower levels than
104 control groups by 23.95 %, 55.98 %, 51.39%, 70.89%, 72.56% and 84.95%, respectively. Increased
105 lead exposure causes a significant reduction in carotene levels compared to control from 49.08 % to
106 94.56 % respectively (Table. 4).

107 After 40th days, the rate of carotenoid and total chlorophyll decreased significantly at the dose of 2500
108 mg.l^{-1} and 3000 mg.l^{-1} compared to the control; mean values are significantly reduced ($p < 0.05$) by -
109 27%, -41% and -24%, -34%, respectively, and no difference was noted in the other concentrations.

110 *Lipid peroxidation and hydrogen peroxide*

111 At 20th days of exposure to increased levels of lead (500, 1000, 1500, 2000, 2500 and 3000 mg.l^{-1}),
112 we noted a significant increase ($p < 0.05$) level of hydrogen peroxide in sheets compared to control. In

rods, the level value of hydrogen peroxide was higher ($p<0.05$) by 2 to 4 times in all groups exposed to lead compared to control. According our results (fig. 1), we noted that the hydrogen peroxide content was very lower in roots of control, however in contrast the level of hydrogen peroxide was 60 times more important ($p<0.05$) in stems at 3000 mg.l⁻¹ compared to control. After 40 days, the level of hydrogen peroxide is significantly higher (400 times) in the sheets of plant exposed to 3000 mg.l⁻¹ of lead compared to control.

Fig. 1, shows a significantly increased ($p<0.05$) content of lipid peroxidation by 1.5 to 2.5 compared to control in sheets after 20 day of plant exposed to 2000 mg.l⁻¹ to 3000 mg.l⁻¹, in rods the level of lipids peroxidation was increased from 1.8 to 5 times in all groups exposed to different concentration of lead compared to control. It has been noted that the rate of TBARS increases by increasing the lead content, however in roots, after 20 day, we noted that the level of lipids peroxidation was increased by 87.5%, 150%, 250 %, 275% and 300% respectively to a concentration in lead of 1000mg/l, 1500 mg/l, 2000 mg/l, 2500 mg/l and 3000mg/l compared to control ($p<0.05$). After 40 days of exposure to lead, we obtain that the rods and roots indicate a significant increased of level of lipids peroxidation in all groups by 1.2 and 2.7 times compared to control ($p<0.05$); in sheets, we noted the rate of lipid peroxidation decreases significantly with increasing the rate of lead in the ground, from 1.3 to 3.8 times less.

Soluble protein and Proline levels

The results obtained in Table 5 indicated that the level of soluble protein in leaves was significantly increased by 35% and 60% in groups receive 2000 mg.l⁻¹ and 3000 mg.l⁻¹ respectively compared to control, at 500, 1000, 1500 mg/l dose respectively, we noted a significant decrease 12.5% to 50 %. In rods, the rate of soluble protein was 14.6 and 13.6 times more important in plant receive 2500 mg.l⁻¹ and 3000 mg.l⁻¹ compared to control; in roots, soluble protein was 2 to 4 times more increased in groups receive 1000, 1500, 2000, 2500, 3000 mg.l⁻¹ compared to control. In the Table 6, the soluble protein was significantly higher ($p<0.05$) from 28% to 63% in sheets, rods and roots after irrigation of plant with water contain 3000 mg.l⁻¹ of lead.

At level of leaves, rods and roots (Table. 5 and 6), it was observed that the level of proline increase gradually with increase of lead concentration from 10 to 16 times more compared to control. After 40 days of exposition to lead at different concentration, level of proline was significantly higher from 2 to 8 times more than control, in sheets, rods and roots.

Discussion

Lead is an element not very mobile and has a very high persistence in soil (Saddler and Berthelin, 1998). The plants have many systems of detoxification to limit interaction of these ions with biological molecules; lead induce a range of deleterious effects for the agencies, he disrupts the membrane structures and parietal, modified the statute of waterborne, disrupted the absorption and/or the translocation of essential minerals elements (calcium, manganese, zinc, iron etc....) or further reduced photosynthesis (Seregin and Ivanov, 2001; Sharma and Dubey, 2005). Depending to environment conditions, plants can absorb a part of the lead present in the soil and Pb²⁺ ions broadcast in the root, but are blocked by the physical barrier like endoderme. At the macroscopic scale, the plants exposure to lead induces a reduction in growth of biomass, yields and when stress is too severe, led to the emergence of resprouts necrosis and foliar, chlorotic, or even to the death of the plant (Sharma and Dubey, 2005). It also presents a high affinity for proteins which possess thiol groups or of metal cofactors (metallo-enzymes). However, despite the toxicity certain of these two types of interactions, it is not enough to explain the large variety of deleterious effects observed in plants treated with lead and particularly the genotoxicity induced by this metal (pourrut et al., 2008).

In our results indicate that the treatment with lead induces an increase in production of H₂O₂ and O₂^{•-}, we can explain this by fact that it is possible to estimate indirectly the level of ERO production and the generation of oxidative stress, by measuring the activity of antioxidant enzymes, or the rate of lipid peroxidation, which are biomarkers of oxidative stress (Sharma and Dubey, 2005; Chen et al., 2007; Wang and al., 2007). An increase in lipid peroxidation content in coriander grown under Pb stress was observed (Fig. 1). It is possible that increase of MDA concentration in Coriander may be due to a increase in polyunsaturated fatty acid concentration relative to saturated fatty acids, which has also

been reported in some plants under stressful conditions (Rucinska and Gwozdz, 2005; Grappa et al., 2007).

The increased of H₂O₂ may be essentially due to reduction of catalase activity, the case of this enzyme is complex, since half of publications reported an inhibition of activity by lead (Verma and Dubey, 2003; Choudhury and Panda, 2004; Seregin and al., 2004; Chen and al., 2007; Dey and al., 2007; Hu and al., 2007; Qureshi and al., 2007; Gopal and Rizvi, 2008), the other half shows an increase of his activity (Reddy and al., 2005; Mishra and al., 2006). This inhibition seems to be due to the species treated, to duration or to intensity of the treatment. The inhibition of catalase activity has been observed during different stress, for all the organisms studied, this inhibition is not a general rule; the origin of this phenomenon is still widely debated. In addition, when the CAT is inhibited, this mechanism is dose-dependent and the catalase is a metallo-enzyme, its inhibition could be due to a direct interaction with the lead (Landberg and Greger, 2002).

Some molecules non-loaded are also able to spread through the plant walls and can thus cause of multiple damage as oxidation of DNA (Britt, 1997), or even the oxidation of proteins including at the level of the cysteines and methionines (Rinalducci et al. 2008; Bartoli et al., 2004), these are the two amino acids that are most sensitive and since they are quite often involved in the fixing of metals or in the catalytic properties of many enzymes and proteins; reactive oxygen species inactivate enzymes and damage important cellular components. ROS are responsible for protein, lipid and nucleic acid modification and are thought to play a major role in ageing and cell death (Jacobson et al., 1996).

The lead exposure leads to a strong inhibition of photosynthesis, the photosynthetic yield, and to a limitation of the rate of assimilation of CO₂. This inhibition is explained by the decrease of the levels of chlorophylls and carotenoids generally found (Kosobrukhov and al., 2004; Gopal and Rizvi, 2008). The chlorophyll b seems more sensitive than chlorophyll a (Kacabova and Natr, 1986 ; Wozny and al., 1995; Vodnik and al., 1999). Singlet oxygen is the first excited electronic state of O₂. Insufficient energy dissipation during photosynthesis can lead to formation of chlorophyll (Chl) triplet state. And the Chl triplet state can react with 3O₂ to give up very reactive singlet oxygen. It has been proved that singlet oxygen formation during photosynthesis can have damaging effect on photosystem I (PSI) and a photosystem II (PSII) and on whole machinery of photosynthesis.

Lead interacts at this level in two ways:

- In a direct manner, by substituting divalent ions related to the metallo-enzymes. This is particularly the case with the δ-aminolevulinate deshydratase (ALAD) which is at the basis of the synthesis of the chlorophylls and whose ion Zn²⁺ is replaced by Pb²⁺;

- In an indirect way, by inducing a deficiency in these divalent ions.

Proline, α-amino acid is an antioxidant and potential inhibitor of programmed cell death and is considered as an indicator of stress; in several plant, the accumulation of proline has been observed as a response to biotic and abiotic stress ((Boguszewska et al. 2010; Torres 2010; Khatamipour and al., 2011). It has been suggested that free proline act as osmoprotectant, a protein stabilizer, a metal chelator, an inhibitor of lipid peroxidation and OH• and ¹O₂ scavenger. Increased proline accumulation appears especially during salt, drought and metal stresses (Trovato et al., 2008). Therefore proline is not only an important signaling molecule, but also an effective ROS quencher. It has been found that the important role of proline is in potentiating pentose-phosphatase pathway activity as important component of antioxidative defense mechanism (Hare & Cress 1997).

Conclusion

Thus, the finding that *Coriandrum sativum* L. shows that it is good plant material for studying other aspects of abiotic stress resistance mechanisms. Based on the present work, it can be suggested that toxic concentrations of lead cause oxidative stress, as evidenced by increased H₂O₂ formation, lipid peroxidation and proline content in coriander plant. In this study, a significant change in different parameters such as growth of shoot and roots, chlorophyll and carotenoid concentrations coupled with lipid peroxidation, protein oxidation and hydrogen peroxide indicated that high Pb levels in nutrient

solution produced toxic effects. It was proposed that the reduced growth in Pb of coriander exposed to toxic levels of Pb might be induced by an enhanced production of toxic oxygen species and subsequent lipid peroxidation. Moreover, it was possible to observe that Pb-tolerant plants developed some defense mechanisms against oxidative stress.

References

- Arnon D.I., (1949). Copper enzymes in isolated chloroplasts: polyphenoloxidase in *Beta vulgaris*. *Plant Physiol* 24: 1-5
- Bartoli C., Gómez F., Martinez D.E., Guamet J.J., (2004). Mitochondria are the main target for oxidative damage in leaves of wheat (*Triticum aestivum* L). *Journal of Experimental Botany*, 55, pp. 1663-1669.
- Boguszewska D., Grudkowska M., Zagdańska B., (2010). Drought responsive antioxidant enzymes in potato (*Solanum tuberosum* L.), *Potato Research*, 53, pp. 373-382.
- Bradford M.M., (1976). A rapid and sensitive method for the quantification of micro-gram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry* 72: 248-254.
- Briat J.F., Lebrun M., (1999). Plant responses to metal toxicity. *Plant Biology and Pathology*, Academy of Sciences, Elsevier, Paris, 322: 43-54.
- Britt A.B., (1997). Genetic analysis of DNA repair in plants. In Lumsden P Mpoloka 4881 (ed.). *Plants and UV-B : Responses to environmental change*, Cambridge University Press. pp 77-93.
- Cakmak I., (2002) Plant nutrition research: Priorities to meet human needs for food in sustainable ways. *Plant Soil* 247, 3–24.
- Chen J., Zhu C., Li L., Sun Z, Pan X., (2007). Effects of exogenous salicylic acid on growth and H₂O₂-metabolizing enzymes in rice seedlings under lead stress. *J Environ Sci (China)* 19(1):44–49.
- Choudhury S., Panda S., (2004). Toxic effects, oxidative stress and ultrastructural changes in moss *Taxithelium Nepalense* (Schwaegr.) Broth. under chromium and lead phytotoxicity. *Water Air Soil Pollut* 167(1):73–90.
- Dey S.K., Dey J., Patra S., Pothal D., (2007). Changes in the antioxidative enzyme activities and lipid peroxidation in wheat seedlings exposed to cadmium and lead stress. *Braz J Plant Physiol* 19(1):53–60.
- Gopal R, Rizvi A.H., (2008). Excess lead alters growth, metabolism and translocation of certain nutrients in radish. *Chemosphere* 70(9):1539–1544.
- Groppa M.D., Tomaro M.L., Benavides M.P., (2007). Polyamines and heavy metal stress: the antioxidant behavior of spermine in cadmium- and copper-treated wheat leaves. *Biometals* 20:185–195
- Hare P.D., Cress W.A., (1998). Metabolic implications of stress-induced proline accumulation in plants, *Plant Growth Regul.*, 21, pp.79–102.
- Heath R.I., Packer L., (1968). Photoperoxidation in isolated chloroplasts. I-Kinetics and stoichiometry of fatty acid peroxidation. *Arch Biochem Biophysics* 125:189-198.
- Heath RI, Packer L., (1968). Photoperoxidation in isolated chloroplasts. I-Kinetics and stoichiometry of fatty acid peroxidation. *Arch Biochem Biophysics* 125:189-198.

253 Hu J, Shi G, Xu Q, Wang X, Yuan Q, Du K., (2007). Effects of Pb²⁺ on the active oxygenscavenging
 254 enzyme activities and ultrastructure in *Potamogeton crispus* leaves. *Russ J Plant Physl* 54(3):414–
 255 419.

256 Jacobson M.D., Weil M., Raff M.C., (1996). Role of Ced-3/l family proteases in stuarosporine-induced
 257 programmed cell death, *J. Cell Biol.* 133, pp.1041–1051.

258 Kacabova P and Natr L. (1986). Effect of lead on growth characteristics and chlorophyll content in
 259 barley seedlings. *Photosynthetica* 20(4): 411-417.

260 Khatamipour M., E. Piri, Y. Esmaeilian, Tavassoli A., (2011). *Scholars Research Library Annals of*
 261 *Biological Research*, 2 (5) :527-532

262 Kosobrukhov H., I Knyazeva, V. Mudrik (2004). *Plantago major* plants responses to increase content
 263 of lead in soil: Growth and photosynthesis. *Plant Growth Regulation* 42: 145-151.

264 Landberg T and M Team (2002). Differences in: oxidative stress in heavy metal resistant and
 265 sensitive clones of *Salix viminalis*. *Journal of Plant Physiology* 159 (1): 69-75.

266 Loreto F. and Velikova V. 2001. Isoprene produced by leaves protects the photosynthetic apparatus
 267 against ozone damage, quenches ozone products, and reduces lipid peroxidation of cellular
 268 membranes. *PlantPhysiol* 127: 1781-1787

269 Lutts, S., I. Lefevre, Ch. Delperee, S. Kivits, C. Dechamps, A. Robledo, E. Correal, (2004). Heavy
 270 metal accumulation by the halophyte species Mediterranean saltbush. *J. Environ. Qual.*, 33: 1271-
 271 1279.

272 Malecka H., Jarmuszkiewicz W., Tomaszewska B., (2001). Antioxidative defense to lead stress in
 273 subcellular compartments of pea root cells. *ActaBiochimicaPolonica* 48 (3): 687-98.

274 Melo EA, Filhob JM, Guerrac NB: Characterizations of antioxidant compounds in aqueous coriander
 275 extract (*Coriandrum sativum* L.). *Lebensm.-Wiss u.-Technol* 2005; 38:15-19.

276 Mhemdi H, Rodier E, Kechaou N, Fages J., (2011). A supercritical tuneable process for the selective
 277 extraction of fats and essential oil from coriander seeds. *J. Food Engg.* 105(4):609-616

278 Mishra S, Srivastava S, Tripathi R, Kumar R, Seth C, Gupta D., (2006). Lead detoxification by coontail
 279 (*Ceratophyllum demersum* L.) involves induction of phytochelatins and antioxidant system in response
 280 to its accumulation. *Chemosphere* 65(6):1027–1039.

281 Mishra W., Choudhuri MY., (1999). Effects of Salicylic Acid on Heavy Metal-Induced Membrane
 282 Deterioration Mediated by Lipxygenase in Rice. *Biologia Plantarum* 42 (3): 409-415.

283 Nguyen S.T., Paquin R., (1971). Methodes d'extraction et de purification des acides amines libras et
 284 des proteines de tissus vegetaux. *J. Chromatography*, 61, 349-351

285 Parys E., Romanowska E., Siedlecka M., Poskuta JW., (1998). The effects of lead on photosynthesis
 286 and respiration in detached leaves and mesophyll protoplasts of
 287 *Pisumsativum*. *ActaPhysiologiaePlantarum* 20: 313-322.

288 Patra M., Bhowmick N., Bandopadhyay B., Sharma A., (2004). Comparison of mercury, lead and
 289 arsenic with respect to genotoxic effects on plant systems and the development of genetic tolerance.
 290 *Environmental and Experimental Botany*, 52: 199-223.

291 Perchet, G., Sangely, M., Goñi, M., Merlina, G., Revel, J.C., Pinelli, E., (2008). Microbial population
 292 changes during bioremediation of nitroaromatic- and nitraminecontaminated lagoon. International
 293 Biodeterioration & Biodegradation 61, 304-312.
 294 Pourrut B, Perchet G, Silvestre J, Cecchi M, Guiesse M, Pinelli E., (2008). Potential role of NADPH-
 295 oxidase in early steps of lead-induced oxidative burst in *Vicia faba* roots. J Plant Physiol 165(6):571–
 296 579.
 297 Qureshi M, Abdin M, Qadir S, Iqbal M., (2007). Lead-induced oxidative stress and metabolic
 298 alterations in *Cassia angustifolia* Vahl. Biol Plantarum 51(1):121–128.
 299 Reddy AM, Kumar SG, Jyothsnakumari G, Thimmanaik S, Sudhakar C., (2005). Lead induced
 300 changes in antioxidant metabolism of horsegram (*Macrotyloma uniflorum* (Lam.) Verdc.) and
 301 bengalgram (*Cicer arietinum* L.). Chemosphere 60(1):97–104.
 302 Rinalducci S., Murgiano L., Zolla L., (2008). Redox proteomics: basic principles and future
 303 perspectives for the detection of protein oxidation in plants. J Exp Bot 59:3781– 3801.
 304 Rucinska, R., Gwozdz, E.A., (2005). Influence of lead on membrane permeability and lipoxygenase
 305 activity in lupine roots. Biol. Plant. 49, 617–619.
 306 Saddler P and J Berthelin (1998). Contamination of soils by trace elements: the risks and their
 307 management. Report of the academy of sciences Ed. Lavoisier, Paris, 42 pp.
 308 Saifullah Meers E., Qadir M., de Caritat P., Tack F. M. G., Du Laing G., Zia, M. H. (2009). EDTA-
 309 assisted Pb phytoextraction. Chemosphere. 74, 1279–1291.
 310 Seregin I.V., Ivanov V.B., (2001). Physiological aspects of cadmium and lead toxic effects on higher
 311 plants. Russ J Plant Physiol 48(4):523–544
 312 Seregin I.V., Shpigun L.K., Ivanov V.B. (2004). Distribution and toxic effects of cadmium and lead on
 313 maize roots. Russ J Plant Physiol 51(4):525–533.
 314 Seregin I.V., Shpigun L.K., Ivanov V.B., (2004). Distribution and toxic effects of Cadmium and Lead on
 315 maize roots. Russian Journal of Plant Physiology 51 (4): 525-533.
 316 Sharma P, Dubey R.S., (2005). Lead toxicity in plants. Braz J Plant Physiol 17(1):35–52
 317 Torres, M.A. (2010). ROS in biotic interactions, Physiol. Plant. 138, pp. 414-429.
 318 Trovato M., Mattioli R., Costantino P. (2008). Multiple roles of proline in plant stress tolerance and
 319 development, Rendiconti Lincei, 19, pp. 325-346.
 320 Verma S., Dubey R.S. (2003). Lead toxicity induces lipid peroxidation and alters the activities of
 321 antioxidant enzymes in growing rice seedlings. Plant Science 164 (4): 645-655.
 322 Vodnik D, Jentschke G., Fritz E., Gogala N., Godbold D.L., (1999). Root-applied cytokinin reduces
 323 lead uptake and affects its distribution in norway spruce seedlings. Physiologia Plantarum 106(1): 75-
 324 81.
 325 Wang H, Shan X, Wen B, Owens G, Fang J, Zhang S., (2007). Effect of indole-3-acetic acid on lead
 326 accumulation in maize (*Zea mays* L.) seedlings and the relevant antioxidant response. Environ Exp
 327 Bot 61(3):246–253.
 328 Wangenstein H, Samuelsen AB, Malterud KE: Antioxidant activity in extracts from coriander. Food
 329 Chem 2004; 88:293–97.

Wozny A., Schneider J., Gwozdz E.A., (1995). The effects of lead and kinetin on greening barley leaves. *Biologia Plantarum* 37(4): 541-552.

Young AP (1926). Facultative parasitism and host range of fungi. *Am J Bot* 13:502-520.

Table 1: General condition of culture in the greenhouse.

Parameters	Brightness L	Temperature °C	Air W/M2	Sunshine W/m2	Wind km/h	Humidity %	Side
Day 1	10000	23.9±1.1	474±2.4	468	0	100	Right
Day 2	9915	15.6±1.3*	50±3.2*	51*	0	100	Left
Day 3	9911	15±1.2*	50±2.8*	48*	0	100	Left
Day 4	9663	15±1.1*	49±1.5*	46*	0	100	Left
Day 5	9410	14.8±1.0*	50±2.6*	44*	0	100	Left
Day 6	9174	14.6±1.0*	50±3.3*	43*	0	100	Left
Day 7	10000	24±1.3	474±4.4	466	0	100	Right
Day 8	10000	24.9±1.2	475±4.2	470	0	100	Right
Day 9	8914	14.7±1.2*	50±2.2*	43*	0	100	Left
Day 10	8876	14.3±1.3*	47±2.0*	41*	0	100	Left

(*) the average of three replication are significantly different compared to control at p<0.05

Table 2: Plant morphology modification after 20 days to lead-exposed at different level

Concentration mg/l	Length (cm)	Number of Sheets	Weight (g)		
			Sheets	Rods	Roots
Control	11.5	113	1.64±0.23	1.32±0.26	0.56±0.07
500	7.04	67	0.46±0.01*	1.12±0.11	0.35±0.06*
1000	4.02	47	0.34±0.01*	0.68±0.08*	0.33±0.03*
1500	3.07	33	0.28±0.01*	0.26±0.02*	0.10±0.04*
2000	2.6	29	0.16±0.02*	0.17±0.02*	0.84±0.03*
2500	1.5	17	0.12±0.01*	0.08±0.01**	0.75±0.02*
3000	1.01	9	0.12±0.01*	0.36±0.01*	0.34±0.03*

(*) the average of three replication are significantly different compared to control at p<0.05.

Table 3: Plant morphology modification after 40 days to lead-exposed at different level

Concentration Mg/l	Length Cm	Number of sheets			Weight (g)		
		Green	Clear Green	Yellow	Sheets	Rods	Roots
0	23	595	0	0	17.96±0.50	5.73±0.20	2.83±0,16
500	19	567	158	0	15.46±0.54*	5.22±0.12*	3.44±0,08*
1000	17.5	487	182	28	15.47±0.51*	5.59±0.12	4.67±0,08*
1500	18	334	298	131	17.64±0.39	5.38±0.11	3.85±0,17*
2000	18.5	266	184	141	18.81±0.36	5.43±0.14	4.64±0,09*
2500	19	158	398	157	19.32±0.47*	5.25±0.35	4.21±0,35*
3000	17.8	165	390	149	17.52±0.67	5.32±0.36	5.42±0,11*


(*) the average of three replication are significantly different compared to control at p<0.05.

343

Table 4: Effect of lead on different pigments in *Coriandrum sativum* L.

	20 th Day				40 th Day			
	Chlorophyll mg.g ⁻¹ Fresh tissue weight			Total Carotenoid	Chlorophyll mg.g ⁻¹ Fresh tissue weight			Total Carotenoid
	a	b	Total		a	b	Total	
Control	9,2±1.1	7,2±1.2	16,4±2.6	6.0±2.9	7,5±1.0	3,6±0.4	11,1±1.1	3.0±0.8
500	8,3±0.7	5,5±2.4	13,7±2.1	3,1±1.3	7,2±1.6	3,4±1.0	10,6±1.6	2,7±0.6
1000	6,9±1.4	3,2±0.9*	10,1±2.0*	1,7±1.8	7,1±1.6	3,2±0.3	10,4±1.9	2,7±0.5
1500	5,2±1.3*	3,5±1.1*	8,7±1.2*	1,7±2.4	6,8±2.6	3,2±1.1	10,0±2.4	2,6±0.2
2000	4,8±0.8*	2,1±1.7*	6,9±2.5*	1,4±1.1*	6,3±2.1	2,9±0.8	9,2±1.4	2,4±0.5
2500	3,2±1.3*	2,0±1.76*	5,2±1.7*	0,5±0.2*	5,7±1.8	2,6±0.7	8,3±1.3*	2,1±0.7
3000	2,2±0.8*	1,1±0.9*	3,3±2.6*	0,3±0.1*	5,0±1.2	2,2±0.4*	7,2±1.8*	1,7±0.6

344 (*) the average of three replication are significantly different compared to control at p<0.05.



345

346 **Figure 1** : Effect of different lead level on lipid peroxidation and hydrogen peroxide of *Coriandrum*
 347 *sativum* L. ((*) the average of three replication are significantly different compared to control at
 348 p<0.05.)

349

350

351

352

353

354

355

356

357

358

359

360

Table 5: Effect of lead on soluble proteins ($\mu\text{g/g}$ fresh tissue weight) and proline (mM.g^{-1}) in *Coriandrum sativum* L. after 20 days

	Sheets		Rods		Roots	
	Soluble Protein	Proline	Soluble Protein	Proline	Soluble Protein	Proline
Control	114,8 \pm 6.4	1,8 \pm 0.5	8,4 \pm 1.1	1,1 \pm 0.6	103,6 \pm 10.2	0,6 \pm 0.4
500	58,8 \pm 4.8*	2,3 \pm 0.7	16,8 \pm 1.0*	2,9 \pm 0.4*	98,0 \pm 9.5	1,3 \pm 0.5
1000	67,2 \pm 5.7*	3,3 \pm 0.5*	47,6 \pm 3.2*	3,8 \pm 0.7*	224,0 \pm 20.2*	1,9 \pm 0.9
1500	98,0 \pm 4.9*	4,8 \pm 0.5*	22,4 \pm 2.2*	4,9 \pm 1.0*	302,4 \pm 34.3*	4,7 \pm 1.4*
2000	151,2 \pm 15.5*	6,3 \pm 1.07*	25,2 \pm 2.7*	5,9 \pm 0.9*	355,6 \pm 40.1*	9,7 \pm 2.2*
2500	112,0 \pm 16.4	10,9 \pm 2.2*	123,2 \pm 15.5*	6,9 \pm 1.6*	291,2 \pm 35.5*	12,6 \pm 2.8*
3000	179,2 \pm 25.5*	12,8 \pm 3.3*	114,8 \pm 13.4*	9,0 \pm 2.2*	431,3 \pm 55.5*	16,9 \pm 4.6*

(*) the average of three replication are significantly different compared to control at $p < 0.05$.

Table 6: Effect of lead on soluble proteins ($\mu\text{g/g}$ fresh tissue weight) and proline (mM.g^{-1}) in *Coriandrum sativum* L. after 40 days

	Sheets		Rods		Roots	
	Soluble Protein	Proline	Soluble Protein	Proline	Soluble Protein	Proline
Control	30,8 \pm 4.5	0,009 \pm 0.001	28,0 \pm 2.22	1,01 \pm 0.29	19,6 \pm 3.12	1,1 \pm 0.44
500	61,6 \pm 6.6*	0,014 \pm 0.004	22,4 \pm 1.75*	1,72 \pm 0.42	19,6 \pm 4.42	2,3 \pm 0.82
1000	58,8 \pm 3.7*	0,019 \pm 0.003*	14,0 \pm 0.95*	1,90 \pm 0.72	14,0 \pm 2.71	2,7 \pm 0.42*
1500	50,4 \pm 3.1*	0,021 \pm 0.007*	8,4 \pm 1.40*	3,51 \pm 1.21*	14,0 \pm 3.65	3,0 \pm 0.75*
2000	75,4 \pm 7.4*	0,021 \pm 0.005*	11,2 \pm 0.75*	4,85 \pm 1.53*	11,2 \pm 1.71*	4,0 \pm 1.09*
2500	58,8 \pm 5.2*	0,022 \pm 0.006*	25,2 \pm 2.33	6,50 \pm 1.95*	19,6 \pm 3.10	5,3 \pm 1.17*
3000	81,2 \pm 8.8*	0,027 \pm 0.004*	30,8 \pm 4.41	8,31 \pm 2.12*	25,2 \pm 2.73	7,4 \pm 1.54*

(*) the average of three replication are significantly different compared to control at $p < 0.05$.