

Remote Sensing and Geographic Information System for Optimizing Land Use Base on Fertility Capability Classification

Abdel-rahman A. Mustafa

Soil & Water Department, Faculty of Agriculture, Sohag University, Sohag, Egypt

a_mustafa32@yahoo.com

ABSTRACT

Soil is one of the most precious national resources and the knowledge of soil resources of an area is vital for optimizing land use and any developmental activities. Remote sensing and GIS have emerged as extremely valuable tools to study the soil resources, their potential for various use and problems. Hence an attempt has been made to study the soils of some soils of the Eastern Desert Part of Sohag Governorate and map them based on the fertility capability classification (FCC) using remote sensing and GIS. False color composite (FCC) of Landsat ETM imageries were visually interpreted incorporated with Digital Elevation Model (DEM) which generated from the Shuttle Radar Topographic Mission (SRTM). Different imaging interpretation units were identified and soil pedons were examined in each unit. Horizon wise soil samples were collected and analyzed for physiochemical properties by adopting standard procedures. Based on the results, the major landforms of the studied area were described as Wadi Bottom (WB), Bajada (B), Alluvial Fans (AF), Tableland (T), Gently Undulating Sand Sheet (GUS) and Undulating Sand Sheet (US). The type, substrata type and condition modifiers were also identified for each landform. The main condition modifiers of the study area were texture (S), low CEC (e), K deficiency (k), calcareous (b), salinity (s), dry condition (d), gravels (r) and low organic matter (m). Relevant FCC units were assigned to various landforms based on the type, substrata type and condition modifiers. A utility map was prepared using GIS with the FCC units, their limitations and extent distribution. Generally, the fertility of these soils was poor on account of low organic matter, total nitrogen, available phosphorus, potassium and micronutrients. Also, the water retentively was not satisfactory by the virtue of poor organic matter and higher percentage of coarser fraction. Based on the fertility constrains various soil management practices have been suggested to optimize the land use.

Keywords: Remote sensing, GIS, Land used, Fertility capability classification, Landforms.

1. INTRODUCTION

Soils are one of the most precious natural resources and the basic soil resource information is a prerequisite for planning sustainable agriculture and for optimizing land use and developmental activities. Natural soil classification systems such as Soil Taxonomy place

E-mail address: a_mustafa32@yahoo.com

more emphasis on subsurface than on topsoil properties, because of their permanent nature, whereas most soil managements practices are largely limited to the plowed layer. To bridge the gap between soil classification and soil fertility, fertility capability classification (FCC) system has been used. The need for Fertility Capability Classification (FCC) therefore arose out of identified technical problems of soil fertility maintenance in our fragile soils and the need for appropriate technology to improve fertility management of soils. FCC is a technological tool for agricultural land management that shows graphically the different fertility limitation sites in an area and the kinds of fertility management problems faced by users of the land. The FCC focuses attention on surface soil properties most directly related to management of field crops and is best used as an interpretative classification in conjunction with a more inclusive natural soil classification. The FCC, or some modification thereof, can serve as the basic for grouping soils for specific soil management evaluations and land use planning. Remote sensing and GIS have emerged as an extremely valuable tool to study the soil resources, their potential for various land use and problems. [1] classified some soils of Akwa Ibom State in South Eastern Nigeria and they found that gleying (g), low potassium reserve (k) and acidic reaction (h) were the general constraints of these soils. Also, some other inland depression soils had sandy (S) top soils while Bku had loamy top (L) soils but the three sub soils were loamy (L). All the floodplain soils had sandy (S) soils at both top and subsoils. For effective management of these soils, application of organic manure (including cattle manure) would supply the basic cations including K, as well as reduce soil acidity. [2] identified, with the aim of identifying three fertility capability classes dominate in the soils of Sokoto-Rima flood plains at Sokoto Nigeria, namely Lgm (Loamy soils low in organic matter with gleying limitations) ; Lghm (Loamy soils, low in organic matter and with gleying and pH limitation) and Sgm (Sandy soils low in organic matter and with gleying limitations). The three classes were then resolved to form the three mapping units shown in the FCC map. Soil class Lghm has higher fertility/yield potential followed by the class Lgm then Sgm class. Periodic monitoring of soil quality, adding organic manure and applying ameliorative measures such as liming can improve and sustain productive capacity of the soils. [3] studied the soils of Cameroon lowlands and classified them into fertility capability classes. Also, they identified soil fertility limitations as Fe- and Al-toxicities (a), low nutrient capital reserves (k), high leaching potential (e), and micronutrient deficiencies (Fe and Zn). The lowland soils were classified as: Lagk, Cagk, Laegk, Cbgm, Caeg, Lbg, Lgk, Cgv, LCg and Cgv, which reflect these limitations. Hence, an attempt has been made to study the soils of the Eastern Desert Part of Sohag Governorate and map them based on fertility capability classification using remote sensing and GIS. This will be the key for applying the efficient soil management practices for sustainable agriculture production especially in newly reclaimed area.

2. MATERIAL AND METHODS

2.1. Study area

The study area is a part of the Eastern Desert of Sohag, Egypt and located between geo-coordinates 26° 25' to 26° 45' latitudes (N) and 32° 40', 33° 00' longitudes (E) covering about 121,316 feddan. It is situated between the Nile Valley in the West and the Red Sea mountains in the East. The location map of the studied area is shown in figure (1).

The area under study is characterized by hot dry sub-humid to semi-arid transition with intense hot summer, cold winter and general dryness throughout the year except during July and September. The maximum temperature goes up to 45° C in the month of June. The lowest temperature goes down to 6.5° during January and February. The relative humidity (RH) ranges between 30% and 56% and the average about 43% in summer and 48% in winter. Prevailing winds are dominantly from the northwest to the southeast with an average maximum speed of 10 knots/h. The area receives mean annual rainfall ranging between 2.75 and 50 mm at the extreme Southeastern zone, while heavy showers are recorded occasionally during winter causing flash floods [4,5].






Fig.1 Location map of the studied area

2.2.Methodology

2.2.1.Remote Sensing data and processing

In the present study the Landsat ETM+ satellite data of 2010 was used. The study area is covered by one image (172Path /42 Row). The false color composite of the study area is presented in figure (2). The digital data of geo-coded cloud free of three images was downloaded from <http://glcf.umd.edu/data/landsat/>[6]. Table 1 presents the principle specifications of the sensor used in the investigation. The Shuttle Radar Topographic Mission (SRTM) images of 30 pixel size resolution have been used to generate the DEM for the study area and its surrounding were consulted to represent the area landscape. The study area was extracted from the whole image (Fig.2) of through on screen digitization of the area of interest (AOI) and masking out using subset module of ENVI software ver.4.8 (Research Systems Inc., Boulder, CO, USA).

Table. 1 Satellite and sensor specifications

	Bands	Spatial resolution (m)	Spectral resolution (μm)
1	Blue	30	0.414 – 0.514
2	Green	30	0.519 - 0.601
3	Red	30	0.631 – 0.692
4	NIR	30	0.772 – 0.898
5	SWIR-1	30	1.547 – 1.749
6	TIR	60	10.31 – 12.36
7	SWIR-2	30	2.064 – 2.345
8	Pan	15	0.515 – 0.896




Fig.2 False color composite of Landsat image of the studied area

2.2.2.Delineation of different landforms:

The delineation of the landform units from the satellite data needs a high spatial resolution images; therefore the spatial resolution of the used Landsat ETM+ was enhanced through the data merge process. This process is commonly used to enhance the spatial resolution of multi-spectral datasets using higher spatial resolution panchromatic data or single band (band 8). In this study merged data were performed using multi-spectral bands (30 m) as a low spatial resolution with panchromatic band 8 of ETM+ satellite image as a high spatial resolution (15 m) resulting in multi-spectral data with high spatial resolution (15 m). The landforms map has been generated from the SRTM (30 m) and enhanced Landsat ETM+ images using the ENVI 4.8 software [7].

By using the image elements such as texture, parcelling, pattern, shape, size, color, site and situation, many information about the terrain have been extracted from enhanced ETM+ image. Moreover, The SRTM data has been used in conjunction with enhanced ETM+ to provide a better visualization of the topographic features, namely surface elevation, slope, aspect, shaded relief and convexity. The topographic features have extracted using ENVI 4.8 software. Afterwards, the landform units were defined and classified and the map legend was established. DEM of the study area has been generated from the SRTM image using ArcGIS 9.3 software. The extracted data generates a preliminary geomorphologic map which was checked and completed through field observation.

2.2.3. Field work and samples collection:

A rapid reconnaissance survey of the area under study was conducted in order to achieve more detailed information of the soil patterns, land forms and characteristic of the landscape and landforms occurring in the study area.

Twelve soil profiles were selected representing various types of landforms occurring in the study area. The morphological examination of soil profiles was carried out in the field as per procedures laid out in the Soil Survey Manual [8]. Horizon wise disturbed soil samples (1 Kg) as well as core samples (diameter 2.5 cm and length 6 cm) were collected from each profile and kept separately in polyethylene bags for further analysis. Location coordinates were recorded with hand held GPS under WGS 84 (Lat-Lon) coordinate system (Fig.3).




Fig.3 Location of the representative soil profiles laid on studied area

2.2.4. Laboratory analysis and soil classification:

The collected soil samples were subjected for the following analyses: Particle size distribution [9], calcium carbonate, electric conductivity (ECe) in the soil paste extract, soluble cations and anions, soil pH, organic matter content [10]; cation exchange capacity and exchangeable sodium [11].

The American Soil taxonomy [12] was followed to classify the different soils of the studied area up to the family level. Then the correlation between the physiographic and taxonomic units, were identified [13].

2.2.5. Fertility capability classification:

Each landform were further classified under FCC system proposed by [14] and later modified by [15]. The FCC system consists of three categories viz., Type (topsoil texture or upper 20 cm depth), substrata type (subsoil texture between 20 and 50 cm depth) and condition modifiers (physical or chemical properties which influence the interaction between soil and fertilizer materials).

2.2.6. Generation of thematic maps

Inverse Distance Weighted (IDW) interpolation determines cell values using a linearly weighted combination of a set of sample points. The weight is a function of inverse distance. IDW lets the user control the significance of known points on the interpolated values, based on their distance from the output point. Thematic maps were generated using IDW interpolation provided in Arc GIS 9.3 software [16].


3. RESULTS AND DISCUSSION

3.1. Characterization of map units

The visual interpretation of the Landsat data and DEM integrated with Soil Taxonomy and soil field data using GIS have been used to generate the slope map and physiographic soil map (Fig. 5 and 6). The studied soils are classified according to USDA (2010) as TypicHaplocalcids, TypicTorripsamment and TypicTorriorthents (Table 2). The main soil characteristics of the mapping units are shown in Table (3).

The physiography of the studied area was identified based on the Landsat ETM+ images, the Digital Elevation Model (DEM) and slope map (Fig 4 and 5). The obtained results revealed that, there were six physiographic units in the area under studied (Fig. 6) viz. the Wadi Bottom (WB), Bajada (B), Alluvial Fans (AF), Tableland (T), Gently Undulating Sand Sheet (GUS) and Undulating Sand Sheet (US). The detailed characteristics of these physiographic units were discussed by [17].

161



162





Fig.4 DEM of of the studied area

163

164

165



166




Fig.5 Slope map of the studied **ied** area

167

168

169

170




Fig. 6 Physiographic units map of the studied area

Table2. Legend of the physiographic map of the studied area

Landscape	Lithology	Relief	Landform	Land use	Map unit symbol	Sub group	Area Feddan (1000)	%
Wadi Plain (WP)	Eocene Deposits (1)	Almos t Flat (1)	Wadi Bottom (WB)	Barren	WP11 _{WB}	TypicHaplocalcid s	26.426	21.78
			Alluvial Fans (AF)	Barren	WP11 _{AF}	TypicTorriorthent s	33.457	27.58
			Bajada (B)	Barren	WP11 _B	TypicHaplocalcid s	15.785	13.02
			Tableland (T)	Barren	WP11 _T	TypicTorriorthent s	16.648	13.72
		Gently Undul ating (2)	Gently Undulatin g sand sheet (GUS)	Barren	WP12 _{GUS}	TypicTorripsamm ents	16.500	13.60
		Undul ating (3)	Undulatin g sand sheet (US)	Barren	WP13 _{US}	TypicTorripsamm ents	12,500	10.30
Total							121.316	100

3.2. Fertility capability classification:

Based on analytical results (Table 3), the FCC units were established. The type, substrata type and condition modifiers were also identified. The main condition modifiers of the study

areawere texture (S), low CEC (e), K deficiency (k), calcareous (b), salinity (s), dry condition (d), gravels (r) and low organic matter (m). Relevant FCC units were assigned to various map units (Tables 4 & 5) and fig. 7 .

The results of FCC units of WP11WB, WP11AF, WP11B and some parts of WP12GUS and WP13US were classified as Sekbsdr (1-2%) only an area of 3125 feddan of WP11B was classified as Sekbsdr (2-4%). This implies that these map units have sandy (S) soils at both top and subsoils. The soils also have constraints of high leaching potential (e), low nutrients reserve (k), basic reaction (b) and salinity (s). As the soil exhibit ustic or xeric soil moisture regime, the Soil moisture stress constraint (d) has been recognized. The other modifiers are because of gravels content (r) and low organic matter (m). The soils of WP11T were classified as SekbsdrSRm (8-10%) and SekbsdrSRm (10-12%) which having the same condition modifiers but different slope grade. These soils are characterized by a high risk of soil erosion (SR) that erosion can negatively affect plant productivity and ecosystem functions . The FCC unit Sekbdr (1-2%) has been found in some areas belongs to WP12GUS and WP13US.

Table 3.The main soil characteristics of the mapping units

Unit		WP11WB		WP11AF		WP11T		WP11B		WP12GUS		WP13US	
Profile No.		7	12	9	4	10	11	5	8	2	1	6	3
1-Climate (c)													
Annual rainfall	mm	0	0	0	0	0	0	0	0	0	0	0	0
Mean temperature	°C	32.0	32.0	32.0	32.0	32.0	32.0	32.0	32.0	32.0	32.0	32.0	32.0
Relative humidity	%	54	54	54	54	54	54	54	54	54	54	54	54
Actual sunshine	hrs	11.2	11.2	11.2	11.2	11.2	11.2	11.2	11.2	11.2	11.2	11.2	11.2
2-Soil physical characteristics													
Depth	cm	75	80	100	95	100	100	90	100	85	90	95	90
Gravels	%	4.65	5.36	11.65	12.86	28.37	34.64	5.79	5.59	5.48	6.37	6.06	7.24
Coarse sand	%	75.10	75.71	82.73	83.41	86.11	84.26	75.73	76.74	91.51	91.21	91.47	93.20
Fine sand	%	6.70	6.21	5.35	5.37	5.43	6.88	9.25	8.91	1.44	1.83	1.90	2.00
Silt	%	12.00	11.03	7.24	7.91	5.03	4.99	10.61	10.31	5.14	4.86	4.61	3.10
Clay	%	6.20	7.05	4.68	3.31	3.44	3.88	4.42	4.04	1.91	2.10	2.02	1.70
Texture	ls	ls	ls	ls	s	GS	GS	ls	ls	s	s	s	s
3-Topography													
Slope	%	1-2	1-2	1-2	1-2	8-10	10-12	1-2	2-4	1-2	1-2	1-2	1-2
4-Wetness													
Drainage		well	well	well	well	well	well	well	well	well	well	well	well
Flood duration	Months	F0	F0	F0	F0	F0	F0	F0	F0	F0	F0	F0	F0
5-Fertility													
pH		8.09	7.83	8.16	8.28	8.39	8.36	8.09	8.41	8.45	8.37	8.24	7.95
Total Nitrogen	%	0.01	0.02	0.05	0.07	0.02	0.01	0.04	0.03	0.02	0.02	0.03	0.02
Organic carbon	%	0.16	0.15	0.08	0.10	0.08	0.08	0.11	0.11	0.11	0.08	0.13	0.07
Available P	mg/kg	6.0	5.4	5.6	7.1	6.5	4.4	4.6	5.0	3.4	5.0	4.6	4.3
Exchangable Na	Cmol+/kg	0.41	0.37	0.24	0.24	0.27	0.24	0.27	0.31	0.21	0.20	0.29	0.30
Exchangable K	Cmol+/kg	0.19	0.14	0.19	0.18	0.12	0.16	0.19	0.17	0.12	0.13	0.16	0.15
Exchangable Ca	Cmol+/kg	2.06	2.67	1.93	2.52	1.53	2.26	2.87	1.62	1.73	1.52	1.91	1.84
Exchangable Mg	Cmol+/kg	1.45	1.46	0.87	0.70	0.68	0.69	0.76	1.02	1.65	0.90	0.75	0.81
CEC	Cmol+/kg	4.26	4.76	3.38	3.75	2.68	3.46	4.19	3.20	3.82	2.81	3.19	3.18
Base saturation	%	96.71	97.54	95.57	96.76	97.61	96.58	97.49	97.05	97.12	97.62	97.21	97.48
ESP	%	9.67	7.76	7.33	6.43	10.35	6.76	6.69	9.84	4.85	7.94	9.18	9.51
DTPA extractable Fe	mg/kg	1.2	1.7	1.7	1.3	0.9	1.2	1.5	1.3	0.7	0.7	0.6	0.7
DTPA extractableMn	mg/kg	0.5	0.4	0.3	0.3	0.5	0.4	0.4	0.3	0.3	0.4	0.5	0.3
DTPA extractable Zn	mg/kg	0.3	0.4	0.4	0.5	0.3	0.4	0.5	0.5	0.2	0.2	0.2	0.2
DTPA extractable Cu	mg/kg	0.2	0.2	0.3	0.2	0.1	0.07	0.2	0.1	0.05	0.07	0.1	0.2
Salinity (ECe)	dS/m	5.69	5.41	10.67	7.38	5.38	5.85	5.02	6.45	5.58	3.83	4.28	3.99
ESP	%	9.67	7.76	7.33	6.43	10.35	6.76	6.69	9.84	4.85	7.94	9.18	9.51
CaCO ₃	%	12.36	13.41	17.08	13.24	8.59	9.19	13.62	17.68	7.01	3.81	8.65	5.44

203

Table 4. Soil fertility limitations and fertility capability classification units

Map unit	Profile No.	Type	Substrata Type	Condition modifiers										Area Feddan (1000)	FCC unit
				e	k	b	s	n-	d+	r+	SR	m	%		
WP11WB	7	S	S	+	+	+	+	-	+	+	-	+	0.8	26.426	Sekbsdrm (1-2%)
	12	S	S	+	+	+	+	-	+	+	-	+	0.7		Sekbsdrm (1-2%)
WP11AF	9	S	S	+	+	+	+	-	+	+	-	+	0.4	33.457	Sekbsdrm (1-2%)
	4	S	S	+	+	+	+	-	+	+	-	+	0.1		Sekbsdrm (1-2%)
WP11T	10	S	S	+	+	+	+	-	+	+	+	+	9.8	11.108	SekbsdrSRm (8-10%)
	11	S	S	+	+	+	+	-	+	+	+	+	10.5	5.54	SekbsdrSRm (10-12%)
WP11B	5	S	S	+	+	+	+	-	+	+	-	+	0.6	12.66	Sekbsdrm (1-2%)
	8	S	S	+	+	+	+	-	+	+	-	+	3	3.125	Sekbsdrm (2-4%)
WP12GUS	2	S	S	+	+	+	+	-	+	+	-	+	0.9	12.75	Sekbsdrm (1-2%)
	1	S	S	+	+	+	-	-	+	+	-	+	0.6	3.75	Sekbdrm (1-2%)
WP13US	6	S	S	+	+	+	+	-	+	+	-	+	1	7.85	Sekbsdrm (1-2%)
	3	S	S	+	+	+	-	-	+	+	-	+	0.7	4.65	Sekbdrm (1-2%)

S: sandy, e: low CEC, k: low nutrient reserves, b: calcareous, s: salinity, n⁻: nitric, d⁺: dry soil moisture condition, r⁺: gravels, SR: erosion, m: low organic matter and %: slope.

204
205
206
207
208**Table 5. Interpretation of Soil fertility capability classification units**

Map unit	FCC unit	Description
WP11WB , WP11AF	Sekbsdrm (1-2%)	Sandy surface and subsurface soils having low cation exchange capacity, low nutrients reserves, calcareous reaction, salinity. Soils with dry conditions, gravels and deficient in soil organic carbon.
WP11T	SekbsdrSRm (8-10%)	Sandy surface and subsurface soils having low cation exchange capacity, low nutrients reserves, calcareous reaction, salinity. Soils with dry conditions, gravels , erosion risk and deficient in soil organic carbon with steep slope.
	SekbsdrSRm (10-12%)	Sandy surface and subsurface soils having low cation exchange capacity, low nutrients reserves, calcareous reaction, salinity. Soils with dry conditions, gravels , erosion risk and deficient in soil organic carbon with steep slope.
WP11B	Sekbsdrm (1-2%)	Sandy surface and subsurface soils having low cation exchange capacity, low nutrients reserves, calcareous reaction, salinity. Soils with dry conditions, gravels and deficient in soil organic carbon.
	Sekbsdrm (2-4%)	Sandy surface and subsurface soils having low cation exchange capacity, low nutrients reserves, calcareous reaction, salinity. Soils with dry conditions, gravels and deficient in soil organic carbon.
WP12GUS	Sekbsdrm (1-2%)	Sandy surface and subsurface soils having low cation exchange capacity, low nutrients reserves, calcareous reaction, salinity. Soils with dry conditions, gravels and deficient in soil organic carbon.
	Sekbdrm (1-2%)	Sandy surface and subsurface soils having low cation exchange capacity, low nutrients reserves, calcareous reaction. Soils with dry conditions, gravels and deficient in soil organic carbon.
WP13US	Sekbsdrm (1-2%)	Sandy surface and subsurface soils having low cation exchange capacity, low nutrients reserves, calcareous reaction, salinity. Soils with dry conditions, gravels deficient in soil organic carbon.
	Sekbdrm (1-2%)	Sandy surface and subsurface soils having low cation exchange capacity, low nutrients reserves, calcareous reaction. Soils with dry conditions, gravels deficient in soil organic carbon.

209
210
211
212
213
214
215
216
217
218
219
220
221
222





Fig.7 Fertility capability classification (FCC) units of the studied area

3.3 Suggestive plausible soil managements:

Now, there is a raised question i.e. at what time scales are FCC attributes refer today, months, years, decades or centuries? And hence the scientific management technologies can be applied for mitigating these constrains. Experience in using FCC indicates that some of the condition modifiers can be changed with management at different time scales. In the current study, the possibility of overriding constrains is presented in table 6.

Table 6. The temporal scale dimension of FCC attributes

FCC attribute	Can be changed by management with time (years)				Means of change
	<1	1-10	10-100	>100	
Type/substrata type - S				√	inherent, unless severely eroded
High leaching potential- e				√	inherent
Low nutrient reserves -k				√	inherent
Calcareous -b			√		by sustained leaching in slightly calcareous ones
Saline - s		√			by effective leaching
Sodic - n		√			by effective leaching
Soil moisture stress - d	√				temporarily by irrigation
Gravels- r				√	inherent, unless severely eroded
High erosion - SR				√	inherent; can be mitigated by soil conservation practices
Low organic matter - m		√			by organic input application rates that exceed decomposition rate

From the previous table, some of the soil constrains cannot be changed in less than century (inherent) such as type/substrata type, high leaching potential, low nutrient reverses, gravels and high erosion risk. Whereas, condition modifiers can change at the decadal scale (10–

100 years) include calcareous reaction by sustained irrigation and subsequent leaching, salinity and sodicity by applying effective leaching and low level of organic matter which can be maintained under certain levels by supplying soil with different rates and sources of organic inputs. The soil water stress can be managed by applying the water through irrigation using the effective method of application such as trickle irrigation [18]. Some soil management considerations are mentioned hereunder:

3.3.1.Low organic matter (m) and low nutrient reserves (k)

Low organic matter content which is prevailing in all soil profiles can be improved through application of organic manure, green manuring, mulching, crop rotation and so on. Also base saturation can be improved by applying fertilizers and amendments. Use of nitrogen and phosphorus fertilizers to mitigate major nutrient deficiencies is a must.

3.3.2.Salinity (s)

This can be removed by applying leaching and supplying the affected area with efficient drainage system in case of good quality water. Whereas, if the quality of irrigation water is poor due to either high salinity or high alkalinity or both, some suggestive management plans can be adopted such:

(1) In case of saline area and high salinity irrigation water, subsurface drainage system is a useful tool for desalinization.

(2) In case of saline area and high sodic irrigation water, subsurface drainage system along with application of gypsum could be used for improving the productivity. The gypsum amount to be added is determined by quality and quantity of water to be added per year by applying the simple equation [19]:

$$GR = ((RSC - 2.5) \times N \times 36)$$

Where:

GR: Gypsum requirement (tons/acre), RSC: Residual sodium carbonate, N: number of irrigations.

Thus, for soils irrigated with water having RSC 10.9, 10.4, 8.4 and 5.5 me/l and needing 5 irrigations, the GR will be 1512, 1422, 1062 and 540 kg/acre.

3.3.3.High ESP soils (n):

Application of gypsum to soils along with deep ploughing and subsurface drainage is recommended. GR can be calculated by using the following equation:

$$GR(tons / ha) = \frac{(ESP_I - ESP_F) \times CEC \times 25.8 \times P}{100}$$

Where

ESPI: Exchangeable Sodium Percentage initial (ESP) of soil, ESPF: ESP final, CEC: Cation exchange capacity of soil, P: purity factor of gypsum.

3.3.4.High erosion (SR)

Following measures are suggested to reclaim high erosion land:

- Leveling and construction of contour bunds.
- Pipe outlets or ramps with suitable grasses for draining excess run off.
- Perennial vegetation like fuel, fodder trees and grasses may help effectively to conserve the soil.

3.3.5.Rocky and quarried (r):

Following measures can be adopted

- Enclosures of the hilly area with barbed wire.

- Prohibition of grazing.
- Locally suited tree species may be grown to conserve soils.
- Rehabilitation of quarry lands- plantation of suitable tree species.

4. CONCLUSION

According to the results, the major landforms of the studied area were described as Wadi Bottom (WB), Bajada (B), Alluvial Fans (AF), Tableland (T), Gently Undulating Sand Sheet (GUS) and Undulating Sand Sheet (US). The results of FCC units of WP11WB, WP11AF, WP11B and some parts of WP12GUS and WP13US were classified as Sekbsdr (1-2%) only an area of 3125 feddan of WP11B was classified as Sekbsdr (2-4%). This implies that these map units have sandy (S) soils at both top and subsoils. The soils also have constraints of high leaching potential (e), low nutrients reserve (k), basic reaction (b) and salinity (s). As the soil exhibit ustic or xeric soil moisture regime, the Soil moisture stress constraint (d) has been recognized. The other modifiers are because of gravels content (r) and low organic matter (m). The soils of WP11T were classified as SekbsdrSR (8-10%) and SekbsdrSRm (10-12%) which having the same condition modifiers but different slope grade. These soils are characterized by a high risk of soil erosion (SR) that erosion can negatively affect plant productivity and ecosystem functions. The FCC unit Sekbsdr (1-2%) has been found in some areas belongs to WP12GUS and WP13US. By following the scientific technologies, the fertility constraints can be improved.

Authors' Contributions

CONSENT (WHERE EVER APPLICABLE)

ETHICAL APPROVAL (WHERE EVER APPLICABLE)

REFERENCES

Reference to a journal:

1. Udo BY, Utip U, Kufre EI, Monday T, Idungafa MA. Fertility Assessment of Some Inland Depression and Floodplain (Wetland) Soils in Akwa Ibom State. J of Tropical Agri, Food, Env and Extension. 2009;8(1):14-19.
2. Yakubu AA, Ibrahim SA, Ojanuga AG, Singh A. Fertility Capability Classification of Soils of the Sokoto-Rima Flood Plain, Nigeria. Nig. J. Soil and Env. Res. 2012;10: 20: 63 – 70.
3. Tabi FO, Ngobesing ESC, Yinda GS, Boukong A, Omoko M, Bitondo D, Mvondo Ze AD. Soil fertility capability classification (FCC) for rice production in Cameroon lowlands. African J of Agri.Res. 2013;8(119):1650-1660.
4. Awad AA. Integration of Remote Sensing, Geophysics and GIS to evaluate groundwater potentiality – A Case study in Sohag region. Egypt.Geology Department, Fac. of Science, Assiut University, Assiut, Egypt. 2008.
5. EGMA. Egyptian Meteorological Authority database. 2004.
6. Global Land Cover Facility (GLCF). <http://glcf.umd.edu/data/landsat/>. 4321 Hartwick Building · College Park, Maryland 20740. 2014.
7. Dobos E, Norman B, Bruce W, Luca M, Chris J, Erika M. The Use of DEM and Satellite Data for Regional Scale Soil Databases. 17th World Congress of Soil Science (WCSS), Bangkok, Thailand. 2002; 649:14-21.
8. FAO. Guidelines for Soil Description. Fourth edition, Food and Agriculture Organization of the United Nations, Rome, Italy. 2006.

- 338 9. USSL Staff. Diagnosis and improvement of saline and alkali soils. Agriculture Handbook
339 60, Richards LA (ed.). USDA: Washington, DC. 1954.
340 10. Jackson ML . Soil Chemical Analysis. Prentice Hall of India Pvt. Ltd, New Delhi;1973.
341 11. Black CA. Methods of soil analysis. 2nd edition. Chemical and microbiological
342 properties. Agronomy series no. 9, ASA, SSSA, Madison, Wis., USA. 1982.
343 12. Soil Survey Staff. Keys to Soil Taxonomy, 9th edition, USDA National Resource
344 Conservation Services. 2003.
345 13. Elbersen GWW, Catalan R. Portable Computer in Physiographic Soil Survey Unit. Proc.
346 Inte. Soil Sci., Cong. Homburg. 1987.
347 14. Buol SW, Sanchez PA, Cate RB, Granger MA. Soil fertility capability classification: A
348 technical soil classification system for fertility management. In: Bornemisza, E, and A.
349 Alvarado (Eds), Soil management in tropical America. N. C. State University, Raleigh, N. C.
350 1975; 126-145.
351 15. Sanchez PA, Couto W, Buol SW. The Fertility Capability Classification System (FCC):
352 Interpretation, applicability and modification. Geoderma. 1982; 27: 283 – 309.
353 16. ESRI. ArcGIS Desktop: Release 9.3. Redlands, CA: Environmental Systems Research
354 Institute. 2009.
355 17. Mustafa AA, Negim OE. Geomatics Based Soil Mapping of The Eastern Desert Part of
356 Sohag Governorate, Egypt. J. Soil Sci. and Agric. Eng., Mansoura Univ. 2015;6 (12): 1527 –
357 1543.
358 18. Mustafa A A. Integration of parametric approach and GIS for optimum irrigation method
359 in soils of Eastern desert part of Sohag, Egypt. The 12th International Conference of The
360 Egyptian Society of Soil Science, 7-9 March, 2016, Ismailia, Egypt. 2016.
361 19. Rajput RP. Potentials and utilization of Brackish water in irrigation management. In
362 Proceeding of the training on “Diagnostic and improvement of problematic soils”, Aug 18 to
363 Sep 16, 2000), India. 2000.

364 DEFINITIONS, ACRONYMS, ABBREVIATIONS

365 **GIS**

366 **FCC**

367 **DEM**

368 **SRTM**

369 **WB**

370 **AF**

371 **GUS**

372 **US**

373 **FCC UNITS**

374 **USDA**