Remote Sensing and Geographic Information System for Optimizing Land Use Base on Fertility Capability Classification

Abdel-rahman A. Mustafa Soil & Water Department, Faculty of Agriculture, Sohag University, Sohag, Egypt a_mustafa32@yahoo.com

ABSTRACT

12

Soil is one of the most precious national resources and the knowledge of soil resources of an area is vital for optimizing land use and any developmental activities. Remote sensing and GIS have emerged as extremely valuable tools to study the soil resources, their potential for various use and problems. Hence an attempt has been made to study the soils of some soils of the Eastern Desert Part of Sohag Governorate and map them based on the fertility capability classification (FCC) using remote sensing and GIS. False color composite (FCC) of Landsat ETM imageries were visually interpreted incorporated with Digital Elevation Model (DEM) which generated from the Shuttle Radar Topographic Mission (SRTM). Different imaging interpretation units were identified and soil pedons were examined in each unit. Horizon wise soil samples were collected and analyzed for physiochemical properties by adopting standard procedures. Based on the results, the major landforms of the studied area were described as Wadi Bottom (WB), Bajada (B), Alluvial Fans (AF), Tableland (T), Gently Undulating Sand Sheet (GUS) and Undulating Sand Sheet (US). The type, substrata type and condition modifiers were also identified for each landform. The main condition modifiers of the study area were texture (S), low CEC (e), K deficiency (k), calcareous (b), salinity (s), dry condition (d), gravels (r) and low organic matter (m). Relevant FCC units were assigned to various landforms based on the type, substrata type and condition modifiers. A utility map was prepared using GIS with the FCC units, their limitations and extent distribution. Generally, the fertility of these soils was poor on account of low organic matter, total nitrogen, available phosphorus, potassium and micronutrients. Also, the water retentively was not satisfactory by the virtue of poor organic matter and higher percentage of coarser fraction. Based on the fertility constrains various soil management practices have been suggested to optimize the land use.

13 14

Keywords: Remote sensing, GIS, Land used, Fertility capability classification, Landforms.

15 16 17

18 **1. INTRODUCTION**

19 Soils are one of the most precious natural resources and the basic soil resource information 20 is a prerequisite for planning sustainable agriculture and for optimizing land use and developmental activities. Natural soil classification systems such as Soil Taxonomy place 21

E-mail address: a mustafa32@yahoo.com

22 more emphasis on subsurface than on topsoil properties, because of their permanent nature, 23 whereas most soil managements practices are largely limited to the plowed layer. To bridge 24 the gap between soil classification and soil fertility, fertility capability classification (FCC) 25 system has been used. The need for Fertility Capability Classification (FCC) therefore arose 26 out of identified technical problems of soil fertility maintenance in our fragile soils and the 27 need for appropriate technology to improve fertility management of soils. FCC is a technological tool for agricultural land management that shows graphically the different 28 29 fertility limitation sites in an area and the kinds of fertility management problems faced by 30 users of the land. The FCC focuses attention on surface soil properties most directly related to management of field crops and is best used as an interpretative classification in 31 conjunction with a more inclusive natural soil classification. The FCC, or some modification 32 thereof, can serve as the basic for grouping soils for specific soil management evaluations 33 and land use planning. Remote sensing and GIS have emerged as an extremely valuable 34 35 tool to study the soil resources, their potential for various land use and problems. [1] classified some soils of Akwa Ibom State in South Eastern Nigeria and they found that 36 37 gleying (g), low potassium reserve (k) and acidic reaction (h) were the general constraints of 38 these soils. Also, some other inland depression soils had sandy (S) top soils while Bku had 39 loamy top (L) soils but the three sub soils were loamy (L). All the floodplain soils had sandy 40 (S) soils at both top and subsoils. For effective management of these soils, application of 41 organic manure (including cattle manure) would supply the basic cations including K, as well 42 as reduce soil acidity. In other study conducted in Sokoto-Rima flood plains at Sokoto 43 Nigeria [2] identified three fertility capability classes dominate in the soils, namely Lgm 44 (Loamy soils low in organic matter with gleving limitations); Lghm (Loamy soils, low in 45 organic matter and with gleying and pH limitation) and Sqm (Sandy soils low in organic matter and with gleving limitations). The three classes were then resolved to form the three 46 47 mapping units shown in the FCC map. Soil class Lghm has higher fertility/yield potential 48 followed by the class Lgm then Sgm class. Periodic monitoring of soil quality, adding organic 49 manure and applying ameliorative measures such as liming can improve and sustain 50 productive capacity of the soils. [3] classified the lowland soils of Cameroon as Lagk, Cagk, 51 Laegk, Cbgm, Caeg, Lbg, Lgk, Cgv, LCg and Cgv. In addition, they concluded that the main 52 soil fertility limitations were Fe- and Al-toxicities (a), low nutrient capital reserves (k), high 53 leaching potential (e), and micronutrient deficiencies. Hence, an attempt has been made to 54 study the soils of the Eastern Desert Part of Sohag Governorate and map them based on 55 fertility capability classification using remote sensing and GIS. This will be the key for 56 applying the efficient soil management practices for sustainable agriculture production 57 especially in newly reclaimed area.

58

59 2. MATERIAL AND METHODS

60 **2.1.Study area**

The study area is a part of the Eastern Desert of Sohag, Egypt and located between geocoordinates 26° 25` to 26° 45`latitudes (N) and 32° 40`, 33° 00` longitudes (E) covering about121,316 feddan. It is situated between the Nile Valley in the West and the Red Sea mountains in the East. The location map of the studied area is shown in figure (1).

65 The area under study is characterized by hot dry sub-humid to semi-arid transition with intense hot summer, cold winter and general dryness throughout the year except during July 66 and September. The maximum temperature goes up to 45° C in the month of June. The 67 lowest temperature goes down to 6.5° during January and February. The relative humidity 68 69 (RH) ranges between 30% and 56% and the average about 43% in summer and 48% in 70 winter. Prevailing winds are dominantly from the northwest to the southeast with an average 71 maximum speed of 10 knots/h. The area receives mean annual rainfall ranging between 72 2.75 and 50 mm at the extreme Southeastern zone, while heavy showers are recorded 73 occasionally during winter causing flash floods [4,5].





- 75 76
- --
- 77

Fig.1 Location map of the studied area

79 2.2.Methodology

80 2.2.1.Remote Sensing data and processing

In the present study the Landsat ETM+ satellite data of 2010 was used. The study area is 81 82 covered by one image (172Path /42 Row). The false color composite of the study area is presented in figure (2). The digital data of geo-coded cloud free of three images was 83 downloaded from http://glcf.umd.edu/data/landsat/[6]. Table 1 presents the principle 84 specifications of the sensor used in the investigation. The Shuttle Radar Topographic 85 Mission (SRTM) images of 30 pixel size resolution have been used to generate the DEM for 86 87 the study area and its surrounding were consulted to represent the area landscape. The 88 study area was extracted from the whole image (Fig.2) of through on screen digitization of the area of interest (AOI) and masking out using subset module of ENVI software ver.4.8 89 (Research Systems Inc., Boulder, CO, USA). 90

91 92

Table. 1 Satellite	e and sensor	specifications	
Bands	Spatial	Spectral	
	resolution	resolution (um)	

		resolution (m)	resolution (µm)
1	Blue	30	0.414 – 0.514
2	Green	30	0.519 - 0.601
3	Red	30	0.631 – 0.692
4	NIR	30	0.772 – 0.898
5	SWIR-1	30	1.547 – 1.749
6	TIR	60	10.31 – 12.36
7	SWIR-2	30	2.064 - 2.345
8	Pan	15	0.515 – 0.896





Fig.2 False color composite of Landsat image of the studied area

96 2.2.2.Delineation of different landforms:

97 The delineation of the landform units from the satellite data needs a high spatial resolution 98 images; therefore the spatial resolution of the used Landsat ETM+ was enhanced through 99 the data merge process. This process is commonly used to enhance the spatial resolution of 100 multi-spectral datasets using higher spatial resolution panchromatic data or single band (band 8). In this study merged data were performed using multi-spectral bands (30 m) as a 101 102 low spatial resolution with panchromatic band 8 of ETM+ satellite image as a high spatial 103 resolution (15 m) resulting in multi-spectral data with high spatial resolution (15 m). The 104 landforms map has been generated from the SRTM (30 m) and enhanced Landsat ETM+ 105 images using the ENVI 4.8 software [7].

106 By using the image elements such as texture, parcelling, pattern, shape, size, color, site and 107 situation, many information about the terrain have been extracted from enhanced ETM+ 108 image. Moreover, The SRTM data has been used in conjunction with enhanced ETM+ to 109 provide a better visualization of the topographic features, namely surface elevation, slope, aspect, shaded relief and convexity. The topographic features have extracted using ENVI 110 111 4.8 software. Afterwards, the landform units were defined and classified and the map legend 112 was established. DEM of the study area has been generated from the SRTM image using 113 ArcGIS 9.3 software. The extracted data generates a preliminary geomorphologic map which 114 was checked and completed through field observation.

115 **2.2.3. Field work and samples collection:**

A rapid reconnaissance survey of the area under study was conducted in order to achieve
 more detailed information of the soil patterns, land forms and characteristic of the
 landscapeand landforms occurring in the study area.

Twelve soil profiles were selected representing various types of landforms occurring in the study area. The morphological examination of soil profiles was carried out in the field as per procedures laid out in the Soil Survey Manual [8]. Horizon wise disturbed soil samples (1 Kg) as well as core samples (diameter 2.5 cm and length 6 cm) were collected from each profile and kept separately in polyethylene bags for further analysis. Location coordinates were recorded with hand held GPS under WGS 84 (Lat-Lon) coordinate system (Fig.3).





Fig.3 Location of the representative soil profiles laid on studied area

128 **2.2.4. Laboratory analysisand soil classification:**

The collected soil samples were subjected for the following analyses: Particle size distribution [9], calcium carbonate, electric conductivity (ECe) in the soil paste extract, soluble cations and anions, soil pH, organic matter content [10]; cation exchange capacity and exchangeable sodium [11].

133 The American Soil taxonomy [12] was followed to classify the different soils of the studied 134 area up to the family level. Then the correlation between the physiographic and taxonomic 135 units, were identified [13].

136 **2.2.5.Fertility capability classification:**

Each landform were further classified under FCC system proposed by [14] and later modified
by [15]. The FCC system consists of three categories viz., Type (topsoil texture or upper 20 cm depth), substrata type (subsoil texture between 20 and 50 cm depth) and condition
modifiers (physical or chemical properties which influence the interaction between soil and fertilizer materials).

142 2.2.6.Generation of thematic maps

143 Inverse Distance Weighted (IDW) interpolation determines cell values using a linearly 144 weighted combination of a set of sample points. The weight is a function of inverse 145 distance. IDW lets the user control the significance of known points on the interpolated 146 values, based on their distance from the output point. Thematic maps were generated

147 using IDW interpolation provided in Arc GIS 9.3 software [16].

1483. RESULTS AND DISCUSSION

149 **3.1. Characterization of map units**

The visual interpretation of the Landsat data and DEM integrated with Soil Taxonomy and soil field data using GIS have been used to generate the slope map and physiographic soil map (Fig. 5 and 6). The studied soils are classified according to USDA (2010) as TypicHaplocalcids, TypicTorripsamment and TypicTorriorthents (Table 2). The main soil characteristics of the mapping units are shown in Table (3).

- 155 The physiography of the studied area was identified based on the Landsat ETM+ images,
- the Digital Elevation Model (DEM) and slope map (Fig 4 and 5). The obtained results revealed
- that, therewere six physiographic units in the area under studied (Fig. 6) viz. theWadi Bottom (WB), Baiada (B), Alluvial Fans (AF), Tableland (T), Gently Undulating Sand Sheet (GUS)
- 158 (WB), Bajada (B), Alluvial Fans (AF), Tableland (T), Gently Undulating Sand Sheet (GUS)
- and Undulating Sand Sheet (US). The detailed characteristics of these physiographic units
 were discussed by [17].





Table2. Legend of the physiographic map of the studied area

Landscape	Lithology	Relief	Landform	Land	Map unit	Sub group	Are	ea
				use	symbol		Feddan (1000)	%
Wadi Plain (WP)	Eocene Deposits (1)	Almos t Flat (1)	Wadi Bottom (WB)	Barren	WP11 _{WB}	TypicHaplocalcid s	26.426	21.78
			Alluvial Fans (AF)	Barren	WP11 _{AF}	TypicTorriorthent s	33.457	27.58
			Bajada (B)	Barren	WP11 _B	TypicHaplocalcid s	15.785	13.02
			Tableland (T)	Barren	WP11 _T	TypicTorriorthent s	16.648	13.72
		Gently Undul ating (2)	Gently Undulatin g sand sheet (GUS)	Barren	WP12 GUS	TypicTorripsamm ents	16.500	13.60
		Undul ating (3)	Undulatin g sand sheet (US)	Barren	WP13 _{US}	TypicTorripsamm ents	12,500	10.30
Total		~ /					121.316	100

3.2. Fertility capability classification:

179 Based on analytical results (Table 3), the FCC units were established. The type, substrata

type and condition modifiers were also identified. The main condition modifiers of the study

areawere texture (S), low CEC (e), K deficiency (k), calcareous (b), salinity (s), dry condition
(d), gravels (r) and low organic matter (m). Relevant FCC units were assigned to various
map units (Tables 4 & 5) and fig. 7.

The results of FCC units of WP11WB, WP11AF, WP11B and some parts of WP12GUS and 184 185 WP13US were classified as Sekbsdrm (1-2%) only an area of 3125 feddan of WP11B was 186 classified as Sekbsdrm (2-4%). This implies that these map units have sandy (S) soils at both top and subsoils. The soils also have constraints of high leaching potential (e), low 187 188 nutrients reserve (k), basic reaction (b) and salinity (s). As the soil exhibit ustic or xeric soil 189 moisture regime, the Soil moisture stress constraint (d) has been recognized. The other modifiers are because of gravels content (r) and low organic matter (m). The soils of WP11T 190 were classified as SekbsdrSRm (8-10%) and SekbsdrSRm (10-12%) which having the same 191 condition modifiers but different slope grade. These soils are characterized by a high risk of 192 soil erosion (SR) that erosion can negatively affect plant productivity and ecosystem 193 functions. The FCC unit Sekbdrm (1-2%) has been found in some areas belongs to 194 195 WP12GUS and WP13US.

196 Table 3.The main soil characteristics of the mapping units

	Unit	WP11 _{WB}		WP	11_{AF}	WI	P11 _T	WE	P11 _B	WP	12 _{GUS}	WP13 _{US}	
Profile No.		7	12	9	4	10	11	5	8	2	1	6	3
1-Climate (c)													
Annual rainfall	mm	0	0	0	0	0	0	0	0	0	0	0	0
Mean temperature	^{0}C	32.0	32.0	32.0	32.0	32.0	32.0	32.0	32.0	32.0	32.0	32.0	32.0
Relative humidity	%	54	54	54	54	54	54	54	54	54	54	54	54
Actual sunshine	hrs	11.2	11.2	11.2	11.2	11.2	11.2	11.2	11.2	11.2	11.2	11.2	11.2
2-Soil physical													
charateristics													
Depth	cm	75	80	100	95	100	100	90	100	85	90	95	90
Gravels	%	4.65	5.36	11.65	12.86	28.37	34.64	5.79	5.59	5.48	6.37	6.06	7.24
Coarse sand	%	75.10	75.71	82.73	83.41	86.11	84.26	75.73	76.74	91.51	91.21	91.47	93.20
Fine sand	%	6.70	6.21	5.35	5.37	5.43	6.88	9.25	8.91	1.44	1.83	1.90	2.00
Silt	%	12.00	11.03	7.24	7.91	5.03	4.99	10.61	10.31	5.14	4.86	4.61	3.10
Clay	%	6.20	7.05	4.68	3.31	3.44	3.88	4.42	4.04	1.91	2.10	2.02	1.70
Texture		ls	ls	ls	s	GS	GS	ls	ls	s	s	s	s
3-Topography													
Slope	%	1-2	1-2	1-2	1-2	8-10	10-12	1-2	2-4	1-2	1-2	1-2	1-2
4-Wetness													
Drainage		well	well	well	well	well	well	well	well	well	well	well	well
Flood duration	Months	F0	F0	F0	F0	F0	F0	F0	F0	F0	F0	F0	F0
5-Fertility													
pH		8.09	7.83	8.16	8.28	8.39	8.36	8.09	8.41	8.45	8.37	8.24	7.95
Total Nitrogen	%	0.01	0.02	0.05	0.07	0.02	0.01	0.04	0.03	0.02	0.02	0.03	0.02
Organic carbon	%	0.16	0.15	0.08	0.10	0.08	0.08	0.11	0.11	0.11	0.08	0.13	0.07
Available P	mg/kg	6.0	5.4	5.6	7.1	6.5	4.4	4.6	5.0	3.4	5.0	4.6	4.3
Exchangable Na	Cmol+/kg	0.41	0.37	0.24	0.24	0.27	0.24	0.27	0.31	0.21	0.20	0.29	0.30
Exchangable K	Cmol+/kg	0.19	0.14	0.19	0.18	0.12	0.16	0.19	0.17	0.12	0.13	0.16	0.15
Exchangable Ca	Cmol+/kg	2.06	2.67	1.93	2.52	1.53	2.26	2.87	1.62	1.73	1.52	1.91	1.84
Exchangable Mg	Cmol+/kg	1.45	1.46	0.87	0.70	0.68	0.69	0.76	1.02	1.65	0.90	0.75	0.81
CEC	Cmol+/kg	4.26	4.76	3.38	3.75	2.68	3.46	4.19	3.20	3.82	2.81	3.19	3.18
Base saturation	%	96.71	97.54	95.57	96.76	97.61	96.58	97.49	97.05	97.12	97.62	97.21	97.48
ESP	%	9.67	7.76	7.33	6.43	10.35	6.76	6.69	9.84	4.85	7.94	9.18	9.51
DTPA extractable	mø/kø	1.2	1.7	1.7	1.3	0.9	1.2	1.5	1.3	0.7	0.7	0.6	0.7
Fe													
DTPA	mø/kø	0.5	0.4	0.3	0.3	0.5	0.4	0.4	0.3	0.3	0.4	0.5	0.3
extractableMn	88												
DTPA extractable	mg/kg	0.3	0.4	0.4	0.5	0.3	0.4	0.5	0.5	0.2	0.2	0.2	0.2
Zn	88												
DTPA extractable	mg/kg	0.2	0.2	0.3	0.2	0.1	0.07	0.2	0.1	0.05	0.07	0.1	0.2
Cu		5 60		10.55	7.00	5.00	5.05	5.00		0	2.02	4.00	2 00
Salinity (ECe)	dS/m	5.69	5.41	10.67	7.38	5.38	5.85	5.02	6.45	5.58	3.83	4.28	3.99
ESP	%	9.67	7.76	7.33	6.43	10.35	6.76	6.69	9.84	4.85	7.94	9.18	9.51
CaCO ₃	%	12.36	13.41	17.08	13.24	8.59	9.19	13.62	17.68	7.01	3.81	8.65	5.44

197

198

199

200

201

Map unit	Map unit Profile _ Substrata			Condition modifiers								Area	FCC unit		
	No. Type Type	Туре	e	k	b	s	n-	d+	r+	SR	m	%	Feddan (1000)		
WP11WB	7	S	S	+	+	+	+	-	+	+	-	+	0.8	26.426	Sekbsdrm (1-2%)
	12	S	S	+	+	+	+	-	+	+	-	+	0.7		Sekbsdrm(1-2%)
WP11AF	9	S	S	+	+	+	+	-	+	+	-	+	0.4	33.457	Sekbsdrm(1-2%)
	4	S	S	+	+	+	+	-	+	+	-	+	0.1		Sekbsdrm(1-2%)
WP11T	10	S	S	+	+	+	+	-	+	+	+	+	9.8	11.108	SekbsdrSRm (8-10%)
	11	S	S	+	+	+	+	-	+	+	+	+	10.5	5.54	SekbsdrSRm (10-12%)
WP11B	5	S	S	+	+	+	+	-	+	+	-	+	0.6	12.66	Sekbsdrm (1-2%)
	8	S	S	+	+	+	+	-	+	+	-	+	3	3.125	Sekbsdrm (2-4%)
WEIGONG	2	S	S	+	+	+	+	-	+	+	-	+	0.9	12.75	Sekbsdrm(1-2%)
WP12GUS	1	S	S	+	+	+	-	-	+	+	-	+	0.6	3.75	Sekbdrm(1-2%)
	6	S	S	+	+	+	+	-	+	+	-	+	1	7.85	Sekbsdrm(1-2%)
WP13US	3	S	S	+	+	+	-	-	+	+	-	+	0.7	4.65	Sekbdrm(1-2%)

Table 4.Soil fertility limitations and fertility capability classification units

S:sandy, e:low CEC, k:low nutrient reserves, b: calcareous, s: salinity, \mathbf{n} : nitric, \mathbf{d}^+ : dry soil moisture condition, \mathbf{r}^+ : gravels, SR: erosion, m: low organic matter and %: slope.

Table 5.Interpretation of Soil fertility capability classification units

Map unit	FCC unit	Description
WP11WB, WP11AF	Sekbsdrm (1-2%)	Sandy surface and subsurface soils having low cation exchange capacity, low nutrients reserves,
		calcareous reaction, salinity. Soils with dry conditions, gravels and deficient in soil organic carbon.
WP11T	SekbsdrSRm (8-10%)	Sandy surface and subsurface soils having low cation exchange capacity, low nutrients reserves,
		calcareous reaction, salinity. Soils with dry conditions, gravels, erosion risk and deficient in soil
		organic carbon with steep slope.
	SekbsdrSRm (10-12%)	Sandy surface and subsurface soils having low cation exchange capacity, low nutrients reserves,
		calcareous reaction, salinity. Soils with dry conditions, gravels, erosion risk and deficient in soil
		organic carbon with steep slope.
WPIIB	Sekbsdrm (1-2%)	Sandy surface and subsurface soils having low cation exchange capacity, low nutrients reserves,
		calcareous reaction, salinity. Soils with dry conditions, gravels and deficient in soil organic carbon.
	Sekbsdrm (2-4%)	Sandy surface and subsurface soils having low cation exchange capacity, low nutrients reserves,
WD10CUC	\mathbf{C} -1-h - d-m $(1, 20/)$	calcareous reaction, salinity. Soils with dry conditions, gravels and deficient in soil organic carbon.
WP12GUS	Sekbsdrm(1-2%)	Sandy surface and subsurface soils having low cation exchange capacity, low nutrients reserves,
	Salah drama (1, 20)	calcareous reaction, salinity. Soils with dry conditions, gravels and delicient in soil organic carbon.
	Sekbullii(1-2%)	Sandy surface and subsurface sorts having low carbon exchange capacity, low numeric serves,
WD12US	Salthedrm (1.20%)	Calcaleous feaction. Sons with dry conditions, gravels and deficient in son organic carbon.
WP1505	SekOsullii(1-2%)	salicy surface and subsurface soils naving low cation exchange capacity, low nutrients reserves, calcareous reaction salinity. Soils with dry conditions, gravels deficient in soil organic carbon
	$S_{alth} d_{max}(1, 20/)$	calcaledus reaction, samitly. Sons with dry conditions, graves derictent in son organic carbon.
	Sekbullit(1-2%)	sandy surface and subsurface sorts naving low cation exchange capacity, low nutrients reserves,
		carcareous reaction. Sons with dry conditions, gravels deficient in son organic carbon.





227 **3.3 Suggestive plausible soil managements:**

Now, there is a raised question i.e. at what time scales are FCC attributes refer todays, months, years, decades or centuries? And hence the scientific management technologies can be applied for mitigating these constrains. Experience in using FCC indicates that some of the condition modifiers can be changed with management at different time scales. In the current study, the possibility of overriding constrains is presented in table <u>6</u>.

234

235 236

Table 6. The temporal scale dimension of FCC attributes

FCC attribute	Can be ch	anged by mai	nagement with	Means of change	
	<1	1-10	10-100	>100	
Type/substrata					inherent, unless severely eroded
type - S					
High leaching					inherent
potential- e					
Low nutrient				\checkmark	inherent
reserves -k					
Calcareous -b					by sustained leaching in
					slightlycalcareous ones
Saline - s					by effective leaching
Sodic - n					by effective leaching
Soil moisture	\checkmark				temporarily by irrigation
stress - d					
Gravels- r				V.	inherent, unless severely eroded
High erosion - SR				\checkmark	inherent; can be mitigated by
					soil conservation practices
Low organic		\checkmark			by organic input application rates
matter - m					that exceed decomposition rate

237 From the previous table, some of the soil constrains cannot be changed in less than century

238 (inherent) such as type/substrata type, high leaching potential, low nutrient reverses, gravels

and high erosion risk. Whereas, condition modifiers can change at the decadal scale (10-

100 years) include calcareous reaction by sustained irrigation and subsequent leaching,
salinity and sodicity by applying effective leaching and low level of organic matter which can
be maintained under certain levels by supplying soil with different rates and sources of
organic inputs. The soil water stress can be managed by applying the water through
irrigation using the effective method of application such as trickle irrigation [18]. Some soil
management considerations are mentioned hereunder:

246 3.3.1.Low organic matter (m) and low nutrient reserves (k)

Low organic matter content which is prevailing in all soil profiles can be improved through application of organic manure, green manuring, mulching, crop rotation and so on. Also base saturation can be improved by applying fertilizers and amendments. Use of nitrogen and phosphorus fertilizers to mitigate major nutrient deficiencies is a must.

252 3.3.2.Salinity (s)

This can be removed by applying leaching and supplying the affected area with efficient drainage system in case of good quality water. Whereas, if the quality of irrigation water is poor due to either high salinity or high alkalinity or both, some suggestive management plans can be adopted such:

(1) In case of saline area and high salinity irrigation water, subsurface drainagesystem is a useful tool for desalinization.

(2) In case of saline area and high sodic irrigation water, subsurface drainage
system along with application of gypsum could be used for improving the
productivity. The gypsum amount to be added is determined by quality and quantity
of water to be added per year by applying the simple equation [19]:

$$GR = ((RSC - 2.5) \times N \times 36)$$

263 264 Where:

265 GR: Gypsum requirement (tons/acre), RSC: Residual sodium carbonate, N: number 266 of irrigations.

267 Thus, for soils irrigated with water having RSC 10.9, 10.4, 8.4 and 5.5 me/l and

268 needing 5 irrigations, the GR will be 1512, 1422, 1062 and 540 kg/acre.

269 3.3.3.High ESP soils (n):

Application of gypsum to soils along with deep ploughing and subsurface drainage is recommended. GR can be calculated by using the following equation:

$$GR(tons / ha) = \frac{(ESP_I - ESP_F) \times CEC \times 25.8 \times P}{(ESP_I - ESP_F) \times CEC \times 25.8 \times P}$$

100

272 273

Where
ESPI: Exchangeable Sodium Percentage initial (ESP) of soil, ESPF: ESP final, CEC: Cation
exchange capacity of soil, P: purity factor of gypsum.

276 **3.3.4.High erosion (SR)**

277

279

278 Following measures are suggested to reclaim high erosion land:

- Leveling and construction of contour bunds.
- Pipe outlets or ramps with suitable grasses for draining excess run off.
- Perennial vegetation like fuel, fodder trees and grasses may help
- 282 effectively to conserve the soil.
- 284 3.3.5.Rocky and guarried (r):

285

287

- 286 Following measures can be adopted
 - Enclosures of the hilly area with barbed wire.

- 288 Prohibition of grazing.
- 289 Locally suited tree species may be grown to conserve soils.
- 290 • Rehabilitation of quarry lands- plantation of suitable tree species.

4. CONCLUSION 292

293

291

294 According to the results, the major landforms of the studied area were described as Wadi 295 Bottom (WB), Bajada (B), Alluvial Fans (AF), Tableland (T), Gently Undulating Sand Sheet 296 (GUS) and Undulating Sand Sheet (US). The results of FCC units of WP11WB, WP11AF, 297 WP11B and some parts of WP12GUS and WP13US were classified as Sekbsdrm (1-2%) 298 only an area of 3125 feddan of WP11B was classified as Sekbsdrm (2-4%). This implies 299 that these map units have sandy (S) soils at both top and subsoils. The soils also have 300 constraints of high leaching potential (e), low nutrients reserve (k), basic reaction (b) and 301 salinity (s). As the soil exhibit ustic or xeric soil moisture regime, the Soil moisture stress 302 constraint (d) has been recognized. The other modifiers are because of gravels content (r) 303 and low organic matter (m). The soils of WP11T were classified as SekbsdrSRm (8-10%) 304 and SekbsdrSRm (10-12%) which having the same condition modifiers but different slope 305 grade. These soils are characterized by a high risk of soil erosion (SR) that erosion can 306 negatively affect plant productivity and ecosystem functions. The FCC unit Sekbdrm (1-2%) 307 has been found in some areas belongs to WP12GUS and WP13US. By following the 308 scientific technologies, the fertility constrains can be improved. 309 Authors' Contributions

310

311 CONSENT (WHERE EVER APPLICABLE) 312

313 ETHICAL APPROVAL (WHERE EVER APPLICABLE)

315 REFERENCES

316

314

317

318 Reference to a journal:

- 1. Udo BY, Utip U, Kufre EI, Monday T, Idungafa MA. Fertility Assessment of Some Inland 319 320 Depression and Floodplain (Wetland) Soils in Akwa Ibom State. J of Tropical Agri, Food, Env 321 and Extension. 2009;8(1):14-19.
- 322 2. Yakubu AA, Ibrahim SA, Ojanuga AG, Singh A. Fertility Capability Classification of Soils 323 of the Sokoto-Rima Flood Plain, Nigeria. Nig. J. Soil and Env. Res. 2012;10: 20: 63 – 70.
- 324 3. Tabi FO, Ngobesing ESC, Yinda GS, Boukong A, Omoko M, Bitondo D, Mvondo Ze AD
- 325 .Soil fertility capability classification (FCC) for rice production in Cameroon lowlands. African 326 J of Agri.Res. 2013;8(119):1650-1660.
- 327
- 4. Awad AA. Integration of Remote Sensing, Geophysics and GIS to evaluate groundwater 328 potentiality - A Case study in Sohag region. Egypt.Geology Department, Fac. of Science, 329 Assiut University, Assiut, Egypt. 2008.
- 330 5. EGMA. Egyptian Meteorological Authority database. 2004.
- 331 6.Global Land Cover Facility (GLCF). http://glcf.umd.edu/data/landsat/. 4321 Hartwick 332 Building · College Park, Maryland 20740. 2014.
- 333 7. Dobos E, Norman B, Bruce W, Luca M, Chris J, Erika M.The Use of DEM and Satellite Data forRegional Scale Soil Databases. 17th World Congress of Soil Science (WCSS),
- 334 335 Bangkok, Thailand. 2002; 649:14-21.
- 8. FAO. Guidelines for SoilDescription.Fourth edition, Food and Agriculture Organization of 336 337 the United Nations, Rome, Italy. 2006.

338 339 340 341 342 343	 USSL Staff. Diagnosis and improvement of saline and alkali soils. Agriculture Handbook 60, Richards LA (ed.). USDA: Washington, DC. 1954. Jackson ML . Soil Chemical Analysis.Prentice Hall of India Pvt. Ltd, New Delhi;1973. Black CA. Methods of soil analysis.2nd edition. Chemical and microbiological properties. Agronomy series no. 9, ASA, SSSA, Madison, Wis., USA. 1982. Soil Survey Staff. Keys to Soil Taxonomy, 9th edition, USDA National Resource
344	Conservation Services. 2003.
345 346 347 348 349 350 351 352 353 354	 Elbersen GWW, Catalan R. Portable Computer in Physiographic Soil Survey Unit.Proc. Inte. Soil Sci.,Cong. Homburg.1987. Buol SW, Sanchez PA, Cate RB, Granger MA. Soil fertility capability classification: A technical soil classification system for fertility management. In: Bornemisza, E, and A. Alvarado (Eds), Soil management in tropical America. N. C. State University, Raleigh, N. C. 1975; 126-145. Sanchez PA, Couto W, Buol SW. The Fertility Capability Classification System (FCC): Interpretation, applicability and modification. Geoderma. 1982; 27: 283 – 309. ESRI. ArcGIS Desktop: Release 9.3. Redlands, CA: Environmental Systems Research Institute 2009.
354 355 356	17. Mustafa AA, Negim OE. Geomatics Based Soil Mapping of The Eastern Desert Part of Sohag Governorate, Egypt. J. Soil Sci. and Agric. Eng., Mansoura Univ. 2015;6 (12): 1527 –
357 358 359 360 361 362 363 364	 1543. 18. Mustafa A A. Integration of parametric approach and GIS for optimum irrigation method in soils of Eastern desert part of Sohag, Egypt. The 12th International Conference of The Egyptian Society of Soil Science, 7-9 March, 2016, Ismailia, Egypt.2016. 19. Rajput RP. Potentials and utilization of Brackish water in irrigation management. In Proceeding of the training on "Diagnostic and improvement of problematic soils", Aug 18 to Sep 16, 2000), India.2000.
365	DEFINITIONS, ACRONYMS, ABBREVIATIONS
367 368 369	GIS FCC DEM
370 371	SRTM WB
372	AF
373	GUS
374	US
375	FCC UNITS
3/6	