

Antimicrobial effects of four plant extracts against post harvest spoilage fungi of yam (*Dioscorea rotundata* Poir).

¹Anukwuorji, C.A., ²Chukwuma, Maureen, Obianuju, ³Ezebo, R.O. and ³Anuagasi, C.L.

¹Department of Applied Biology and Biotechnology, Enugu State University of Science and Technology, P.M.B. 01660 Enugu State, Nigeria.

²Department of Biology, Nwafor Orizu College of Education, Nsugbe.

³Department of Botany, Nnamdi Azikiwe University, P.M.B. 5025, Awka, Anambra State.

Corresponding author. E-mail: dozygreat2k2@yahoo.co.uk

ABSTRACT

This study investigated the effects of plant extract on fungal pathogens responsible for yam rot in storage. Among the eight fungal pathogens isolated from yams with symptoms of post harvest rot, the inhibitory effects of four plant materials on four of these organisms (*Fusarium solani*, *Aspergillus niger*, *Botryodiplodia theobromae* and *Rhizopus stolonifer*) with the highest prevalence were examined. Phytochemicals test of these plant materials showed the presence of alkaloid, flavonoid, glycosides, saponin and tannins at different quantities. The pathogenicity test revealed that all the organisms tested were pathogenic on healthy yam tubers with *Aspergillus niger* being the most pathogenic. All the plant extracts inhibited the growth of the test organisms at varying degrees. The degree of inhibition was dependent on concentration of extract, extraction medium and the test organism. The highest inhibitory values were obtained from ethanol extracts of *Moringa oleifera* and *Azadirachta indica* at 7.5% and 10.0% concentration each, while *Gongronema latifolium* and *Xylopia aethiopicum* gave lower inhibitory values. This suggests that *Moringa oleifera* and *Azadirachta indica* are good bio killers and their biological active ingredients can be exploited for the control of yam rot.

Key words: Yam, Plant extracts, Fungi, Post harvest,

INTRODUCTION

Yam (*Dioscorea rotundata*) of the family Dioscoreaceae is a perennial herb cultivated for the consumption of its starchy tuber (Ezeibekwe *et al.*, 2009). Yam is one of the highly rated staple food crops of the tropical world and serves as an important carbohydrate food for millions of people in the tropical and subtropical countries which includes but not limited to West Africa, Caribbean, China, Malaysia, Oceania and Japan (Okigbo and Ikediugwu, 2000; Okigbo and Nmeko, 2005). Yam is of great importance to the economy of Nigeria as food and medicine (Okigbo and Ogbonna, 2006) hence FAO (2008) recorded that Nigeria produces 66.6% of total World's yam production every year. Yam contributes significantly to food security and its

availability in the market for a considerable part of the year helps prevent food shortages because it stores relatively longer than other food crops (Opara and Nwokocho, 2015).

According to FAO (2008) more than 25% of yams produced in Nigeria every year are lost to various kinds of diseases and pests. Deterioration of cultivated yam usually starts in the field from seedling stage, through harvesting and progressed in storage, which occur when infected tubers do not have any sign of external symptoms (Amusa *et al.*, 2003; Okigbo and Ogbonnaya, 2006). Yams are subjected to several diseases and these disease causing agents reduce the quantity and quality of yam by making them unmarketable and unappealing to the consumer (Ezeibekwe *et al.*, 2009). Some of the fungi associated with post harvest rot of yam and tuber crops are: *Aspergillus flavus*, *Aspergillus niger*, *Ervinia caratovora*, *Botrydiopodia theobromae*, *Fusarium solani*, *Fusarium oxysporium*, *Trichoderma viride*, *Rhizopus nodosus* *etc* (Adeniji, 1970; Aidoo, 2015; Anukwuorji, *et al.*, 2013; Ogundana *et al.*, 1970; Okigbo and Ikediugwu, 2000; Okigbo 2004).

Over the years, the use of chemicals has helped to control rot of root and tuber crops but not without challenges. With the constant use of chemicals, the target organisms can develop resistance, the use of chemical can also result to death by accumulating in man, poison and concentrate in food chain as they are usually not eco-friendly. Consequently, it is necessary to search for dependable and sustainable remedies that regard the requirement of man and its environment, hence plant extracts have been successfully used as viable alternative to control diseases in plants because they are relatively cheap, readily available and easily biodegradable (Amadioha, 2000; Amadioha and Obi, 1999; Anukwuorji, *et al.*, 2012; Ijato, 2011; Okigbo and Nmeka, 2005; Olufolaji, 1999). In other words, it is pertinent to find new types of antifungal for root and tuber crops which are not synthetic chemical; hence this research is on the antifungal activity of *Gongronema latifolium*, *Moringa oleifera*, *Xylopia aethiopicum* and *Azadirachta indica* on the primary organisms that induce rot of yam.

MATERIALS AND METHODS

Sources of Materials.

Yam (*Dioscorea rotundata*) tubers that show symptom of rot were obtained from Garriki Market in Enugu State. Healthy yam tubers for pathogenicity test were also purchased from the same market. They were packed into a polythene bag and taken to the laboratory for analysis. *Gongronema latifolium*, *Azadirachta indica*, *Moringa oleifera* and *Xylopia aethiopicum* were collected from Agbani Village.

Plant Extract Preparation

The plant materials were closely examined for the presence of extraneous materials and deteriorated leaves were removed. The sorted ones were first washed in running water and then

in sterile distilled water. They were allowed to drain and then sun-dried separately for five days. Each of the plant materials was grounded separately and sieved in 1 mm diameter sieve to obtain a powdered processed sample used for the extraction. For ethanol extract preparation, 20 g of each ground leaf material was mixed (separately) with 100 ml of 70% ethanol in a 500 ml beaker. The extract was sieved through four layers of sterile cheese cloth. Different concentrations of 10, 25, 50 and 100% were prepared. The same process as for ethanol was followed for aqueous extract preparation.

Isolation, Purification, Characterization and Identification of Fungal Pathogens of Yam

Yam tubers with symptoms of rot were washed in tap water. They were then cut into 3mm diameter at the boundary area between healthy and rotten portion of the yam tuber. The cut tissues were surface sterilized with 70% ethanol and rinsed twice, then dried on sterile tissue paper and four sections of the sterilized tissue pieces plated out on PDA. The inoculated plates were incubated for five days at a room temperature and observed daily for emergence of colonies. When growth has established, subcultures were prepared by transferring hyphal tips from the colony edge to fresh plate of PDA with the aid of flame sterilized blade. The resulting pure cultures were used for characterization and subsequent identification of the isolates using identification guide and compound microscope.

Antifungal sensitivity of plant extracts on fungal growth (*in vitro*)

The effect of extracts on fungal growth was determined using the growth inhibition test *in vitro*. Before dispensing PDA into each of the plates, four equal sections were created on each Petri-dish by drawing two perpendicular lines on the reverse of the plate, the point where the two lines meet is taken to be the center. One ml from each of the concentrations was dispensed into 9ml of melted PDA in a Petri dish, shaken together and allowed to solidify; this gave rise to 2.5%, 5.0%, 7.5% and 10% extract concentrations. Four millimeter diameter mycelia disc obtained from the colony edge of 7-day old pure cultures of each of the four test fungi was inoculated at the center of the plate containing the PDA-extract mixture. Negative control experiments were set up without the addition of any plant material while positive control PDA plates were added 1 ml of 0.5g Mancozeb. Three replicates were setup for each treatment and all the plates were incubated at 28 ± 2 °C for five days. Percentage inhibition was calculated from the data obtained using the formula below:

$$\text{Growth inhibition (\%)} = \frac{DT-DC}{DC} \times \frac{100}{1}$$

Where DC – Average Diameter of control and
DT – Average Diameter of fungal colony
With treatment

Pathogenicity Test

Four test fungi (*Fusarium solani*, *Aspergillus niger*, *Botryodiplodia theobromae* and *Rhizopus stolonifer*) isolated from the rotten yam tubers were tested on their ability or otherwise to induce rot on healthy yam tubers. The method of Okigbo and Ikediugwu (2000) was adopted. Fresh healthy yam tubers were washed with tap water and distilled thereafter sterilized with 70% ethanol. Cylindrical 1cm deep was removed aseptically from the healthy yam tuber with the aid

of sterile 5mm cork borer and then a core sample of the isolate from a 7days old PDA pure culture was inserted into the hole created and was closed back. Part of the tuber removed earlier was cut off to compensate for the thickness of the agar inoculum. The replaced core at the point of inoculation was sealed with sterile petroleum jelly; the same process was done for the control but without any pathogen. All the inoculated tubers were incubated over a period of 7-14 days at the temperature of 28-30 °C; they were checked daily for signs of rot. At the end of the 14days incubation period, the tubers were cut open along the point of each inoculation to expose the inner portion and the extent of rot examined, measured and recorded.

Phytochemical analysis of plant materials:

The presence of biological active ingredients in the four plant materials (*Gongronema latifolium*, *Azadirachta indica*, *Moringa oleifera* and *Xylopia aethiopicum*) was investigated using different standard methods as described by Edeoga *et al*, (2005); Oluduro (2012). The presence and quantity of alkaloid, saponin, tannins, flavonoid and glycoside was tested.

Statistical analysis:

The experimental design used was Complete Randomized Design (CRD) with three replicates. Data collected were subjected to Analysis of Variance (ANOVA) and means were separated using Duncan letters at 0.05 probability level to indicate the level of significance between values.

RESULTS

Isolation, characterization and Identification of fungi pathogens from rotten yam tubers

The following fungi (*Fusarium solani*, *Fusarium oxysporium*, *Aspergillus niger*, *Aspergillus flavus*, *Botryodiplodia theobromae*, *Rhizopus stolonifer*, *Rhizoctonia solani* and *Penicillium oxalicum*) were frequently isolated from rotten yam tubers. There was remarkable variation in their frequency of occurrence, *A. niger* with the frequency of 83.3% being the highest while *Rhizoctonia solani* (8.3%) being the least occurred (Fig. 1).

Pathogenicity Test

All the test organisms (*Fusarium solani*, *Aspergillus niger*, *Botryodiplodia theobromae* and *Rhizopus stolonifer*) were pathogenic to healthy yam tubers after 14 days of inoculation. The most pathogenic was *Aspergillus niger* (63%) while the least virulent was *Fusarium solani* (16%) (Fig. 2).

Effects of plant extracts at different concentrations on the growth of four test organisms *in vitro*

All the plant extracts showed varying degrees of inhibition on the test fungi, this was dependent on the concentration of the extract. With respect to ethanol extract, *Moringa oleifera*

at 10% extract concentration showed the highest inhibitory effect of 74.00 ± 4.583 on *Fusarium oxysporium*, this was significantly higher than other interactions, while the least inhibitory effect was recorded from *Gongronema latifolium* at 2.5% extract concentration. The value observed from the positive control was greater than all other interactions (Table 1).

On the effects of ethanol plant extracts on the inhibition of mycelia growth of *A.niger*, *Moringa oleifera* at 7.5% and 10% extract concentrations showed the highest inhibitory effect of 76.67 ± 7.024 and 76.33 ± 9.452 respectively, these were significantly better than other interactions with the exception of the positive control (82.00 ± 13.115) (Table 2).

For *Penicillium*, ethanol extract of *Moringa oleifera* and *A. indica* each at 10% extract concentration with the inhibitory effects of 72.00 ± 6.245 was the highest while the least inhibitory potent is *Xylopium aethiopicum* at 2.5% extract concentration (Table 3).

The highest inhibitory effect of 77.33 ± 2.083 recorded from ethanol *Moringa oleifera* extract at 7.5% extract concentration was the highest even though not significantly different from its inhibitory effect at 10% extract concentration and the positive control with 75.33 ± 6.110 and 76.00 ± 14.000 inhibition respectively (Table 4).

For aqueous extracts on the inhibition of mycelia growth of *Fusarium oxysporium*, the significantly better inhibitory effect was recorded from *Moringa oleifera* at 10% concentration and *A. indica* at 7.5% and 10% with inhibition percentages of 48.33 ± 7.234 , 46.00 ± 5.875 and 46.00 ± 7.211 respectively (Table 5).

Aqueous extract of *Moringa oleifera* at 7.5% and 10% concentration and *A.indica* at 7.5% and 10% depicted the highest inhibitory effect of 47.67 ± 3.215 , 48.33 ± 6.073 , 44.00 ± 9.539 and 47.00 ± 6.245 respectively. These were significantly higher than other interactions. The least effect of 0.67 ± 1.155 and 1.00 ± 1.732 was recorded from *Gongronema latifolium* and *Xylopium aethiopicum* at 2.5% extract concentration each (Table 6).

Table 7 clearly depicted that *Moringa oleifera* and *Azadirachta indica* all at 7.5% and 10% aqueous extract concentration had the best suppressive effect on the mycelia growth of *Penicillium spp* after 7 days of inoculation. *Gongronema latifolium* and *Xylopium aethiopicum* all at 2.5% extract concentration gave the least effect of 3.00 ± 0.000 and 4.00 ± 0.928 respectively. These were significantly lower than other interactions.

Inhibitory effects of aqueous plant extracts on the mycelia growth of *Rhizopus spp* followed the same trend as table 8 and 9. The least inhibitory effects of 3.67 ± 0.152 and 4.67 ± 0.686 were recorded from *Gongronema latifolium* and *Xylopium aethiopicum* all at 2.5% extract concentration, while the highest inhibitory effects were recorded from *Moringa oleifera* and *Azadirachta indica* all at 7.5% and 10% extract concentration. The positive control proved more potent with 76.00 ± 14.000 inhibition (Table 8).

Phytochemical Analysis

Quantitative analysis showed that *A. indica* contains abundant of Alkaloid and saponin while *M.oleifera* was abundantly rich in tannins. Alkaloid and tannins were absent in *X.aethiopicum* and *G.latifolium*. Glycoside is only reasonably present in *M. oleifera*.

Qualitative analysis revealed that *M.oleifera* was highest in alkaloid (5.31%), *G. latifolium* being the highest in saponin (19.24%), the highest quantity of tannins was observed in *A. indica* (19.24%). *G. latifolium* recorded the highest percentage of flavonoid (12.16%) while the only reasonable quantity of glycoside was found in *M. oleifera* (0.50%).




Fig. 1. Percentage occurrence of isolated fungi pathogens on rotten yam tubers




Fig. 2: Mean Percentage of Rot (Pathogenicity Test) on Healthy Yam Tubers

Table 1: Inhibitory effects of ethanol plant extracts at different concentrations on the growth of *Fusarium oxysporium* at day 7 after inoculation

Plant materials and control	Concentrations (%)			
	2.5	5.0	7.5	10.0
+ve Control	83.00±6.083f	83.00±6.083e	83.00±6.083d	83.00±6.083d
<i>G. latifolium</i>	4.67±0.163b	7.67±1.807b	35.00±5.568b	36.67±5.275b
<i>X.aethiopicum</i>	6.00±1.732c	6.33±1.970b	36.67±5.85b	37.33±3.215b
<i>M. oleifera</i>	45.00±5.000e	46.67±5.275d	69.67±5.508c	74.00±4.583d
<i>A.indica</i>	25.00±5.000d	28.67±2.767c	60.67±8.021c	61.67±7.638c
-ve Control	0.00±0.000a	0.00±0.000a	0.00±0.000a	0.00±0.000a
P-value	0.000	0.000	0.000	0.000

Results are in Mean ±Std

The same letter in a column is not significantly different

Table 2: Inhibitory effects of ethanol plant extracts at different concentrations on the growth of *Aspergillus niger* at day 7 after inoculation

Plant materials and control	Concentrations (%)			
	2.5	5.0	7.5	10.0
+ve Control	82.00±13.115e	82.00±13.115e	82.00±13.115f	82.00±13.115e
<i>G. latifolium</i>	2.33±0.517b	9.67±2.082b	30.33±4.240b	31.67±5.408b
<i>X.aethiopicum</i>	5.67±0.658c	10.00±2.292b	40.33±2.517c	37.00±6.083b
<i>M. oleifera</i>	32.67±0.429d	51.67±3.786d	76.67±7.024e	76.33±9.452d
<i>A.indica</i>	32.67±2.517d	35.67±5.452c	49.00±5.568a	57.67±5.774c
-ve Control	0.00±0.000a	0.00±0.000a	0.00±0.000a	0.00±0.000a
P-value	0.000	0.000	0.000	0.000

Results are in Mean ±Std

The same letter in a column is not significantly different

Table 3: Inhibitory effects of ethanol plant extracts at different concentrations on the growth of *Penicillium* spp at day 7 after inoculation

Plant materials and control	Concentrations (%)			
	2.5	5.0	7.5	10.0
+ve Control	86.67±10.732d	86.67±10.732d	86.67±10.73d	86.67±10.732d
<i>G. latifolium</i>	16.00±4.000b	17.00±2.646b	37.67±4.041b	35.00±3.606b
<i>X.aethiopicum</i>	14.00±1.464b	13.00±3.606b	34.00±5.292b	36.33±4.676b
<i>M. oleifera</i>	40.67±1.155c	43.33±4.041c	62.33±4.933c	72.00±6.245c
<i>A.indica</i>	38.00±2.646c	40.00±0.000c	60.00±4.000c	72.00±6.245c
-ve Control	0.00±0.000a	0.00±0.000a	0.00±0.000a	0.00±0.000a
P-value	0.000	0.000	0.000	0.000

Results are in Mean ±Std

The same letter in a column is not significantly different

Table 4: Inhibitory effects of ethanol plant extracts at different concentrations on the growth of *Rhizopus* spp at day 7 after inoculation

Plant materials and control	Concentrations (%)			
	2.5	5.0	7.5	10.0
+ve Control	76.00±14.000e	76.00±14.000d	76.00±14.000d	76.00±14.000d
<i>G. latifolium</i>	10.67±3.786b	10.00±2.718b	35.00±2.646b	38.00±4.359b
<i>X.aethiopicum</i>	18.00±4.568b	18.33±1.528c	37.33±2.517	39.67±0.577b
<i>M. oleifera</i>	48.33±2.082d	51.00±8.933c	77.33±2.083d	75.33±6.110d
<i>A.indica</i>	47.33±3.786d	48.33±8.622c	69.00±1.539c	65.00±6.288c
-ve Control	0.00±0.000a	0.00±0.000a	0.00±0.000a	0.00±0.000a
P-value	0.000	0.000	0.000	0.000

Results are in Mean ±Std

The same letter in a column is not significantly different

Table 5: Inhibitory effects of aqueous plant extracts at different concentrations on the growth of *Fusarium oxysporium* at day 7 after inoculation

Plant materials and control	Concentrations (%)			
	2.5	5.0	7.5	10.0
+ve Control	83.00±6.083d	83.00±6.083f	83.00±6.083e	83.00±6.083e
<i>G. latifolium</i>	0.67±1.155b	2.67±0.517b	12.00±1.732b	12.67±1.015b
<i>X.aethiopicum</i>	1.33±1.155b	7.67±1.528c	22.00±2.646c	22.67±3.506c
<i>M. oleifera</i>	20.33±4.505c	26.33±4.163e	44.33±3.512d	48.33±7.234d
<i>A.indica</i>	17.00±4.359c	19.67±4.505d	46.00±5.875d	46.00±7.211d
-ve Control	0.00±0.000a	0.00±0.000a	0.00±0.000a	0.00±0.000a
P-value	0.000	0.000	0.000	0.000

Results are in Mean ±Std

The same letter in a column is not significantly different

Table 6: Inhibitory effects of aqueous plant extracts at different concentrations on the growth of *Aspergillus niger* at day 7 after inoculation

Plant materials and control	Concentrations (%)			
	2.5	5.0	7.5	10.0
+ve Control	82.00±13.115e	82.00±13.115f	82.00±13.115d	82.00±13.115d
<i>G. latifolium</i>	0.67±1.155b	3.67±0.041b	18.33±2.887b	18.00±6.557b
<i>X.aethiopicum</i>	1.00±1.732b	7.67±1.506c	15.67±4.041b	17.67±3.662b
<i>M. oleifera</i>	19.00±5.533d	37.33±5.504e	47.67±3.215c	48.33±6.073c
<i>A.indica</i>	13.00±4.937c	23.00±4.550d	44.00±9.539c	47.00±6.245c
-ve Control	0.00±0.000a	0.00±0.000a	0.00±0.000a	0.00±0.000a
P-value	0.000	0.000	0.000	0.000

Results are in Mean ±Std

The same letter in a column is not significantly different

Table 7: Inhibitory effects of aqueous plant extracts at different concentrations on the growth of *Penicillium spp* at day 7 after inoculation

Plant materials and control	Concentrations (%)			
	2.5	5.0	7.5	10.0
+ve Control	86.67±10.732e	86.67±10.732d	86.67±10.732d	86.67±10.732d
<i>G. latifolium</i>	3.00±0.000b	6.67±2.082d	18.33±2.887b	21.67±3.505b
<i>X.aethiopicum</i>	4.00±0.928b	5.67±2.371b	15.67±4.041b	21.67±3.970b
<i>M. oleifera</i>	18.67±3.234d	24.33±3.506c	47.67±3.215c	50.00±5.292c
<i>A.indica</i>	14.33±2.371c	24.67±4.238c	44.00±9.539c	47.00±8.888c
-ve Control	0.00±0.000a	0.00±0.000a	0.00±0.000a	0.00±0.000a
P-value	0.000	0.000	0.000	0.000

Results are in Mean ±Std

The same letter in a column is not significantly different

Table 8: Inhibitory effects of aqueous plant extracts at different concentrations on the growth of *Rhizopus spp* at day 7 after inoculation

Plant materials and control	Concentrations (%)			
	2.5	5.0	7.5	10.0
+ve Control	76.00±14.000d	76.00±14.000e	76.00±14.000d	76.00±14.000d
<i>G. latifolium</i>	3.67±0.152b	6.00±1.000b	20.33±3.215b	24.67±4.619b
<i>X.aethiopicum</i>	4.67±0.686b	14.33±7.024c	19.33±2.506b	26.67±5.686b
<i>M. oleifera</i>	22.67±4.933c	30.67±2.082d	46.33±2.327c	48.67±2.517c
<i>A.indica</i>	19.33±3.506c	30.00±0.817d	45.33±4.041c	46.33±5.686c
-ve Control	0.00±0.000a	0.00±0.000a	0.00±0.000a	0.00±0.000a
P-value	0.000	0.000	0.000	0.000

Results are in Mean ±Std

The same letter in a column is not significantly different

Table 9: Qualitative phytochemicals present in *X. aethiopicum*, *M.olerifera*, *G. latifolium* and *A. indica*

Phytochemicals	Plant materials			
	<i>X. aethiopicum</i>	<i>M.oleifera</i>	<i>G. latifolium</i>	<i>A. indica</i>
Alkaloid	-	+	-	+++
Saponin	+	++	++	++
Tannins	-	+++	-	+++
Flavonoid	++	++	++	++
Glycoside	-	+	-	-

Key: +++= Abundantly Present, ++= Moderately Present, += Fairly Present, -= Absent.

Table 10: Quantitative phytochemicals present in *X. aethiopicum*, *M.olerifera*, *G. latifolium* and *A. indica*

Plant materials	Phytochemicals (%)				
	Alkaloid	Saponin	Tannins	Flavonoid	Glycoside
<i>X. aethiopicum</i>	0.4	2.41	0	3.61	0
<i>M.oleifera</i>	5.31	13.86	15.46	7.30	0.50
<i>G. latifolium</i>	1.21	19.24	0.93	12.16	0

<i>A. indica</i>	4.06	13.86	19.42	8.31	0.005
------------------	------	-------	-------	------	-------

DISCUSSION AND CONCLUSION

Organisms that were consistently associated with yam tubers spoilage in this research were: *Aspergillus niger*, *Aspergillus flavus*, *Botrydiplodia theobromae*, *Rhizopus stolonifer*, *Rhizoctonia solani* and *Penicillium oxalicum*, this is in tandem with the reports of several scientists (Anukwuorji, *et al.*, 2013; Okigbo *et al.*, 2009; Ameinyo and Atanga 2007) as the organisms that are mainly prevalent in the post harvest deterioration of root and tuber crops such as potato, cocoyam, cassava etc. The most prevalent organism was *A. niger* (83.3%) followed by *B. theobromae* (75%) and *F. solani* (66.7%). This does not completely agree with the result of Okigbo *et al.*, (2009) who reported that *B. theobromae* was the most prevalent. Pathogenicity study revealed that all the test pathogens were pathogenic on healthy yam tubers (*in vitro*) with different degrees of virulence. The most virulent was *A. niger* (63%) while the least virulent was *F. solani* (16%), this agrees with the result of Anukwuorji *et al.*, (2013) and Okigbo *et al.*, (2014) on potato and yam respectively who reported that *F. solani* was the least virulent while *A. niger* was the most virulent. This is in sharp contrast with the reports of Okigbo and Emeka, (2010) who recorded that *B. theobromae* was the most virulent. Rot of root and tuber crops often starts in the field and progresses in storage (Ezeibekwe *et al.*, 2009). Pathogenic microorganisms mainly penetrates yam tubers through natural openings and sometimes through wounds that occur during harvesting, handling and transportation from field to barn and sometimes to market (Ameinyo and Ataga, 2007). The soils adhering to harvested tubers have been confirmed to contain a good quantity of microorganisms that could induce rot in the tubers (Okigbo and Ogbonna, 2006).

The antifungal potential of some plant materials in controlling different organisms pathogenic to food crops has been reported by some scientists such as Okigbo *et al.*, (2009), Anukwuorji *et al.*, (2013). Other biocontrol measures like the use of microbial antagonists such as *B. subtilis* and *T. viride* were also well documented (Okigbo and Ikediugwu, 2000; Okigbo, 2002; Okigbo and Emeka, 2010).

The inference of this research work depicted that the radial growth (*in vitro*) of the entire test organisms were greatly inhibited by all the plant extracts tested at varying degrees. This is an indication that fungitoxic compounds abounds in the plant materials. This agrees perfectly with the reports of Amienyo and Ataga (2007) and Anukwuorji, *et al.*, (2013) all on rot of potato tubers.

Results obtained from the effects of various plant materials showed that *M. olerifera* and *A. indica* at 7.5% and 10% concentration were better inhibitors than *G. latifolium* and *X. aethiopicum* at 2.5% and 5.0% extract concentration. This is in tandem with the documentation of Ramesh *et al.*, (2009) on deterioration of cassava who reported that *A. indica* significantly

controlled the rot inducing pathogens. The differences in the inhibition ability can be linked to the differences in the nature, quantity and quality of their biological active ingredient (Okigbo *et al.*, 2009; 2013; Onifade, 2002), this suggest the potential of plant materials as alternative to synthetic chemicals.

Quantitative and qualitative phytochemical screening of test plants proved positive to most of the phytochemicals tested, with few exceptions. The presence of biological active substances have been showed to bestow resistance to plants against pathogenic microorganisms (Srinwasan *et al.*, 2001). Therefore the antimycotic potentials of these plants can be linked to the presence of different phytochemicals inherent in them (Okwu and Joshia, 2006). Pharmacological and medicinal values of these phytochemicals was proved by the documentations of several scientists (Okwu, 2004; Okigbo *et al.*, 2009).

The fact that about 25% of yam produced annually in Africa is lost due to rot is appalling; hence post harvest spoilage of yam should be of global concern. However, if Nigeria will still top the list of yam producing countries, then all stakeholders in Nigeria (farmers, government, research institutes, Non-governmental organizations etc) must be involved in finding a lasting solution to this menace. No doubt, plant extract is fast becoming a more reliable solution to the problem of rot; hence the demonstrated antifungal potential of these plant materials recommends them as natural fungicides. Therefore, to overcome the problem of yam rot and rot of other tuber crops, urgent effort and attention should be channeled towards harnessing the potential of these plant materials (extracts).

REFERENCES

- Adeniji, M.O. (1970). Fungi associated with storage decay of yam in Nigeria. *Phytopathology*, 60: 590-592.
- Aidoo, A.T. (2015). Bioactivity of *Zingiber officinale* and *Piper nigrum* plant extracts in controlling post-harvest white yam (*Dioscorea rotundata*) tuber rot fungi. *African Journal of Microbiology Research*. 9(22): 1499 -1503.
- Amusa, N.A., Adegbite A.A., Muhammed, S. and Baiyewu R.A. (2003). Yam diseases and their management in Nigeria. *African Journal of Biotechnology* 2(2): 33.
- Amadioha, A.C and Obi, V.I. (1999). Control of anthracnose disease of cowpea by *Cymbopogon citratus* and *Ocimum gratissimum*. *Acta Phytopathol Entomol. Hungarica*. 34(2): 85 – 89.
- Amadioha, A. C. (2000). Fungitoxic effects of some leaf extract against *Rhizopus Oryzae* . *Int. J. pest. Manage*. 52: 311 – 314.

Amienyo, C.A. and Ataga, A.E. (2007). Use of indigenous plant extracts for the protection of mechanically injured sweet potato (*Ipomoea batatas* (L) Lam) tubers. *Acad. J. Sci. Res. Essay*. 2: 167-170.

Anukwuorji, C.A., Anuagasi, C.L. and Okigbo. R.N. (2013). Occurrence and control of fungal pathogens of potato (*Ipomoea batatas* L. Lam) with plant extracts. *International Journal of pharmTech Research*. 2 (3): 273-289.

Anukwuorji, C.A., Ramesh, R. Putthi and Okigbo, R.N. (2012). Isolation of fungi causing rot of cocoyam (*Colocasis esculata* (L.) schott) and control with plant extracts: (*Allium sativum* (L.), *Garcinia kola*, heckle; *Azadirachta indica* L. and *Carica papaya* L. *Global Advanced Research Journal of Agricultural sciences* 1(2): 033 – 048.

Edeoga,, H.O., Okwu, D.E. and Mbaebie, B.O. (2005). Phytochemical constituents of some Nigerian medicinal plants, *African Journal of Biotechnology*, 4 (7): 685-688.

Ekwealor, K.U., Nnabude, P.C., and Anukwuorji, C.A. (2012). Effects of different irrigation levels and mulch rates on the performance of *Amaranthus* hybrids (H). *African Journal of Sciences*.

Ezeibekwe, I.O., Opara, M.I. and Mbagwu, F.N. (2009). Antifungal effect of Aloe-vera gel on fungal organisms associated with yam (*Dioscorea rotundata* Poir) rot. *Journal of Molecular Genetics* 1(1): 11-17.

F.A.O. (2008) Food and Agricultural Organization corporate statistical database.

Ijato, J. Y. (2011). Evaluation of antifungal effects of extracts of *Allium sativum* and *Nicotiana tobacum* against soft rot of yam (*Dioscorea alata*). *Researcher* 3(2):1-5.

Ogundana, S.K, Naqvi, S.H.Z and Ekundayo, J.A. (1970). Fungi associated with soft rot of yams in storage in Nigeria. *Trans. Br. Mycol . Soc.* 54:445 – 451.

Okigbo, R.N. (2002). Mycoflora of tuber surface of white yam (*Dioscorea rotundata* Poir) and post harvest control of pathogens with *Bacillus substilis*. *Mycopathologia* 156: 81-85.

Okigbo, R.N., (2004). A review of biological control methods for post harvest yams (*Dioscorea* spp.) in storage in south Eastern Nigerian KMITL. *Sci. J.*, 4(1): 207-215.

Okigbo, R.N., Anukwuorji, C.A. and Eguae, C.T. (2012). Control of microorganism causing the

- deterioration of yam chips with *Vernonia amygdalina* (L) and *Zingiber officinale* (L). *Nigerian Journal of mycology* 4:45 – 46.
- Okigbo, R.N., Anukwuorji, C.A. and Ezeabara, C.A. (2013). Effects of leaf extracts of *Chromolaena odorata* (L) and *Terminalia catappa* (L) on mycelia growth of yam tuber pathogens. *Niger. J. Plt. Prot.* 27 (1): 74-84.
- Okigbo, R.N., Anukwuorji, C.A. and Igweonu, M.Z. (2014). Fungitoxic effect of *Azadirachta indica* and *Vernonia amygdalina* extracts on *Aspergillus niger* and *Fusarium oxysporium* the causal organisms of yam (*Dioscorea rotundata*) rot. *The Bioscientist* 2(1): 70- 86.
- Okigbo, R.N. and Emeka, A.N. (2010). Biological control of rot-inducing fungi of water yam (*Dioscorea alata*) with *Trichoderma harzianum*, *Pseudomonas syringae* and *Pseudomonas chlororaphis*. *Journal of Stored Products and Postharvest Research* 1(2): 18-23
- Okigbo, R.N. and Ikediugwu, F.E.O. (2000). Studies on biological control of post harvest rot of yams (*Dioscorea spp*) with *Trichoderma viride* . *J. Phytopathol.* 148; 351 – 355.
- Okigbo, R.N. and Nmeka, A. (2005). Control of yam tuber with leaf extracts of *Xylopia aethiopica* and *Zingiber officinale*. *Afr. J. Biotechnol.* 4(8). 804 – 807.
- Okigbo, R.N. and Ogbonnaya, O.U. (2006). Antifungal effects of two tropical plant extracts (*Ocimum gratissimum* and *Aframomum melegueta*) on postharvest yam (*Dioscorea spp.*) rot. *Afr. J. Biotechnol.*, 5(9): 727-731.
- Okigbo, R.N. Ramesh Putheti and Achusi, C.T. (2009) Post-Harvest Deterioration of Cassava and its Control Using Extracts of *Azadirachta Indica* and *Aframomum Melegueta*. *E-Journal of Chemistry.* 6(4), 1274-1280.
- Okwu, D.E. (2004). Phytochemical and vitamin content of indigenous species of south eastern Nig. *J. Sustainable Agric and Environment.* 6(1): 30-37.
- Okwu, D.E. and Joshia, C. (2006). Evaluation of the chemical composition of two Nigeria medicinal plants. *Afri.J.Biotechnol.* 5(4): 13-16.
- Oluduro, A.O. (2012). Evaluation of antimicrobial properties and nutritional potentials of *Moringa oleifera* Lam. leaf in South Western Nigeria, *Malaysian Journal of Microbiology*, 8(2), 2012, 59-67.
- Olufolaji, D.B. (1999). Control of wet rot disease of *Amaranthus spp* caused by *Chonehora curcubitaricum* with extracts of *Azadirachta indica*. *J.Sust. Agric. Environ* 1(22): 183-190
- Onifade, A.K., 2002. Antifungal effect of *Azadirachta indica* A. Juss extracts on *Collectotricum*

lindemathianum. *Global J. Pure. Appl. Sci.*, **6**(3): 423-428.

Opara, E. U. and Nwokocha, N. J. (2015). Antimicrobial activities of some local medicinal plants against post harvest yam rot pathogens in humid south eastern Nigeria. *Journal of Microbiology Research and Reviews*. 3(1): 1-9.

Ramesh, P. P. N.P, Singh, A.B., Ramana, S. and Subba, R.A. (2009). Impact of organic –manure combinations on the productivity and soil quality in different cropping systems in central India. *J. Plant. Nutr.Soil. Sci.*172:577-585.

Srinwasan, D.,Perumalsamy, L.P. and Nathan, S.T. (2001). Antimicrobial activity of certain Indian medicinal plants used in folkloric med. *J.ethnopharm.*94:217-222.