

Original Research Article**Effect of Wastewater and *Moringa oleifera* Leaf Extract Irrigation on Mineral Composition of Maize (*Zea mays* L) Plant in a Pot Experiment****ABSTRACT**

A study was conducted in the screen house at the FAO/TCP teaching and research farm of Adamawa state University, Mubi to assess the effect of wastewater from fish pond and *Moringa oleifera* leaf extract on the mineral composition of maize plant. Results showed that the concentration of minerals in maize plant were in deficient range for P (0.11-0.12%), Ca (0.12 – 0.22%), Cu (1.65 – 2.55 mg/kg), and Mn (17.5 – 19.0 mg/kg) in all irrigations except K (2.05 – 3.02%) that was in sufficient range. Wastewater irrigation had Mg (0.70%) and Zn (31.93 mg/kg) composition in sufficient range and Fe (45 mg/kg) in deficient range. *Moringa* leaf extract irrigation had maize plant composition of Fe (66.7 mg/kg) in sufficient range while combined wastewater and *Moringa* leaf extract irrigation had Mg (0.31%), Fe (51.67 mg/kg) and Zn (33.60 mg/kg) in sufficient range. Irrigating maize plant with wastewater + *Moringa* leaf extract improved P, K and Zn composition by 9.1, 43.8 and 5.2%, respectively. Regression and correlations analyses indicate that Na concentration in the soil is the element affecting greater number of elements composition in the plant and subsequently the growth Characters. Therefore, wastewater from fish pond and *Moringa* leaf extract irrigation either in combination or separately did not increase the nutrient elements and heavy metals to toxic limits and therefore can be used for irrigation purposes especially maize crop.

Key words: Wastewater, *Moringa* leaf extract, mineral composition, maize, irrigation

1. INTRODUCTION

24 With the increasing global population, the gap between supply and demand for water
25 is widening and has reached an alarming level such that in some part of the world especially
26 in the arid and semiarid regions, it is posing a threat to human existence. In these regions,
27 waste water is considered a valuable source of irrigation water and a fertilizing material [1].
28 However, there is an increasing awareness on the need to dispose waste water safely and
29 beneficially which is of paramount importance bearing in mind environmental health and
30 water pollution issues.

31 Most sewage wastes contain valuable nutrients that improve soil fertility and crop
32 production [2]. Therefore, the benefits of wastewater use in irrigation are numerous but
33 precautions must be taken to avoid short and long-term environmental risks. So, inappropriate
34 handling and management of wastewater reuse for irrigation can create serious environmental
35 and health hazards [3]. Such a proper use can relatively minimize pollution of the ecosystem
36 which otherwise would be contaminated by direct disposal of wastewater into surface or
37 ground water [4]. Wastewater contains essential nutrients for plant growth like N, P, K, iron
38 (Fe), zinc (Zn), manganese (Mn), and copper (Cu) and a considerable amount of organic
39 matter [4,5].

40 However, improper management of wastewater irrigation may provide the crops with
41 nutrients beyond their specific requirement and subsequently accumulate them at undesirable
42 high levels in the crop. This would lead to reduction in the yield and its quality [5,6].

43 The concentration of macro and micro nutrients in corn plants were affected by
44 chelate fertilizers [7]. It was reported by Lockman [8] that there were sufficient range for
45 different element in leaf and whole corn and sorghum at various stages.

46 One of the sources of foliar fertilizer that is attracting attention is *Moringa oleifera*
47 plant which is termed as ‘tree for life’ due to its importance. The efficacy of *Moringa* as
48 organic fertilizer has already proved by a number of field trials conducted in Nigeria.

49 Application of processed *M. oleifera* fresh leaves and seed cake as organic fertilizer improved
50 the yields of maize and sorghum yield [9].

51 The importance of waste water and *Moringa leaf extract* irrigation on the mineral
52 composition of maize plant needs to be assessed to give more understanding of the dynamics
53 of nutrients in plant and soil. Therefore this work was conducted to assess the effect of waste
54 water from fish pond and *Moringa* leaf extract on mineral composition of maize plant.

55 2. MATERIALS AND METHODS

56 The experiment was conducted in the screen house at the Food Agricultural
57 Organisation/Tree Crop Program (FAO/TCP) teaching and research farm of Adamawa State
58 University, Mubi, Nigeria.

59 The experiment was laid out in a complete randomized design comprising of three (3)
60 treatments; wastewater from fish pond (WW), *Moringa* leaf extract and wastewater +
61 *Moringa* leaf extract (WW+MLE). This was replicated three times. The experiment was done
62 in plastic pots measuring 25 cm diameter and 22 cm height. Top soil at 0-15 cm depth was
63 collected at Shuware along river Yadzaram. It was air dried and sieved through 2 mm screen.
64 The texture of the soil was sandy loam (54% sand, 28% silt and 18% clay), with pH of 7.3,
65 electrical conductivity of 799 μ /cm while the water holding capacity was 28.8%, Organic
66 matter and organic carbon were 4.01% and 2.33%, respectively, while available phosphorus
67 was 13.4 mg/kg. The amounts of exchangeable cations were 0.06, 1.0, 0.04 and 0.08 cmol/kg
68 for K, Na,

69 **Table 1: Some chemical properties of the experimental soil**

| Element | Value |
|-----------|-------|
| P (mg/kg) | 13.44 |

| | |
|--------------|-------|
| K (cmol/kg) | 0.06 |
| Ca (cmol/kg) | 0.084 |
| Mg (cmol/kg) | 0.042 |
| Na (cmol/kg) | 1.00 |
| Fe (mg/kg) | 1.35 |
| Cu (mg/kg) | 0.46 |
| Zn (mg/kg) | 3.064 |
| Mn (mg/kg) | 13.55 |
| Ni (mg/kg) | 0.03 |
| Pb (mg/kg) | 0.29 |
| OC (%) | 2.33 |
| OM (%) | 4.01 |
| EC (μ/cm) | 799 |
| pH | 7.25 |
| WHC (%) | 28.8 |

70

71 **Table 2: Chemical composition of irrigation water**

| Element | Wastewater from fish pond | <i>Moringa oleifera</i> leaf extract |
|-----------|------------------------------|---|
| P (mg/l) | 8.0 | 8.10 |
| K (mg/l) | 54 | 52 |
| Ca (mg/l) | 46.1 | 34.7 |
| Mg (mg/l) | 38.0 | 35.0 |
| Na (mg/l) | 192 | 223 |

| | | |
|-----------|------|------|
| Fe (mg/l) | Nd | 0.75 |
| Cu (mg/l) | Nd | 0.25 |
| Zn (mg/l) | 0.04 | 0.05 |
| Ni (mg/l) | Nd | 0.03 |
| Pb (mg/l) | Nd | 0.42 |
| EC (/cm) | 81.3 | 37.5 |
| pH | 7.8 | 7.0 |

72 Heavy metals like Fe, Cu, M n, Ni, and Pb were beyond limit of detection

73 Mg and Ca, respectively (Table 1). The concentration of elements in wastewater and fresh
74 leave of *Moringa oleifera* are presented in the Table 2.

75 Eight kg of the soil was placed in the plastic pots. The pots were irrigated to field capacity
76 and left over for three days to attain equilibrium. The experimental pots were kept in the
77 screen house. Eight seeds of maize were planted in each pot on 3rd of March 2014. The crops
78 were irrigated 3 times a week. The plant population was thinned to four plants per pot two
79 weeks after crop germination and was harvested after 52 days of planting.

80 The single acid digestion method for the analysis of plant tissues for P, K, Ca, Mg, Na, Fe,
81 Cu, Mn and Zn as described by Marr and Cresser [10] was followed.

82 Soil organic carbon was determined by wet oxidation method as described by Walkley and
83 Black [11]. The Titrimetric method for the determination of Calcium and Magnesium in the
84 soil as described by Black, [12] was followed. Soil pH was determined as described by Bates,
85 [13]. Hydrometer method of Soil Mechanical Analysis was followed as described by
86 Bouyoucos, [14]. Available P in soil was determined as described by Bray and Kurtz, [15] at
87 the wavelength of 660 nm. Potassium and sodium was determined in 1M neutral NH₄Ac soil
88 extract using flame photometry and exchangeable acidity as described by Mclean, [16]. The

89 Di-ethylene triaminepenta acetic acid (0.005M DTPA) [17] and 0.1N HCl [18] method of
 90 extracting available Zn and Mn in the soil was adopted. The determination of total iron in soil
 91 by atomic absorption spectrophotometry, after digestion with strong oxidizing acids and
 92 treatment with hydrofluoric acid to eliminate silica as described by Jackson [19] was
 93 maintained.

94 Waste water used for irrigation was analyzed as described in the Standard Methods for the
 95 Examination of Water and Waste water [20]. Fresh leaves of *Moringa oleifera* were prepared
 96 at the ratio of 1:36 as described by Mathur, [21]. The result of the concentration of different
 97 minerals in maize plant were compared with the data reported by Lockman [8] on corn.

98 The data was subjected to analysis of variance using SAS [22]. The significant difference
 99 among the mean was compared using Duncan Multiple Range Test.

100 **3. RESULTS AND DISCUSSION**

101 **3.1 Response of some growth characters of maize**

102 The effect of wastewater and *Moringa* leaf extract irrigation on the growth of maize plant is
 103 presented in Table 3. Plant height at *Moringa* leaf extract irrigation was superior to plant
 104 height recorded at wastewater + *Moringa* leaf extract and *Moringa* leaf extract. The result
 105 obtained agreed with Emmanuel *et al.*, [9] and Mvumi *et al.*, [23] who reported that organic
 106 manure from *Moringa* leaf extract increased plant height in maize. There were no significant
 107 ($P=0.01$) differences between wastewater and *Moringa leaf* extract application on number of
 108 leaves. However, combination of wastewater and *Moringa* leaf extract had number of leaves
 109 advantage of 5.3 and 8.6% over wastewater and *Moringa* leaf extract, respectively. Increase
 110 in the number of leaves of maize plant from irrigation with *Moringa* leaf extract has been
 111 reported by Emmanuel *et al.*, [9] while that of wastewater has been reported by [24]. Thus,

the combined effect of wastewater and *Moringa* leaf extract gave the number of leaf advantage over wastewater and *Moringa* leaf extract irrigated separately. Leave length shows that application of the treatment did not indicate any significant difference. This differs with the findings of Ramana *et al.* [25] who observed that influence of distillery wastewater on the physiology of maize increased leave length. Same response was recorded with the width of leaf. However, there were no significant differences between the treatments.

Table 3: Effect of wastewater and *Moringa* leaf extract on growth parameter of maize

| Treatment | Plant height (cm) | Number of leave | Leave length (cm) | Width of leave (cm) |
|-----------|----------------------|--------------------|----------------------|---------------------------|
| WW | 71.51±3.34b | 11.67±0.58a | 83.17±3.26a | 4.38±0.13a |
| MLE | 86.53±4.19a | 12.00±1.00a | 83.05±4.59a | 4.63±0.39a |
| WW+MLE | 69.60±7.28b | 12.67±0.58a | 87.18±1.72a | 4.78±0.20a |
| LSD | 10.70 | 1.49 | 6.79 | 0.53 |
| <i>P</i> | * | NS | NS | NS |
| CV (%) | 7.06 | 6.15 | 4.03 | 5.73 |

Figures in the same column followed by the same letter are not significantly different by

LSD_{0.05}

*- significant at 5% level of probability

CV-Coefficient of variation

127 **Table 4: Effect of wastewater and *Moringa* leaf extract on the mineral composition**
 128 **of maize plant.**

| Treatment | P | K | Ca | Mg | Na |
|----------------|-------------|------------|------------|------------|-------------|
| | ← | | % | → | |
| WW | 11.00±0.02a | 2.45±0.13 | 0.14±0.18a | 0.70±0.02a | 0.131±0.004 |
| | | b | | | a |
| MLE | 11.00±0.01a | 2.10±0.10 | 0.22±0.11a | 0.13±0.01b | 0.131±0.004 |
| | | b | | | a |
| WW+ML | 12.00±0.01a | 3.02±0.08a | 0.12±0.17a | 0.31±0.17b | 0.126±0.005 |
| E | | | | | a |
| LSD | 0.02 | 0.21 | 0.32 | 0.20 | 0.009 |
| P | NS | ** | NS | * | NS |
| R ² | 0.20 | 0.90 | 0.11 | 0.89 | 0.29 |

| | Fe | Cu | Zn | Mn | Ni | Pb |
|----------------|-------------|------------|-------------|-------------|------------|-----------|
| | ← | | mg/ kg | → | | |
| WW | 45.00±15.00 | 1.65±0.11 | 31.93±1.75a | 17.50±0.87b | 2.33±1.15a | 4.18±0.07 |
| | b | b | | | | a |
| MLE | 66.67±5.77a | 2.55±0.29a | 24.93±3.02 | 17.83a±0.29 | 3.33±0.58a | 3.61±0.07 |
| | | | b | b | | a |
| WW+ML | 51.67±7.64b | 2.00±0.58 | 33.60±2.39a | 19.00±0.87a | 3.33±2.08a | 3.61±0.03 |
| E | | b | | | | a |
| LSD | 20.50 | 0.59 | 0.78 | 0.54 | 0.14 | 0.25 |
| P | * | ** | * | * | NS | NS |
| R ² | 0.54 | 0.59 | 0.78 | 0.54 | 0.14 | 0.25 |

129 Means in the same column followed by the same letter are not significantly different by

130 LSD_{0.05}

131 *-significant at $P=0.05$

132 **- significant at $P=.01$

133 NS-not significant

134

135 **3.2 Mineral composition of maize plants.**

136 The effect of wastewater and Moringa leaf extract on mineral composition of maize plant is
137 presented in Table 4. Phosphorus concentration was highest at wastewater + *Moringa* leaf
138 extract with corresponding value of 0.12% while wastewater and *Moringa* leaf extract had
139 lower values of 0.11% phosphorus each. Similar results were reported by Sadiq and Hussain
140 [7]

141 who found varying concentrations and responses by maize for the uptake of different element
142 receiving fertilizer. Similarly, wastewater + *Moringa* leaf extract application had the highest
143 K concentration of 2.20% significantly ($P=.01$) higher than K concentration from *Moringa*
144 leaf extract (2.10% and wastewater (2.05%). Therefore, the combination of wastewater and
145 *Moringa* leaf extract produced higher concentration of K than irrigating them separately.
146 Calcium concentration decreased from the application of *Moringa* leaf extract (0.22%) to
147 wastewater and wastewater + *Moringa* leaf extract with corresponding composition of 0.14
148 and 0.12%, respectively. Magnesium concentration at wastewater irrigation was 2.3 and 5.4
149 folds that of wastewater + *Moringa* leaf extract and *Moringa* leaf extract irrigation,
150 respectively. This was significant at $P=.01$. Sodium concentration was not significantly
151 ($P=0.05$) different between the treatments as it ranged between 0.10 and 0.13%. Iron and
152 Copper concentrations of 67 and 25 mg/kg at *Moringa* leaf extract were significantly
153 ($P=0.05$) higher with 28.8 and 25% over wastewater + *Moringa* leaf extract and 48.8 and
154 47.1% over wastewater. The concentrations of the two elements decreased with the
155 application of the treatment and their concentrations were deficient according to Lockman

[8]. Zinc, Manganese and Nickel were highest at wastewater + *Moringa* leaf extract irrigation with the corresponding values of 34, 19 and 3 mg/kg, respectively. The result agreed with the finding of Al-Jaloud *et al.*, [27] who revealed increase in concentration of zinc with application of sewage. Manganese and Ni showed no significant difference due to combined effects of the treatment. Reports by [6] indicated that accumulation of heavy metals such as cadmium and lead in Maize shows a little variation in concentration of the elements. Zinc concentration increased with increases in the concentration of respective elements in the treatment. There was no significant difference in the concentration of Pb, however it was highest (0.42 mg/kg) at wastewater application. The difference in mean concentration of each element between different treatments was significant ($P=.05$) for K, Mg, Fe, Cu, Zn and Mn while P, Ca, Na, Ni and Pb were not. The coefficient of determination (R^2) show that 70-99% variability in the concentration of Ca, Na, Zn, Mn, Cu, Ni and Pb in maize plants was affected receiving wastewater, wastewater + *Moringa* leaf extract and *Moringa* leaf extract irrigation. However, P, K, Mg, and Fe correlation matrix (r) was poor and ranged from 0.02 – 0.59. The concentration of P, K, Ca, Fe and Cu were in deficient range in all treatments except for Zn and Mg at wastewater and wastewater + *Moringa* leaf extract. The variation in the concentration of minerals agrees with the report of Sadiq and Hussain [7] who found varying concentrations and responses by maize for the uptake of different elements receiving fertilizers.

3.3 Mineral concentration in soil

The effect of wastewater and *Moringa* leaf extract irrigation on mineral concentration in soil under maize plant is presented in Table 5. Phosphorus concentration was highest (5.34 mg/kg) in soils irrigated with wastewater + *Moringa* leaf extract, which are 1.2 and 3.3 folds of that accumulated by *Moringa* leaf extract and wastewater. There was no significant difference in the residual soil concentration of P irrigated with wastewater + *Moringa* leaf

181 extract and *Moringa* leaf extract. This was in agreement with the observation of Dastorani *et*
 182 *al.*, [27] who reported that, application of wastewater increased P content in the soil as well
 183 as K, where K concentration at wastewater irrigation was 0.07 cmol/kg while wastewater +
 184 *Moringa* leaf extract and *Moringa* leaf extract had the least value with significant difference
 185 at $P=0.01$. Calcium content, at wastewater + *Moringa* leaf extract was highest (0.16 cmol/kg)
 186 and was significantly different at $P=0.05$ while Magnesium content was significantly ($P=0.01$)
 187 high (0.05

188 Table 5: Effect of wastewater and *Moringa* leaf extract on mineral composition of soil
 189 under irrigated maize plants.

| Treatment | P | K | Ca | Mg | Na | OC | OM |
|-----------|------------|-------------|-------------|--------------|-------------|------------|-------------|
| | ←mg/kg→ | ←cmol/kg→ | | | | ←% | → |
| WW | 1.61±0.06b | 0.07±0.004a | 0.09±0.003b | 0.03±0.001b | 3.19±0.07a | 6.61±0.41a | 11.40±0.70a |
| MLE | 4.55±1.04a | 0.06±0.003b | 0.08±0.015b | 0.05±0.013a | 1.87±0.04b | 2.70±0.15b | 4.67±0.27b |
| WW+MLE | 5.39±0.19a | 0.06±0.002b | 0.16±0.044a | 0.04±0.003ab | 3.10±0.07a | 6.95±0.15a | 11.98±0.25a |
| LSD | 1.23 | 0.006 | 0.05 | 0.02 | 0.12 | 0.53 | 0.91 |
| <i>P</i> | * | * | * | * | * | | |
| R^2 | 0.91 | 0.84 | 0.75 | 0.53 | 0.99 | 0.99 | 0.99 |
| | Fe | Cu | Zn | Mn | Ni | Pb | |
| | ←mg/kg→ | | | | | | |
| WW | 1.38±0.10a | 0.62±0.11a | 2.96±0.01a | 13.43±0.35a | 0.03±0.001a | 0.32±0.09a | |
| MLE | 1.15±0.23a | 0.44±0.08b | 2.86±0.16a | 13.40±0.60a | 0.03±0.006a | 0.38±0.03a | |
| WW+MLE | 1.48±0.25a | 0.43±0.05b | 2.84±0.27a | 13.76±0.62a | 0.04±0.012a | 0.36±0.09a | |
| LSD | 0.41 | 0.16 | 0.36 | 1.08 | 0.02 | 0.15 | |
| <i>P</i> | NS | * | NS | NS | NS | NS | |
| R^2 | 0.41 | 0.64 | 0.12 | 0.12 | 0.17 | 0.13 | |

190 Figures in a column above followed by the same letter are not significantly ($P=0.05$) different.

191 *-significant at 5% level of probability

192 **- significant at 1% level of probability

193 NS-not significant

194 CV-Coefficient of variance

195

196 cmol/kg) at *Moringa* leaf extract. This shows that Mg present in *Moringa* leaf extract
 197 increased the Mg content of the soil. The result of Schipper *et al.*, [28] revealed that
 198 wastewater application increased contents of Mg and Ca in the soil. Sodium concentration
 199 was significantly ($P=0.01$), highest at waste water (3.19) application followed by wastewater +
 200 *Moringa* leaf extract (3.10). According to Al-Jaloud *et al.*, [26] macro- and micronutrients in
 201 the effluent neutralize the undesirable effect of high Na concentrations in waste water.
 202 Moreover, antagonistic effect between Na and K was more pronounced under low K
 203 concentrations in soil [29]. Application of the treatment did not significantly influence the
 204 mineral composition of soil Fe. A similar result was reported by Lindsay and Norvell [17].
 205 Analysis of variance shows that application of the treatment was significant ($P=0.01$) in Cu
 206 concentration with highest value of 0.62 mg/kg at wastewater application. The DTPA-
 207 extractable Zn did not show any significant difference. In a similar work, some wastewater
 208 used for irrigation increased trace metals in soils and plants, especially Zn levels in soils [30].
 209 Manganese, Ni, and Pb did not show any significant difference between the treatment and the
 210 concentration were highest (13.76, 0.04 and 0.38 mg/kg) at wastewater + *Moringa* leaf
 211 extract irrigation, respectively. This agrees with Dastorani *et al.* [28] who reported that the
 212 use of some wastewater for irrigation increased trace metals in soils and plants, most
 213 importantly Mn, Ni, and Pb levels in soils.

Regression and correlation analyses (Table 6) indicate that Na concentration in the soil affected Fe and Cu composition in maize plant which differs with the findings of Crammer and Spur [31] who reported that Ca concentration in the soil as the element affecting Ca and Mg concentrations in lettuce plants. However, Na was positively correlated with Mg ($r = 0.720$) and Zn ($r = 0.847$) while the relationship between K and Mg was the strongest ($R^2 = 0.808$) as the remaining R^2 ranged between 0.436 and 0.614. The negative correlation between K and Cu

Table 6: Regression analyses of significant correlation coefficient among soil and plant minerals

| Mineral in soil | Mineral in Plant | Regression equation | R^2 |
|-----------------|------------------|-------------------------|---------|
| P | K | $y = 1.1427x + 2.2260$ | 0.520* |
| P | Mg | $y = -0.1156x + 0.8269$ | 0.614* |
| K | Mg | $y = 38.508x - 1.966$ | 0.808** |
| K | Cu | $y = -52.059x + 5.4505$ | 0.444* |
| Ca | K | $y = 4.4715x + 2.1662$ | 0.513* |
| Na | Mg | $y = 0.2985x - 0.4295$ | 0.518* |
| Na | Fe | $y = -14.431x + 93.68$ | 0.497* |
| Na | Cu | $y = -0.5842x + 3.6549$ | 0.538* |
| Na | Zn | $y = 5.9708x + 13.922$ | 0.717** |
| Cu | K | $y = -1.5814x + 3.4515$ | 0.436* |

*= $P=0.05$, **= $P=0.01$

($r = -0.666$) is similar to the finding of Smith [32] who reported that K top dressing reduced Cu levels in Alfalfa plants.

4. CONCLUSION

The finding in the present study indicate that the concentration of minerals in maize plant were in deficient range for P, Ca, Cu, and Mn in all irrigations except K. Wastewater irrigation had Mg and Zn composition in sufficient range and Fe in deficient range. *Moringa* leaf extract irrigation had maize plant composition of Fe in sufficient range while combined wastewater and *Moringa* leaf extract irrigation had Mg, and Zn in sufficient range. Irrigating maize plant with wastewater + *Moringa* leaf extract improved P, K and Zn composition by 9.1, 43.8 and 5.2%, respectively. Therefore, wastewater from fish pond and *Moringa* leaf extract irrigation either in combination or separately did not increase the nutrient elements and heavy metals to toxic limits and therefore can be used for irrigation purposes especially maize crop.

REFERENCES

1. Al-Rashed M.F, Sherif MM. Water resources in the GCC countries: an overview. *Water Resource. Management*, 2000;14(1): 59-73.
2. Ahmad A, Inam A, Ahmad IH, Azam SZM. Response of sugarcane to treated wastewater of oil refinery. *Journal of Environmental Biology*. 2003;24 (2): 141-146.
3. Angelakis AN, Marecos-do-Monte MHP, Bontoux L, Asano T. The status of wastewater reuses practice in the Mediterranean Basin: need for guidelines. *Water Resource*, 1999;33(10): 2201-2217
4. Gori R, Ferrini F, Nicese FP, Lubello C. Effect of reclaimed wastewater on the growth and nutrient content of three landscape shrubs. *Journal of environmental horticulture*, 2000;18(2): 108-114.

- 253 5. Weber B, Avnimelech Y, Juanico M. Salt enrichment of municipal sewage: new
254 prevention approaches in Israel Environment. *Management*, 1996;20(4): 487-495.
- 255 6. Schalscha EB, Escudero P, Salgado P, Ahumada I. Chemical forms and sorption of
256 copper and zinc in soils of Central Chile. Communication *Soil Science Plant Analysis*,
257 1999;30(3/4): 497-507.
- 258 7. Sadiq M, Hussain G. Effect of chelate fertilizers on metal concentrations and growth
259 of corn in a pot experiment. *Journal of plant nutrition*. 1993;16(4): 699-711.
- 260
- 261 8. Lockman RB. Mineral composition of grain sorghum plants samples. Part 11:
262 suggested nutrient sufficiency limits at various stages of growth. Commun. *Soil Science*
263 *plant Analysis*. 1972;3:271-304
- 264 9. Emmanuel SA, Emmanuel BS, Zaku SG, Thomas SA. Biodiversity and agricultural
265 productivity enhancement in Nigeria: application of processed *Moringa oleifera* seeds
266 for improved organic farming. *Agriculture and biology journal of North America*.
267 2011;2(5): 867-871.
- 268 10. Marr IL, Cresser MS. *Environmental Chemical Analysis*. International Text Book
269 Company, USA. Chapman and Hall, New York. 184. 1983.
- 270 11. Walkley A., Black IA. An Examination of the Degtjarelf method To determine soil
271 organic Matter and proposed modification of the Chromic Acid Titration method. *Soil*
272 *Science*, 1934;37: 29-38
- 273 12. Black CA. (ed) *Methods of soil Analysis Agronomy* No. 9 part 2 Amer. SOC.
274 Agronomy, Madison Wisconsin. 1965.
- 275 13. Bates RG. *Electrometric pH determination*, John Wiley and Sons Inc. New York. 1954.
- 276 14. Bouyoucos GH. *A Recalibration of the Hydrometer for making mechanical Analysis of*
277 *soils Journal*. 1951;43: 434-438

- 278 15. Mclean EO. *Aluminium in method of soil Analysis* (ed. C.B. Black), 978-998.
279 agronomy, Madison, Wisconsin. 1965.
- 280 16. Bray RH, Kurtz LT. Determination of Total organic and available forms of phosphorus
281 in soils. *Soil Science*.1945;59: 39-45
- 282 17. Lindsay WL, Norvell WA. Development of DTPA soil test for zinc, iron, manganese
283 and copper. *Journal of Soil Science*. 1978;42: 421-428
- 284 18. Nelson JL, Brown LC,. VietsJr FG. Methods of assessing Zn status of soils using acid
285 extractable Zn: In C.A. Black *et al.*, (ed): *Methods of soil Analysis*. Part 2.*Agronomy*
286 1959;9:323-336. *Am. Soc. of Agron.And Soil Science.of Am.* Inc, Madison, Wisconsin,
287 USA.
- 288 19. Jackson MC. Soil chemical analysis. Verg: Prentice Hall, Inc., Englwood Cliffs, Nj,
289 1958, 498 s. DM 39.40
- 290 20. American Public Health Association. *Standard methods for the examination of water and*
291 *wastewater*, Published jointly by: American Public Health Association, American
292 Water Works Association, and Water Environment Federation. 2012; 160-172.
- 293 21. Mathur B. *Moringa for cattle fodder and plant growth*. 2006. www.treesforlife.org
294 retrieved on 23/04/2013.
- 295 22. SAS. Proprietary Software Release 9.1.3 (TS1M0) SAS Institute Inc., North
296 Carolina State University-Campuswide-T/R, Site 0027585003. Cary, NC, USA. 2010.
- 297 23. Mvumi C, Tagwira F, Chiteka AZ. Effect of Morianga extract on growth and yield of
298 maize and common beans. *Greener Journal of Agricultural Science*. 2013; 3(1):055-062
- 299 24. Sollenbergs, G *Farming with higher priced fertilizer*. It's back to basics. The furrow
300 1981;6:12-15.

- 301 25. Ramana S, Biswas AK, Kundu S, Saha JK, Yadava RBR Effect of distillery effluent on
302 seed germination in some vegetable crops. *Bioresources Technology*, 2002;82: 273-
303 275.
- 304 26. Al-jaloud AA, Hussain G, Al-saati AJ, Karimulia S. Effect of Wastewater irrigation on
305 mineral composition of corn and sorghum plants in a pot experiment. *Journal of Plant*
306 *Nutrition*, 1995;18: 1677-1692.
- 307 27. Dastorani MT, Hakimzadeh MA, Kalantari S. Evaluation of the effects of industrial
308 wastewater on soil properties and land desertification. *Journal Desert*. 2008; 203-210.
- 309 28. Schipper LA, Williamson JC., Kettles HA, Speir TW. Impact of Land-applied tertiary-
310 treated effluent on soil biochemical properties. *Journal Environment Quality*, 1996;25:
311 1073-1077
- 312 29. Fonseca AF, Melfi AJ, Montes CR. Maize growth and changes in soil fertility after
313 irrigation with treated sewage effluent. II. Soil acidity, exchangeable cations, and
314 sulfur, boron and heavy metals availability. *Communications in Soil Science and Plant*
315 *Analysis*, 2005;36:1983-2003.
- 316 30. Johnson AH, Siccama TG, Friedland AJ. Spatial and temporal patterns of lead
317 accumulation in the forest floor in the north eastern United States. *J. of Environmental*
318 *quality* 1982; 11:577-580
- 319 31. Crammer GR., Spur AA. Salt responses of lettuce to salinity.11. Effect of calcium on
320 growth and mineral status. *Journal of plant Nutrition*. 1986;9 (2): 131-142
- 321 32. Smith SR. *Agricultural recycling of sewage sludge and the environment*. Biddleslimited,
322 Guildford. 1996;79-83.