

# Effect of Urine Sources on Some Soil Health indicators, Maize yield and Its Heavy Metals Uptake in Abakaliki, Southeastern Nigeria

## Abstract:

The need to find alternative to inorganic fertilizer which is costly and not easily affordable to local farmers necessitates research in rare areas. Thus, this experiment was carried out at the Plant and Screen house to study effect of urine sources on some soil health indicators, maize yield and its heavy metals uptake. Completely Randomized Design was used in laying the experiment. 20kg of sieved soil was treated with different urine sources replicated five times. The result indicates that soil pH, total N and organic matter were respectively significantly ( $P<0.05$ ) higher in different urine sources than control. Human urine had significantly ( $P<0.05$ ) higher treatment effect on soil pH, percent total N and organic matter compared to other sources of urine. Similarly, human urine was 9-10%, 15-27%, 10-47% and 6-5% higher in number of leaves, plant height, grain yield and leaf area index when compared to those of cattle and goat urine sources. Significantly ( $P<0.05$ ) higher copper uptake by maize grains was obtained in control relative to those of urine sources. Copper and lead uptake by maize grains were respectively higher by 20, 80, 87% and 87, 47, 7% in control when compared to human, cattle and goat urine sources. Generally, heavy metals uptake by maize grains is below recommended safe limits for toxicity. Urine from adult animals is recommended as credible alternative for improvement of soil health status and sustainable productivity.

**Key words:** effect, maize yield, heavy metals uptake, urine sources, soil health indicators.

## INTRODUCTION

Traditional agriculture relies heavily on mineral fertilizer NPK for crop production in Nigeria and other developing countries (Nwite, 2015) and incidentally, use of fertilizer is confronted with problems of unavailability, high cost and increase in soil acidity. As a result, use of fertilizer is considered to be counterproductive and there is need for its alternative source. This alternative source is urine since it easily affordable as it could be accessed from livestock and man. It has been reported (Adeoluwa and Sulaiman, 2002), that urine contains useful nutrients which if carefully harnessed could sustain soil health status and increase its

36 productivity. Well preserved urine has good quality and could have the same effect  
37 as inorganic fertilizer in optimizing soil fertility status of soil (Nwite, 2015).  
38 Research shows that urine contains major nutrients including nitrogen, phosphorus,  
39 potassium as well as calcium and magnesium which is dependent on age and feed  
40 of the animals (Marino, 2008).

41 When there is no planned disposal of urine it naturally constitutes health hazard  
42 due to its pungent odour which could be curtailed through its proper treatment and  
43 conversion in treating soil for higher productivity (Nwite, 2015). This offensive  
44 odour is attributed to freshly accumulated urine at pH of 6.7 (Hoglung, 2001).  
45 Researchers (Heinonen Tanski and Van Wijk-Sibesma, 2005; Kichman and  
46 Peterson, 1995; Steineck *et al.*, 1999; Richert *et al.*,2002; Malkki and Heinonen-  
47 Tanski, 1999) have shown that human urine source was successfully used as  
48 fertilizer in crop production and raising flowers in Europe and other countries.  
49 Confirmatory studies have been carried out using Barley and under crop and field  
50 trials or even under home gardening (Richert *et al.*, 2002).

51 With the wide spread scarcity of inorganic fertilizer and its associated problems in  
52 food production, there is need for alternative source. If appropriate quantity of  
53 urine is applied to the soil at right time, its nitrogen contents could have the same  
54 value as that of inorganic fertilizer (Adeoluwa and Sulaiman, 2012). For instance,  
55 100 kg N per hectare of urine improved Barley production between 90 – 110 days  
56 of planting in Sweden (Richert *et al.*,2002).

57 Naturally, human being could not easily accept food crops produced with urine due  
58 to suspicion of its health hazard status and safe for consumption. This however  
59 could be overcome by treating urine for quality assurance and safe from health  
60 hazards (Nwite, 2015). In Nigeria food crops that grow around urinals or where  
61 urine is disposed are normally eaten by human beings and animals without any  
62 complaints of health problems. The objective of this experiment was to study effect

63 of urine sources on some soil health indicators, maize yield and its heavy metals  
64 uptake under Abakaliki agroecological environment.

65  
66

## MATERIALS AND METHODS

### 67 **Experimental site**

68 The research was conducted in 2014 at Plant and Screen House of Teaching  
69 and Research Farm, Faculty of Agriculture and Natural Resources  
70 Management, Ebonyi State University, Abakaliki. The area is located **between**  
71 **latitude (06° 4'N and 08° 65'E) in the South-Eastern** zone of Nigeria. The area  
72 experiences bimodal pattern of rainfall which is spread from April-July and  
73 September-November of each year. There is a break in August normally referred  
74 by residents as “August break”. At the beginning of rainfall, it is torrential  
75 and violent and is characterized by thunderstorm and **lightning**. The  
76 minimum and maximum rainfalls are 1700 and 2000 mm with a mean of 1800  
77 mm (ODNRI, 1989). The temperature during rainy season is usually low (27°C)  
78 but increases to 31 °C in dry season. Relative humidity is 80% in rainy season  
79 which declines to 60% during the cold Harmattan periods and dry season of  
80 the year (ODNRI, 1989) being characteristics of tropical climate.

81 The soil is derived from sedimentary deposits from cretaceous and tertiary  
82 periods. According to Federal Department of Agricultural Land Resources  
83 (FDALR, 1985), Abakaliki agricultural zone lies within “Asu River” and is  
84 associated with Olive brown shale, fine grained sandstones and mudstone. It is  
85 unconsolidated within 1m depth (Shale residuum) and belongs to the order  
86 ultisol classified as *typic haplustult*. The area was grown of short vegetation  
87 and medium to tall trees. There is also growth of native grasses, herbs and shrubs  
88 with patches of ground.

89

## 90 **Experimental Design and Treatment Application**

91 The experimental design used in this study was Completely  
92 Randomized Design (CRD). Human urine of male adult was collected from  
93 **prison inmates** while cattle and goat urine was sourced from **Cattle and**  
94 **Goats' market located at Gariki and Hossana, Abakaliki** respectively. **This**  
95 **was to reduce the time needed to collect enough quantity of urine for the**  
96 **study. Plastic containers of 5litres each were provided to both prison and**  
97 **Animal attendance for the purpose of collection of urine.** The animals used  
98 were of matured age. **The choice of these animals was based on ease of**  
99 **accessibility since every farming family in the locality can afford to keep them.**  
100 **These animals too are omnivorous and have common feeding habit at adult age.**  
101 The urine was stored in air-tight plastic containers for 6 months before  
102 application **to ensure sanitation process. The urine treatments were based on**  
103 **hectare equivalence of 20kg soil.**

104 **Human urine = 50,000 mg $ha^{-1}$  equivalent to 100 mg $kg^{-1}$  soil**

105 **Cattle urine = 50000 mg $ha^{-1}$  equivalent to 100 mg $kg^{-1}$  soil**

106 **Goat urine = 50000 mg $ha^{-1}$  equivalent to 100 mg $kg^{-1}$  soil**

107 **Control = 0 mg $ha^{-1}$  equivalent to 0 mg $kg^{-1}$  soil**

108 The urine rates were applied to **20kg** of soil weighed into **perforated**  
109 **polybags** two weeks after germination of maize seeds. These treatments were  
110 replicated six times to give a total of twenty four experimental **polybags** in the  
111 experiment. The **polybags** were watered to field capacity as often as moisture is  
112 required. The **polybags** were separated by 0.5m spaces while replicates were set  
113 1m apart.

## 114 **Planting of maize**

115 Maize variety (Oba super II hybrid) (*Zea mays* L.) collected from  
116 Ebonyi State Agricultural Development Programme (EBADEP), Onu Ebonyi

117 Izzi, Abakaliki was used as a test crop. The maize seeds were planted at two  
118 seeds per hole and at 5 cm depth in each pot. Two weeks after germination  
119 (WAG), thinning was carried out to allow one plant per stand. Weeds were  
120 removed by handpicking at regular intervals till harvest.

### 121 **Agronomic parameters**

122 A total of ten tagged maize plants were used for study. When the husks  
123 were dried, the cobs were harvested, dehusked, shelled and grain yield  
124 adjusted to 14% moisture content. Plant height was measured with metric  
125 ruler from the base of plant to tallest plant leaf at tasseling. Leaf area index  
126 (LAI) was determined by the formula according to Nwite *et al.* (2014).

$$127 \quad LAI = \frac{\text{Leaf area (m}^2\text{)}}{\text{Ground cover (m}^2\text{)}} \dots\dots\dots (1)$$

### 129 **Soil Sampling**

130 Auger sampler was used to collect soil samples at 0-20 cm depth from  
131 site where soil used for experiment was collected. The samples were bulked and  
132 used for routine laboratory analysis. Samples were further collected from each  
133 polybag for some post-harvest chemical properties determination.

134

### 135 **Laboratory methods**

136 The samples were dried, ground and passed through 2 mm sieve and used  
137 to determine soil properties. Particle size distribution of the experimental soil  
138 was determined using the Bouyoucous method as outlined in Gee and Or  
139 (2002) procedure. Soil pH determination was carried out in soil/water solution  
140 ratio of 1:2.5. The pH values were read off using pH meter with glass  
141 electrode (Peech, 1965). Total nitrogen was determined using Micro-kjeldahl  
142 procedure (Bremner, 1996). Available phosphorus determination was done  
143 using Bray-2 method as outlined in Page *et al.* (1982). Organic matter was

144 determined by Walkley and Black (1934) digestion method. Exchangeable  
145 bases of calcium (Ca), Magnesium (Mg), Potassium (K), and Sodium (Na) were  
146 extracted using ammonium acetate (NH<sub>4</sub>OAC) extraction method. Potassium  
147 and sodium were determined using flame photometer. The compositions of  
148 urine were determined by Atomic Absorption spectrophotometer as well as crop  
149 uptake copper (Cu) lead (Pb) using Dewis and Freitas (1976) procedure.

150

### 151 **Data analysis**

152 Data collected from the experiment were subjected to Analysis of  
153 Variance (ANOVA). Means were separated using Fishers' Least Significant  
154 Difference (FLSD) as outlined in Steel and Torrie (1980). Significance was  
155 reported at 5% probability level.

156

157

## RESULTS AND DISCUSSION

### Composition of Urine

Table 1 shows some major nutrients and heavy metals composition of urine source. There were variations in values of nutrients and heavy metals in urine source. Nevertheless, human urine have highest values of nutrients when compared to livestock sources although, comparable. Cattle and goat urine contained 0.10  $\text{mgkg}^{-1}$  each of copper (Cu) and lead (Pb) but was not found in human urine. The comparable composition of elemental concentrations in animal urine could be attributed to their adult age, omnivorous nature as well as similarity in their dietary needs.

**Table 1.** Compositions of some major nutrients and heavy metals in urine sources

| Parameter                       | Human urine | Cattle urine | Goat urine |
|---------------------------------|-------------|--------------|------------|
| pH $\text{kcl}$                 | 9.1         | 9.0          | 8.9        |
| Ammonia $\text{mgkg}^{-1}$      | 0.01        | 0.01         | 0.01       |
| Nitrogen %                      | 4.54        | 4.52         | 4.51       |
| Phosphorus $\text{mgkg}^{-1}$   | 0.04        | 0.03         | 0.02       |
| Potassium $\text{cmol/kg}^{-1}$ | 0.05        | 0.03         | 0.03       |
| Sodium $\text{cmol/kg}^{-1}$    | 0.29        | 0.28         | 0.28       |
| Copper $\text{mgkg}^{-1}$       | -           | 0.10         | -          |
| Lead $\text{mgkg}^{-1}$         | -           | -            | 0.10       |

181 **Properties of Soil before initiation of study**

182 Table 2 shows physicochemical properties of soil before initiation of study.  
183 Sand fraction was dominant in the particle size distribution. The textural  
184 class was sandy loam. The pH was 5.0 which indicate strongly acidic soil  
185 (Landon, 1991). Nitrogen was 0.13% and according to Enwezor *et al.* (1981)  
186 is very low and organic matter which had 2.2% value was moderate using  
187 Benchmark of FMAWRD (2002) for Tropical soils. Phosphorus (20.40  
188 mgkg<sup>-1</sup>) was high (Enwezor *et al.*, 1989). Exchangeable calcium was of  
189 medium value but magnesium, potassium and sodium were very low (Asadu  
190 and Nweke (1999). Cation exchange capacity recorded very low values  
191 (Asadu and Nweke, 1999). It implies that the soil was of low fertility status  
192 as obtained in Abakaliki areas for soils used for maize production as well as  
193 other crops.

194 **Table 2.** Properties of soil before initiation of study

| 195 | <b>Soil properties</b>                            | <b>Values</b> |
|-----|---|---------------|
| 196 | Sand (gkg <sup>-1</sup> )                         | 750           |
| 197 | Silt (gkg <sup>-1</sup> )                         | 140           |
| 198 | Clay (gkg <sup>-1</sup> )                         | 110           |
| 199 | Texture class                                     | Sandy Loam    |
| 200 | pH <b>kcl</b>                                     | 5.0           |
| 201 | Total Nitrogen (%)                                | 0.13          |
| 202 | Organic matter (%)                                | 2.2           |
| 203 | Available phosphorus (mgkg <sup>-1</sup> )        | 20.40         |
| 204 | Calcium (cmol kg <sup>-1</sup> )                  | 3.10          |
| 205 | Magnesium (cmol kg <sup>-1</sup> )                | 0.92          |
| 206 | Potassium (cmol kg <sup>-1</sup> )                | 0.17          |
| 207 | Sodium (cmol kg <sup>-1</sup> )                   | 0.10          |
| 208 | Cation exchange capacity (cmol kg <sup>-1</sup> ) | 7.50          |



## 209 **Effect of Urine Sources on Some Soil Health Indicators**

210 Effect of urine sources on some soil health indicators is shown in Table 3. **Soil**  
211 **health indicators are used in this study to indicate parameters that are**  
212 **determinant of soil fertility status.** Urine sources had significantly ( $P < 0.05$ )  
213 higher treatment effect on pH when compared with the control. Human source  
214 of urine had significantly ( $P < 0.05$ ) higher pH than those of cattle and goat urine  
215 sources, respectively. On the other hand, human urine was 5 and 6% higher in  
216 pH than the urine from cattle and goat. Similarly, significantly ( $P < 0.05$ ) higher  
217 treatment effect was obtained in percent total nitrogen in human and cattle  
218 sources of urine relative to control. Furthermore, human urine showed  
219 significantly ( $P < 0.05$ ) higher treatment effect on percent total nitrogen  
220 compared to those of cattle and goat sources of urine. Available phosphorus  
221 obtained in different urine sources slightly varied from that of the control. The  
222 available phosphorus of human source of urine was 14% higher than control and  
223 generally marginally higher than those of cattle and goat sources of urine. There  
224 was significantly ( $P < 0.05$ ) higher treatment effect of urine sources on percent  
225 organic matter relative to control. Urine obtained from human and goat was  
226 significantly ( $P < 0.05$ ) higher in percent organic matter than the one from cattle.  
227 This represents 21 and 14% increments in percent organic matter in human and  
228 goat sources of urine compared to that of cattle source.

229 The significant increments of pH, percent nitrogen, organic matter and  
230 improvement of available phosphorus show that these soil health indicators  
231 were released into the soil by urine sources. This finding indicates that urine  
232 could substitute mineral inorganic fertilizer as it could be used as fertilizer to  
233 supply essential and major nutrients to soil on one land and on the other  
234 improve soil health status. These findings are in line with the report of  
235 Adeoluwa and Sulaiman (2012) that urine used as fertilizer improved soil health

236 status. Several researchers (Gutser *et al.*, 2005; Schonning, 2001; Adeoluwa and  
 237 Cofie, 2012) reported positive impact of urine on nitrogen which increased and  
 238 sustained soil fertility. Higher significant positive effect of human urine source  
 239 on soil health indicators suggests that it could be more superior than other urine  
 240 sources in improving soil health indicators (Nwite, 2015). This finding had  
 241 earlier been reported by Benge, (2006) and Adeoluwa and Sulaiman (2012).  
 242 This by extension suggests that indeed, urine sources and particularly human  
 243 urine could serve as useful alternative fertilizer for crop production.  
 244 The positive impacts of the urine fertilizer on soil health indicators indicate an  
 245 improvement on the soil health status. The human urine source improved the  
 246 soil pH keeping it within a safe range of 5.6-6.0, nitrogen and organic matter at  
 247 significant levels than other sources. Peverly and Gates (1973) stated that  
 248 organic fertilizers perform better with some crops. This is further supported by  
 249 Adeoluwa and Cofie (2012) findings that urine fertilizer improved some health  
 250 indicators.

251

252 **Table 3.** Effect of urine sources on some soil health indicators

| 253 <b>Treatment</b> | <b>pH H<sub>2</sub>O</b> | <b>Total N%</b> | <b>Pmgkg<sup>-1</sup></b> | <b>OM%</b> |
|----------------------|--------------------------|-----------------|---------------------------|------------|
| 254 Control          | 5.1d                     | 0.10b           | 25.60                     | 1.05d      |
| 255 Human urine      | 6.0a                     | 0.14a           | 29.65                     | 1.76a      |
| 256 Cattle urine     | 5.7b                     | 0.12b           | 28.24                     | 1.40a      |
| 257 Goat urine       | 5.6c                     | 0.11b           | 28.30                     | 1.62b      |
| 258 FLSD(0.05)       | 0.1                      | 0.02            | NS                        | 0.05       |

259 P – Available phosphorus, OM(%) – Percent organic matter, N(%) -Percent  
 260 Total nitrogen. Treatment means with different letters indicate significant  
 261 differences.

262

263

## 264 **Effect of Urine Sources on Agronomic Yield of Maize**

265 Table 4 shows effect of urine sources on agronomic yield of maize. Urine  
266 sources had higher number of leaves, plant height, taller maize plants and larger  
267 leaf area when compared with the control, respectively. Human urine source  
268 was 9-10%, 15%-27%, 10-47% and 6-5% higher in these maize agronomic yield  
269 than those of cattle and goat urine sources. The effect of urine sources not  
270 having any significant effect on agronomic parameters of maize could be linked  
271 to inherent capability of the maize plant (Benge, 2006). Results in Table 1 did  
272 not show much variation on nutrients compositions and this could have  
273 influenced maize agronomic parameters. The generally higher agronomic yield  
274 of maize in urine sources relative to control could be attributed to improved soil  
275 health status by urine treatment. This by implication underscores the usefulness  
276 of urine as credible of alternative to inorganic fertilizer for sustenance of soil  
277 fertility status and in increased maize production that is safe for human  
278 consumption. Human urine increased agronomic yield of maize due to its high  
279 nutrients (Table 1) and its ability to release same into the soil. Benge (2006) and  
280 Adeoluwa and Sulaiman (2012) pointed out that human urine increased soil  
281 fertility and Jathropha production.

282 **Table 4.** Effect of urine sources on agronomic yield of maize

| 283 <b>Treatment</b>   | <b>No of leaf</b> | <b>Plant height (cm)</b> | <b>Grain yield (g/pot)</b> | <b>LAI</b> |
|------------------------|-------------------|--------------------------|----------------------------|------------|
| 284 Control            | 11.2              | 58.12                    | 2.0                        | 0.60       |
| 285 <b>Human urine</b> | 13.4              | 85.08                    | 4.2                        | 0.66       |
| 286 Cattle urine       | 12.2              | 72.14                    | 3.8                        | 0.62       |
| 287 Goat urine         | 12.0              | 61.98                    | 2.2                        | 0.63       |
| 288 FLSD (0.05)        | NS                | NS                       | NS                         | NS         |

289 LAI – Leaf area index.

290  
291  
292

293 **Effect of Urine Sources on Heavy Metals Uptake by Maize Grains**

294 Effect of urine sources on heavy metals uptake by maize grains is shown  
295 in Table 5. The result showed significantly ( $P < 0.05$ ) higher effect of copper  
296 uptake by maize grains in control compared to urine sources of human, cattle  
297 and goat. This accounted for 20, 80 and 37% increments of copper uptake in  
298 maize grains in control respectively when compared to human, cattle and goat  
299 sources of urine. Lead uptake by maize grains was 87, 47 and 7% higher in  
300 control relative to human, cattle and goat sources of urine.

301 The significantly higher copper uptake by maize grains grown in control plot  
302 compared to those grown in urine sources treated plots could be attributed to  
303 inputs from soil rather than urine fertilizer. Analysis of urine sources indicated  
304 very low presence of copper and lead (Table 1). The same trend of higher Cu  
305 uptake by maize grains was shown by control in Pb uptake by maize grains  
306 compared to those obtained under urine sources treatment. These findings show  
307 that urine could be used as fertilizer for crop production without placing man at  
308 a risk of ecotoxicity of heavy metals. This finding could be attributed to  
309 improved health status of soil (Table 3) and low presence of heavy metals in  
310 urine sources. The likelihood of heavy metals to build up in soil amended with  
311 urine fertilizer appears to be higher in soils treated with goat and cattle urine  
312 than human urine. Adewole *et al.* (2008) reported heavy metal uptake by crops  
313 in their work and noted that these heavy metals were stored in crop parts.  
314 Anikwe and Nwobodo (2002) and corroborated by Asadu *et al.* (2008) in their  
315 findings observed that human beings were at risk of heavy metals toxicity if  
316 they could utilize crops grown around areas contaminated with heavy metals  
317 due to eco-toxicity. This could be possible through recycling of heavy metals  
318 through food chain. Heavy metal of lead has the capacity to cause brain, renal or  
319 reproductive disorders in human beings. The heavy metals of copper and lead

320 are below 0.0-2.0 and 0.01 rated as medium to low (LASEPA, 2005) values and  
 321 far below 0.0-5.0 (WHO, 1996) or 2-1500 and 2-300 recommended as normal  
 322 by Alloway (1990). However, the interesting result is that heavy metal uptake  
 323 by maize grains could not be linked to urine treatment of soil.

324  
 325

326 **Table 5.** Effect of urine sources on heavy metals uptake by maize grains

| Treatment    | → mgkg <sup>-1</sup> ← |      |
|--------------|------------------------|------|
|              | Cu                     | Pb   |
| Control      | 0.30a                  | 0.30 |
| Human urine  | 0.04a                  | 0.04 |
| Cattle urine | 0.24b                  | 0.16 |
| Goat urine   | 0.06c                  | 0.28 |
| FLSD (0.05)  | 0.05                   | NS   |

334 Cu – Copper, Pb – Lead, Treatment means with different letters indicate  
 335 significant differences from each other.

336 **Conclusion**

337 This study has shown that urine sources could improve soil health status  
 338 and serve as useful alternative fertilizer for maize crop production. Urine  
 339 sources significantly improved soil health indicators. Agronomic parameters  
 340 responded positively to improved soil health status and performed better in  
 341 urine sources than control. Perhaps, the greatest beneficial aspect of use of urine  
 342 as fertilizer is low input of heavy metals which keep them below safe limits and  
 343 without any danger of eco-toxicity. In view of its superior performance over  
 344 other urine sources, human urine could be harvested for treatment of soil for  
 345 higher productivity rather than be allowed to be wasted through improper  
 346 disposal.

347

## References

- 349 Adeluwa, O. O. and Cofie, O. (2012). Urine as an alternative fertilizer in  
350 Agriculture: *In: Effects on* Amaranths (*Amaranthus caudatus*) Production.  
351 Renewable Agriculture and Food Systems 8:1. doi:  
352 10.1017/51742170511000512.
- 353 Adeluwa, O. O. and Suleiman, O. N. (2012). Effect of human urine on the *growth*  
354 *performances* of *Jathropha curcas* seedlings and some soil health indices.  
355 Nigerian Journal of Soil Science 22 (2):186-193.
- 356 Adewole, M.B. Adeoye, G.O. and Sridhar, M.K.C. (2008). Effect of inorganic  
357 and organo mineral fertilizers on the uptake of selected heavy metals by  
358 *Helianthus annuus L* and *Tithornia diversifolia* (Hems h) under green house  
359 condition. *Journal of Toxicological and Environmental*. Chemistry 91(5):  
360 970-980.
- 361 Alloway, B. J. (1990). Heavy metals on soils. New York: John Wiley and Sons,  
362 Inc 280p.
- 363 Anikwe, M.A.N. and Nwobodo, K. C. A. (2002). Long-term effect of municipal  
364 wastes disposal on soil properties and productivity in sites used for urban  
365 agriculture in Abakaliki, Nigeria. *Bio-resources Technology* 83:24-50.
- 366 Asadu, C. L. A. and Nweke, F. I. (1999). Soils of arable crop fields in sub-Sahara  
367 Africa: Focus on cassava growing areas, collaborative study of cassava in  
368 Africa. Working Paper No. 18. Resources and Crop -Management Division,  
369 IITA, Ibadan, Nigeria. 1782p.
- 370 Benge, M. (2006). Assessment of the potential of *Jathropha curcas*, (*biodiesel*) for  
371 the energy production and other uses in developing countries. Posted on EC  
372 Ho's website with permission of the author. July 2006 and updated August  
373 2006. 2: 22. [http://www.ascension-](http://www.ascension-publishing.com/B/Z/Jathropha)  
374 [publishing.com/B/Z/Jathropha](http://www.ascension-publishing.com/B/Z/Jathropha). Pdf  
download on January 9, 2012.
- 375 Bremner, J. M. (1996). Nitrogen-Total. *In: Sparks, D. L. (ed). Methods of Soil*  
376 *Analysis. Chemical Methods. American Society of Agronomy* 5(3):1085-  
377 1121.
- 378 Dewis, J. and Freitas, F. (1976). Physical and Chemical Methods of Soil and Water  
379 Analysis. Soil Bulletin 18, FAO, UN, Rome.

- 380 Enwezor, W. O., Udo, E. J., Usoroh, N. J. Ayoade, K. A., Adepetu, J. A., Chude,  
381 V. O. and Udegbe, C. J. (1989). Fertilizer Use and Management Practices for  
382 Crops in Nigeria, Series No 2. Fertilizer Procurement and Distribution  
383 Division, Federal Ministry of Agriculture, Water Resources and Rural  
384 Development, Lagos, Nigeria.
- 385 Enwezor, W.O., Udo, F. J. and Sobulo, R.A. (1981). Fertility status and  
386 productivity of acid sands: *In: Acid sands of Southeastern Nigeria*. Soil  
387 Science Society of Nigeria. 1: 56-73.
- 388 Federal Department of Agricultural Land Resources (FDALR) (1985).  
389 Reconnaissance Soil Survey of Anambra State, Nigeria. Soil Report, Kaduna.  
390 3p.
- 391 Federal Ministry of Agriculture, **Water** and Rural Development (**FMAWRD**)  
392 (2002). Fertilizer Use and Management Practice for Crops in Nigeria.  
393 Produced by the Federal Fertilizer Department. *In: Aduayi, E. A., Chude, V.*  
394 *O. Adebusuyi, B. A. and Olayiwola, S. O. (eds), Abuja 2002. 188p.*
- 395 Gee, G. W. and Or, D. (2002). Particle Size Analysis. *In: Dane. J. H. and Topp, G.*  
396 *C. (eds). Methods of Soil Analysis. Physical Methods. Soil Science Society*  
397 *America. 5(4):255-293.*
- 398 Gutser, R., Ebertseder, T., Weber, A., Schrami, M. and Schmmidhlter, U. (2005).  
399 Short-term and residual availability of nitrogen after long-time application of  
400 organic fertilizers on arable land. *Journal of Plant Nutrition and Soil Science*  
401 *168:439-446.*
- 402 Heinnonen-Tanski, and Van Wijk-Sibesma, C. (2005). Human excreta for plant  
403 production. *Bioresource Technology* 96:403-411.
- 404 Hoglung, C. (2001). Evaluation of **Microbial Health Risks Associated** with the  
405 **Reuse** of **Source-Separated Human** urine. **Ph.D Thesis**, Department of  
406 Biotechnology, Applied Microbiology, **Royal Institute of Technology** (KTH)  
407 **Stolkolm Sweden. 62-64.**
- 408 Kirchmann, H. and Peterson, S. (1995). Human urine-chemical composition  
409 and fertilizer use efficiency. *Fertilizer Resources* 40:149-154.
- 410 Lagos State Environmental Protection Agency (LASEPA, 2005). Lagos State  
411 Environmental Reports 3: 28-33.

- 412 Landon, J. R. (eds) (1991). Booker, Tropical Soil Manual. A hand book for  
413 Soil Survey and Agricultural Land Evaluation in Tropics and Subtropics  
414 New York, USA, John Wiley and Sons: Inc. Third Avenue.
- 415 Malkki, S. and Heinonen-Tanski, (1999). Composition of toilets in permanent  
416 houses. *In: 1 Use of municipal organic wastes. Proceedings of Nigerian*  
417 *Journal of Forestry Seminar 292 DIAS-report. 13: 147-154.*
- 418 Marino, C. (2008). Urine composition depends on certain factors. *Journal of*  
419 *Soil Science Society of America. 73:159-219.*
- 420 Nwite, J.N. (2015). Effect of urine source on soil **chemical** properties and maize  
421 yield in Abakaliki, South-Eastern Nigeria. *International Journal of Advance*  
422 *Agricultural Research 3:31-36.*
- 423 Nwite, J. N. (2013). Evaluation of the productivity of a spent Automobile oil  
424 contaminated soil amended with organic wastes in Abakaliki, Southeastern  
425 Nigeria. Ph.D Thesis, University of Nigeria, Nsukka 130p.
- 426 Nwite, J. N., Okolo, C. C., Ezeaku, P. I. and Enyioko, C. (2014). Effect of  
427 Integrated Nutrient Management on Soil Chemical properties and maize  
428 yield on a sandy loam in Abakaliki, Nigeria. *International Journal of*  
429 *Agriculture and Biosciences 3(6):278-282.*
- 430 Overseas Development of Natural Resources Institute (ODNR) (1989). Nigeria  
431 Profile of Agricultural Potential ODA, United Kingdom. 3p.
- 432 Page, A. L., Miller, R. H. and Keeney, D. R. (1982). Methods of Soil  
433 Analysis. *American Society of Agronomy 9:539-579.*
- 434 Peech, M. (1965). Hydrogen activity. *Methods of Soil Analysis. In: Black CA*  
435 *(ed). American Society of Agronomy 9(1):914-926.*
- 436 Perverly, J.H. and Gates, P.B. (1993). Utilization of municipal solid waste and  
437 sludge compost in crop production systems. *In: Sewage sludge, Land*  
438 *utilization and the Environment. Proceedings of a conference by American*  
439 *Society of Agronomy.*
- 440 Richert Stintizing, A., Rodhe, L., Akerhieln, H. and Stenieck, S. (2002).  
441 Human urine as a fertilizer and plant nutrients application technique and  
442 environmental effects. *In: Venglosk, J., Greserova, G. (eds). Proceedings*



- 443 of 10 International Conference Ramiran 2002 Network. FAO European  
444 System of Cooperative Research Network 161-162 Pp.
- 445 Schonning, C. (2001). Urine diversion-hygienic risks and microbial guidelines  
446 for reuse. Department of Parasitology, Mycology and Environmental  
447 Microbiology. Swedish Institute for Infections.
- 448 Steel, G. D. and Torrie J. H. (1980). Procedures of statistics. A biometrical  
449 approach, 2nd ed. New York, McGraw Hill, Book Company 63p.
- 450 Steineck, S., Richert Stintzing, A., Rodhe, L., Elinquist, H. and Jakobsson,  
451 M. C. (1999). Plant nutrients in human and food refuse. *In*: Proceedings  
452 of Nigerian Journal of Forestry Seminar 292. DIAS report. 13:125-130.
- 453 Walkey, A. and Black, A. (1934). An examination of the Degtjareff method for  
454 determining soil organic matter and a proposed modification of the chronic  
455 acid titration method. *Soil Science* 37:29-38.
- 456 World Health Organization (WHO, 1996). Guideline for drinking quality water.  
457 Switzerland 1(2&3):50-57.