

Genetic Diversity and Variability among Papaya (*Carica papaya* L.) Genotypes using Multivariate Analysis

Emede, Timon Ogheneweware^{1*}, Ajayo, Babatope Samuel¹ and Agho Collins²

¹Department of Crop Science, University of Benin, P.M.B. 1154 Benin City, Nigeria 300001

²Plant Breeding Division, Nigerian Institute for Oil Palm Research (NIFOR).P.M.B.1030, Benin City, Edo State, Nigeria.

AUTHORS' CONTRIBUTIONS

The first author collected the papaya seeds, designed and supervised the study. The second author managed the experiment, collected data and wrote the first draft, while the third author performed the statistical analysis, and wrote the final manuscript. All authors read and approved the final manuscript.

ABSTRACT

Multivariate analysis was used to group and study the pattern of genetic variability and relationship among ten papaya genotypes in Benin City, Nigeria. The experiment was conducted at the Teaching and Research Farm of the Faculty of Agriculture, University of Benin, Benin City, Nigeria (Latitude: 6° 33'N, Longitude: 5° 37'E; 79m asl). Field evaluation of the papaya genotypes was carried out from October 2012 to June 2013. The experiment was laid out as a randomized complete block design with three replications. The result of the Principal Component Analysis (PCA) indicated that the contribution of the first three factors with Eigen value greater than one accounted for 93.0% of the total variation. The first component loaded highly for fruit characters except fruit flesh thickness and fruit diameter which loaded highly in the third component. The vegetative components loaded highly in the second component. Thus the papaya genotypes can be distinguished based on either yield or vegetative characters with more reliability on the yield parameters which mostly accounted for the greater proportion of variation. Better resolution can be achieved when both are considered together. The accessions were allocated into three clusters by Cluster Analysis. The three clusters formed indicate intraspecific phenotypic dissimilarity among the ten genotypes especially with the separation of the genotypes that were collected from similar environments. PCA and Cluster Analysis produced similar results in classifying similar genotypes into their separate clusters. Between accessions, CP006 and CP012 had the highest genetic distance while the lowest genetic distance was between CP001 and CP011. The level of variability observed suggested a high diversity among the genotypes. The phenotypic variations could be explored for utilization, conservation and for future genetic improvement by selection of genotypes with promising agronomic characters. CP012 was particularly superior with respect to the studied traits and was the only genotype in its cluster group. Molecular studies would be useful to confirm the genetic diversity and characterize these genotypes for more detailed examination.

Keywords: papaya, phenotypic variation, cluster analysis, principal component analysis.

1. INTRODUCTION

The papaya (*Carica papaya* L.) is native to tropical America and belongs to the small family Caricaceae, which includes 35 species placed in six genera [1]. Among all species, 32 are dioecious, two trioecious and one monoecious [2]. The papaya is the only species of the genus *Carica*, also being the

best known and most economically important within the family [3], showing widespread cultivation in tropical and subtropical regions around the world. It is commonly known for its food and nutritional values throughout the world [4]. Papaya is a major fruit crop worldwide that is primarily consumed as fresh fruit. It is highly abundant and is commonly known as pawpaw in Nigeria. It is an invaluable plant that is prevalent throughout Tropical Africa and Nigeria is the third largest producer globally [5].

Genetic diversity among parental is considered an important factor for obtaining heterotic hybrids. This diversity is one of the restraining tools for breeding programmes based on hybridization, because it generates parameters for identifying superior parental. According to [6], this distance is essential to increase the chance of recovering superior genotypes. Several authors have reported the efficacy of genetic divergence as a criterion for choosing parents to be crossed [7, 8, 9]. It plays an important role in plant breeding because hybrids between lines of diverse origin generally display greater heterosis than those between closely related strains [10] which permits to select the genetically divergent plants to obtain the desirable recombination of the segregating generation.

Multivariate analysis is a useful tool in quantifying the degree of divergence between biological population at genotypic level and to assess relative contribution of different components to the total divergence both at intra- and inter-cluster levels [11, 12]. The multivariate analysis such as the Principal Component Analysis (PCA), cluster and discriminant analysis have been used to uncover similarities between variables and determine the amount of variation and the most appropriate breeding strategy for papaya improvement programme [13, 14]. Papaya exhibits considerable phenotypic variation for many morphological and horticultural traits, including fruit size, fruit shape, flesh color, flavor and sweetness, length of juvenile period, plant stature, stamen anther, and carpel abortion [15, 16]. Some multivariate studies on papaya have reported genetic diversity using morphological data, isozyme, microsatellite markers, SSR (simple sequence repeats) [13, 17, 14]. In another study in papaya, morpho-agronomic and molecular data were used together to estimate the genetic distance between genotypes derived from backcrossing [18]. The objective of this research was to study the genetic variation among ten papaya genotypes by using multivariate analysis to classify the accessions in order to identify divergent parents for breeding programme and select the most suitable combinations which would provide superior segregates, as well as, to investigate the importance of the evaluated characters.

2. MATERIAL AND METHODS

The ten papaya genotypes used in this study (CP001, CP002, CP003, CP004, CP005, CP006, CP007, CP008, CP011 and CP0120) were obtained from Uselu market and home gardens in Benin City metropolis. The collection was undertaken in October, 2011. The seeds of each papaya accessions were sown in October, 2011 in drill rows of 4m by broadcasting and gradually thinned (by discarding all male plants) to three plants per row, spaced at 2m x 2m. Thus, each papaya accession used in this study was either female or bisexual plants and evaluation for sex type for each genotype was not performed during the study. The experiment was conducted at the Teaching and Research Farm of the Faculty of Agriculture, University of Benin, Benin City, Edo State, Nigeria (Latitude: 6° 33'N, Longitude: 5° 37'E; 79m asl). Field evaluation of the papaya genotypes was carried out from October 2012 to June 2013. The experiment was laid out as a randomized complete block design with three replications.

At flowering, about 9 months after planting, NPK 15:15:15 was applied at the rate of 0.6kg per plant. Each papaya stand received additional 0.6kg of NPK 15:15:15 at 18 months after planting. Data were collected on several quantitative traits derived from a subset of the papaya descriptor list published by IPGRI [19] (Table 1).

The stem height was measured from the ground level to the stem tip with a tape, while stem girth was measured at 10th internode when counted downward from the point of attachment of the first fruit. Leaf length was obtained by measuring from the point of attachment of the petiole to the tip of the longest leaflet, while leaf width was the measured widest portion of the leaves. Petiole length was measured as the distance from the point of its attachment to the stem to the point of attachment of the palmate leaves. At harvest, the fruit was detached from the peduncle and the fruit weight was determined by weighing with a weighing scale. Longitudinal sections of the harvested fruits per tree were made, and then the fruit length was determined from pole to pole of the fruits. Fruit diameter was determined from the equator of the sectioned fruit and the flesh thickness was measured with a measuring tape.

Table 1. Descriptor list used for morphological characterization of papaya genotypes

| Organ | Quantitative Traits |
|---------|---------------------|
| Stem | Height |
| | Girth |
| Petiole | Length |
| Leaf | Length |
| | Width |
| Fruit | Length |
| | Diameter |
| | Flesh thickness |
| | Number |
| | Weight |
| | Fruit yield |

2.1 Data analysis

The data were subjected to analysis of variance and basic statistics in order to give initial information about the traits across the genotypes and parameters such as mean, standard deviation (SD) and coefficient of variation (CV) for each one of the 13 studied traits were calculated. Cluster analysis was implemented using the hierarchical clustering techniques, based on UPGMA (unweighted pair-group method using an arithmetic average) by subjecting the 10 x 13 data matrix to cluster analysis based on squared Euclidean distances for similarity measurement. In order to identify the patterns of phenotypic variation, Principal Component Analysis (PCA) was conducted. The PCs with Eigen value >1.0 was considered as inherently more informative than any single variable alone [20]. The PCA was further rotated using the varimax method with Kaiser Normalization. SAS [21] and SPSS version 17 for Windows statistical software packages were used for the analysis.

3. RESULTS

Pattern of variation among the genotypes were different for each traits. The largest variation was observed for fruit yield (t/ha), total fruit yield (kg), and fruit yield per plant (kg) with coefficients of variations of 75.47%, 75.45% and 69.02% respectively. Generally, all the traits show moderate to high variability. Coefficient of variation (CV %) ranged from 13.52 to 75.47% for the various traits. The coefficient of variation was highest for fruit yield (t/ha), while the lowest level was showed by leaf length (Table 2). Based on the agro-morphological characters, the papaya genotypes collected showed variation in most of the characters especially the fruit yield traits, but was not statistically significant (Table 3).

Table 2. Basic statistics of the agro-morphological characters across 10 genotypes of papaya

| Characters | Mean | <u>Mean</u> <u>Min.</u> – <u>Max.</u> | Standard deviation | Coefficient Variation (CV) |
|----------------------------|--------|--|-----------------------|----------------------------------|
| Stem height (cm) | 170.42 | 138.2 – 209.0 | 23.33 | 13.69 |
| Stem girth (cm) | 15.66 | 12.2 - 18.8 | 2.42 | 15.47 |
| Leaf length (cm) | 41.10 | 34.6 - 52.8 | 5.56 | 13.52 |
| Leaf width (cm) | 57.75 | 51.0 - 63.6 | 7.83 | 13.56 |
| Petiole length (cm) | 61.79 | 41.26 – 84.00 | 13.34 | 21.60 |
| Fruit length (cm) | 24.14 | 18.2 - 31.9 | 4.95 | 20.52 |
| Fruit diameter(cm) | 13.44 | 10.8 - 16.7 | 2.14 | 15.95 |
| Fruit flesh thickness(cm) | 3.03 | 2.3 - 3.5 | 0.42 | 14.00 |
| No. of fruits per plant | 14.60 | 8.0 - 26.0 | 4.88 | 33.43 |
| Number of fruits Harvested | 9.90 | 3.0 - 21.0 | 5.09 | 51.38 |
| Total fruit yield (kg) | 22.00 | 5.9 - 60.5 | 16.60 | 75.45 |
| Fruit yield per plant (kg) | 11.52 | 3.0 - 30.3 | 7.95 | 69.02 |
| FruitYield (t/ha) | 30.52 | 8.2 - 84.0 | 23.03 | 75.47 |

Table 3. Mean values of agronomic characters in the ten papaya lines

| Lines | Stem height (cm) | Stem girth (cm) | Leaf length (cm) | Leaf width (cm) | Petiole length (cm) | Fruit length (cm) | Fruit diameter (cm) | Fruit flesh thickness (cm) | No. of fruits per plant | Number of fruits Harvested | Total fruit yield (kg) | Fruit yield per plant(kg) | Fruit yield (t/ha) |
|-------|------------------|-----------------|------------------|-----------------|---------------------|-------------------|---------------------|----------------------------|-------------------------|----------------------------|------------------------|---------------------------|--------------------|
| CP001 | 144.5 | 14.1 | 34.6 | 49.1 | 53.8 | 19.2 | 11.1 | 2.6 | 10.0 | 7.0 | 12.9 | 7.1 | 17.9 |
| CP002 | 138.2 | 14.9 | 37.6 | 54.6 | 53.3 | 25.0 | 13.5 | 3.3 | 15.0 | 12.0 | 23.8 | 11.9 | 33.0 |
| CP003 | 181.0 | 18.2 | 43.6 | 63.2 | 70.5 | 31.9 | 16.2 | 3.4 | 15.0 | 11.0 | 36.8 | 18.4 | 51.0 |
| CP004 | 182.5 | 16.4 | 42.9 | 59.4 | 52.3 | 29.4 | 13.6 | 3.3 | 15.0 | 10.0 | 24.9 | 13.1 | 34.5 |
| CP005 | 175.0 | 14.3 | 37.4 | 51.1 | 62.9 | 21.2 | 10.8 | 2.3 | 17.0 | 14.0 | 21.3 | 10.6 | 29.5 |
| CP006 | 138.3 | 12.2 | 36.1 | 51.0 | 41.3 | 21.1 | 11.2 | 2.8 | 8.0 | 3.0 | 5.9 | 3.0 | 8.2 |
| CP007 | 189.5 | 16.7 | 46.3 | 63.6 | 84.0 | 22.5 | 14.5 | 2.9 | 14.0 | 8.0 | 18.5 | 9.3 | 25.7 |
| CP008 | 209.0 | 18.5 | 41.8 | 59.9 | 67.9 | 18.2 | 14.9 | 3.5 | 15.0 | 5.0 | 9.0 | 6.7 | 12.5 |
| CP011 | 170.7 | 12.5 | 37.8 | 51.5 | 53.4 | 21.9 | 11.9 | 2.7 | 11.0 | 8.0 | 6.4 | 4.8 | 8.9 |
| CP012 | 175.5 | 18.8 | 52.8 | 74.1 | 78.5 | 31.0 | 16.7 | 3.5 | 26.0 | 21.0 | 60.5 | 30.3 | 84.0 |

3.1 Cluster analysis

The dendrogram of the hierarchical cluster analysis (HCA) separated the 10 accessions into three different clusters with Squared Euclidean distance dissimilarities ranging between 3.09 to 108.31 (Table not shown). Phenogram based on squared Euclidean distance coefficients using 13 traits placed the 10 genotypes into three main clusters (Figure 1). First cluster consisted of a total of four genotypes (40%) namely CP001, 011, 006 and 005. The second cluster consisted of five clusters while the remaining genotype (CP012) was in cluster 3. The Euclidean genetic distance between CP006 and CP012 was the highest (108.3) while the lowest Euclidean distance was between CP001 and CP011 (3.09). Except for stem height, petiole length and fruit length, the third cluster (CP012) had highest values for the remaining ten characters (77% of the studied traits).

3.2 Principal Components Analysis (PCA)

The results showed that three principal components with eigenvalues more than one explained 93% of total variability (Table 4). The first (PC1) as mostly fruit characters that explained 70.4% of total variability. Among the property vectors of PC1, leaf length, leaf width, fruit length, fruit per plant, number of fruit harvested, total fruit yield, fruit yield per plant and fruit yield per hectare have higher values. The second principal component (PC2) is plant vegetative characters which explain 14.74% of total variability. Among the property vectors of PC2, stem height, stem girth, leaf length, leaf width and petiole length have higher values. The third principal component (PC3) is the remaining fruit characters that explain about 7.83% of total variability. All the agro-morphological characters in each component were positively correlated. The first two PCs were plotted to observe the relationship among the ten genotypes of papaya (Fig. 2). Genotypes namely CP001, 011, 006 and 005 were found to be cluster together while (CP012) was in separate cluster (a singleton) as also observed in the cluster analysis. Genotypes to the extreme right of the component plot are those genotypes with high PC1 scores and included CP007 and CP008. These genotypes have high fruit yield qualities as PC 1 was particular high in fruit yield loadings. Similarly, CP012 located at the top extreme left of the component plot with high PC2 scores had high vegetative characteristics.

Table 4. Eigen values, variance, cumulative variance and component scores of the first three principal components for 13 quantitative traits in 10 papaya lines.

| Traits | PC1 | PC2 | PC3 |
|----------------------------|-------|-------|-------|
| Stem height | -.081 | .902 | .182 |
| Stem girth (cm) | .347 | .681 | .576 |
| Leaf length (cm) | .575 | .611 | .447 |
| Leaf width (cm) | .565 | .569 | .553 |
| Petiole length (cm) | .392 | .854 | .091 |
| Fruit length (cm) | .686 | -.054 | .596 |
| Fruit diameter(cm) | .389 | .525 | .737 |
| Fruit flesh thickness(cm) | .169 | .235 | .924 |
| No. of fruits per plant | .826 | .443 | .182 |
| Number of fruits Harvested | .967 | .145 | .004 |
| Total fruit yield (kg) | .913 | .182 | .352 |
| Fruit yield per plant (kg) | .895 | .221 | .375 |
| FruitYield (t/ha) | .913 | .182 | .352 |
| Eigen values | 9.16 | 1.92 | 1.06 |
| Cumulativeeigen values | 9.16 | 11.08 | 12.14 |
| Variance (%) | 70.44 | 14.74 | 7.83 |
| Cumulative variance (%) | 70.44 | 85.18 | 93.01 |

PC1 (First Principal Component)

PC2 (Second Principal Component)

PC3 (Third Principal Component)

HIERARCHICALCLUSTER ANALYSIS

Dendrogram using Average Linkage (Between Groups)




Figure 1: Dendrogram based on UPGMA analysis generated using the phenetic relationships among the 10 papaya lines.




Fig.2. Scattered diagram of first two principal components based on mean values of 13 quantitative traits in 10 papaya lines.

4. DISCUSSION

All the 10 studied genotypes differed from one to another for all the characters, although were not statistically significant. However, it was not possible to relate morphological diversity and performance to sex type. It was reported by [13] that sexual type affected leaf and fruit traits and cluster analysis shows little morphological differentiation between origins of papaya accessions. The CV% for the trait number of fruits harvested, total fruit yield, fruit yield per plant and fruit yield (Kg and tons/ha) is high which implies that there is a high level of variability among the evaluated genotypes with respect to these traits. It means that it is possible to improve these fruit characters in the evaluated papaya population through selection. Wide range of variability in number of fruits per plant and fruit yield per plant was also observed in true breeding accessions of papaya [22] and in hermaphrodite papaya genotypes [23]. Therefore, these traits can be used as principal selection criteria for papaya improvement programme. The range of values for most traits was high with the various genotypes having superior performance for each variable with CP012 particularly superior with respect to the studied traits and was the only genotype in its cluster group indicating that genotype of this cluster could be used as parent in future hybridization program for improved fruit yield qualities. Crosses between individuals from different clusters may result in high heterosis. Even though, the genetic mechanisms that explain heterosis are not fully understood, it is well documented that crosses between unrelated and consequently genetically distant parents, show greater hybrid vigor than crosses between closely related parents [24] since it is expected to produce new recombinants with desired traits. One of the important approaches to papaya breeding is hybridization and subsequent selection. Parents' choice is the first step in plant breeding program through hybridization. In order to benefit transgressive segregation, genetic distance between parents is necessary [25]. The higher genetic distance between parents, the higher heterosis in progeny can be observed [26, 27]. The genetic distance between CP006 and CP012 was the highest and thus crosses between these two parents should be considered. The principal component analysis indicated significant contributions in the component loadings of the 13 traits, which underpins their relevance in determining the variability among the 10 genotypes. The sign on the loadings indicates the direction of relationship between the components and the trait measured [28]. Two traits with high weighting in the same

component are expected to be highly correlated. This principle suggests that these traits could be probably influenced by similar gene(s) and may be used to identify variation among genotypes [28].

In spite of the reduction of the characters to only three principal components, it was possible to account for over 70% of the total variations among the papaya accessions. Thus the capacity of PCA in data reduction without loss of information was confirmed [29]. Component one loaded highly for fruit traits and accounted for over 70% of the total variation among the genotypes and therefore measured the importance of fruit characters in distinguishing the papaya genotypes. Fruit qualities were also sufficient in distinguishing papaya accession in Babados as reported by [13]. In the present study, principal component analysis captured most of the variation within the genotypes in higher number of axes compared to discriminant analysis. Thus, a combination of PCA and discriminant analysis would be appropriate for describing the variation among papaya genotypes. [9] also obtained similar result in coconut. Given the food and nutritional values of papaya, the morphological characterization of the papaya genotypes would serve as a good guide for the genetic development, conservation, collection and utilization of germplasm.

5. CONCLUSION

This research work was initiated with the aim of studying the genetic diversity and identifying key morphological traits to distinguish papaya accessions in Benin City, Edo State, Nigeria. The goal was to identify genotypes with desired traits for further genetic improvements. Multivariate analysis was used to group and study the pattern of genetic variation and relationship among the accessions. The level of variability observed suggested a high diversity among the genotypes. Further studies would require collection of papaya germplasm from other ecological zones in Nigeria in order to widen the scope of the study, and evaluation of sex type for each genotype would be needed so as to relate morphological diversity and performance to sex type. Cluster and principal component analysis classified the genotypes into different similar groups with observable morphological characteristics. Fruit yield characters were superior in distinguishing the genotypes. Representative genotypes may be chosen from any particular groups. The characters identified in this study should guide the breeders in identification, selection and combination of genotypes to obtain lines with the desired traits. Grouping of genotypes by multivariate methods in the study is of practical value for the papaya breeders. However, Molecular studies would be useful to confirm the genetic diversity and characterize these genotypes for more detailed examination and further study across years and locations should be done. This may help to emphasize the availability of these genetic resources for future breeding programmes.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Scheldeman X, Kyndt T, Coppens D'Eeckenbrugge G, Ming R, Drew R, Droogenbroeck BV, Van DP, Moore PH. *Vasconcellea*. In : Wild crop relatives : genomic and breeding resources. Tropical and subtropical fruits. Kole C (ed.). Heidelberg : Springer [Allemagne] 2011; 213-249.
2. Ming R, Yu Q, Moore PH. Sex determination in papaya. *Semin. Cell Dev. Biol.* 2007; 18, 401-408.
3. Van Droogenbroeck B, Breyne P, Goetghebeur P, Romeijn-Peeters E. AFLP analysis of genetic relationships among papaya and its wild relatives (Caricaceae) from Ecuador. *Theor. Appl. Genet.* 2002; 105, 289-297.
4. Krishna KI, Paridhav M, Jagruti AP. Review on nutritional, medicinal and pharmacological properties of papaya (*Carica papaya* linn.). *National Prod. Reliance* 2008; 7 (4), 364 – 373.
5. FAO (Food and Agriculture Organization) for the United Nations. :Statistics online website. <http://faostat.fao.org>. 2015.
6. Cruz CD, Carneiro PCS, Modelos biométricos aplicados a melhoramento genético. - Viçosa: Imprensa Universitária. 2003; Vol. 2, 585pp.

7. Dias LASand Kageyama PY. Multivariate genetic divergence and hybrid performance of cacao (*Theobroma cacao* L.). Braz. J. Genet.1997;20, 63-70.
8. Hamza AM, Agho C, Ado SG, Ikuenobe CE, Ataga CD Odewale JO. Proximate Compositions Evaluation and Variability among Cultivars of Date Palm (*Phoenix dactylifera*L.) in Nigeria.International Journal of Plant & Soil Science 3(3): 248-259, 2014
9. Odewale JO, Agho C, Ataga CD, Aisueni NO, Ikuenobe CE, Okoye MN, Odiowaya G, Edokpayi AA, Ahanor MJ, Uwadiae EO. Pattern of genetic diversity and variability in germplasm resources of local and exotic coconut (*Cocosnucifera* L.) cultivars in Nigeria. Scholarly J. Agric. Sci. 2012; 2(8), 202-207.
10. Singh P. Studies on genetic variability and diversity of rice. Madras Agric. J. 1983; 70(7), 436-440.
11. Jatasra DS, Parada RS. Genetic divergence in wheat under different environmental conditions. Cereal Res. Comm. 1978; 6, 307-317.
12. Zahan MI, Bhuiyan MSR, Hossain MS. Genetic divergence in Oleiferous Brassica species. J. Sher-e-Bangla Agric. Univ. 2008; 2(1), 1-6.
13. Ocampo J, d'Eeckenbrugge GC, Bruyere S, De Belliare LL, Ollitrault P. Organization of morphological and genetic diversity of Caribbean and Venezuelan papaya germplasm. Fruits. 2006; 61, 25 – 37.
14. de Oliveira EJ, Amorim BO, Matos ELS, Costa JL, Castellan MS, Padua JG, PantaJLL. Polymorphism of microsatellite markers in papaya (*Carica papaya* L.). Plant Mol. Biol. Rep. 2010; 28, 514 – 530.
15. Kim M, Moore P, Zee F, Fitch MMM, Steiger D, Manshardt R, Paull R, Drew R, Sekioka T, Ming R. Genetic diversity of *Carica papaya* as revealed by AFLP markers. Genome. 2002; 45, 503–512.
16. Ocampo Pérez J, Dambier D, Ollitrault P, d'Eeckenbrugge GC, Brottier P, Froelicher Y, Risterucci AM. Microsatellitemarkers in *Carica papaya* L.: isolation characterization and transferability to *Vasconcellea* species. MolEcol Notes. 2006; 6, 212–217.
17. Ocampo J, d'Eeckenbrugge GC, Ollitrault P, Risterucci AM, Dambier D. Papaya genetic diversity assessed with microsatellite markers in germplasm from the Caribbean region. ActaHorticulturae. 2007; 740, 93 – 102.
18. Ramos HCC, Pereira MG, Pinto FO, Ribeiro EH. Multivariate analysis to determine the geneticdistance among backcross papaya (*Carica papaya* L.) progenies. Genet. and Molecular Res.2012; 11(2), 1280 – 1295.
19. Anon., Descriptors-for-papaya, IBPGR, FAO, Rome, Italy, 1988.
20. Kaiser HF. The application of electronic computers to factor analysis. Educational and Psychological Measurement. 1960; 20, 141–151. <http://dx.doi.org/10.1177/001316446002000116>
21. SAS Institute. Inc. SAS users's guide. Statistics, version 9.00. SAS Institute, Inc., Cary, NC, 2002.
22. Jana BR, Rai M, Das B, Nath V. Genetic variability and association of component characters for fruit yield in papaya (*Carica papaya* L.). The Orissa J. Hort. 2005; 33(1), 90 – 95.
23. Asefa W, Etissa E, Dagne A, Ayele L, Kebede G, Damtew M, Ayalew M. Genetic variability and association of characters of hermaphrodite papaya genotypes. Adv. in Biosci and Bioeng. 2016; 4(2), 10 -15.
24. Stuber CW. Heterosis in plant Breeding. Plant Breed. Rev. 1994; 12:227-251.
25. Joshi BK, Mudwari A, Bhatta MR, Ferrara GO. Genetic diversity in Nepalese wheat cultivars based on agro-morphological traits and coefficients of parentage. Nep. Agric. Res. J. 2004; 5, 7-17.
26. Joshi AB, Dhawan NL. Genetic improvement of yield with special reference to self-fertilizing crops. Indian J. Genet. And Plant Breed. 1966; 26,101-113.
27. Anand IJ, Murrty BR. Genetic divergence and hybrid performance in linseed. Int. J Genet plant Breed. 1968; 28, 178-185.
28. Biabani AR, Pakniyat H. Evaluation of seed yield-related characters in sesame (*Sesamumindicum* L.) using factor and path analysis. Pakistan J. Bio. Sci. 2008; 11, 1157-1160.
29. Ross CJS. Statistical algorithms. Algosithorus AS13-AS15. Appl. Stat. 1969; 18,103-110.