

Potential

ABSTRACT

An experiment was conducted at the Research Farm of the Department of Genetics and Plant Breeding of Hajee Mohammad Danesh Science and Technology University, Dinajpur, Bangladesh during 2015-2016 to study the performance of the selected wheat variety based on somemorphological traits. Twenty four wheat varieties were used in the experiment where they were collected from Wheat Research Centre, Bangladesh Agriculture Research Institute, Dinajpur. The experiment was conducted in randomized completely blocked design with three replications. Different yield contributing traits like thousand grain weight (g), number of grains per spike, number of spikelet's per spike, days to anthesis, heading days, plant height (cm), days to maturity and grain yield (g/plot) were assayed. The result of the analyses of variance for all the traits showed significant differences among the genotypes. The experimental result demonstrated that the variety PYT-15, BARI Gom 25 and PYT-12 performed better among the tested genotypes in relation to yield and yield contributing traits and those could be recommended for further popularization in different parts of Bangladesh.

Key words: Genotypes, Growth, Performance, Yield and Wheat

1. INTRODUCTION

Being the king of cereals, wheat is the staple food all over the world [1] that contributes more calories and proteins than any other cereal crops to the world diet [2, 3, 4]. All over the world wheat is a very nutritious food grain among the all grains and grows across the globe due to its wider genotypic adaptability. It is consumed as staple food by more than 35% of world population [5]. Wheat plays an important role in the nutrition of rapidly growing population both in our country and the world as used for both human and animal nutrition [6]. All over the world wheat product(s) are consumed in one of these forms viz. chapati, bread, biscuit, pasta and fermented products [7]. Besides this, wheat is considered a good sink of protein, minerals, B-group vitamins and dietary fiber [8, 9]. The wheat germ or embryo is relatively rich in protein, fat and several of the B-vitamins [10]. Nowadays the production of wheat is increasing in many countries due to its higher demand as a consequence of faster population growth [11]. After rice, wheat is the second most important cereal crop in Bangladesh [12] and per year its consumption rate is increasing about 3% [13]. But in Bangladesh the annual wheat production is about only 1.4 million tons [14] which is much lower than the national annual demand. Despite to higher yield potentiality the average yield of Bangladeshi wheat varieties are much lower than the other wheat grower countries of the world. The genetic potential of the crops is the vital factor for harvesting suitable environment in grain yield [15]. Under the changed global conditions, we have already experienced that from other countries food may not be affordable through import. It is desired to have a higher yield per unit area

variation. Careful selection may help to obtain lines higher in yields with better quality. Genetic variability can offer opportunity for the effective selection for high yielding wheat variety rich in grain quality. It may require maximizing wheat production rather than economic yield, depending on global food policy and production. In order to explore the varieties potentiality in maximizing wheat yield and to assist breeding program in selecting lines with higher yield potentials, the yield potentiality of newly developed wheat varieties and promising lines are needed to investigate. Keeping these points in mind the present investigation was undertaken to evaluate the performance of some selected genetically diverged wheat genotypes.

2. MATERIALS AND METHODS

By using 24 wheat base materials this field research was conducted at the Research Farm of the department of Genetics and Plant Breeding of Hajje Mohammad Danesh Science and Technology University (Table 1). The seeds were collected from WRC (Wheat Research Centre) of Bangladesh Agricultural Research Institute. The experiment was conducted in a Randomized Complete Block Design (RCBD) with three replications. The experimental soil was sandy loam with the unit plot size 2.0 m x 5.0 m. The plot to plot distance was 0.75 m and block to block was 1.5 m. The manures and fertilizers like Cow dung, Urea, TSP, MOP, Gypsum and Boric acid were applied at the rate of 1000, 163, 170, 100 and 6 kg/ha, respectively. After final land preparation, full doses of P, K, S, Zn, B and one third of N were mixed thoroughly into the soil. The rest amount of N was applied at 21 and 53 days after seedlings emergence split into two equal amounts. The seeds of the selected genotypes were sown on 23 November 2015 in rows of 20 cm apart, at the rate of 120 kg per ha. Recommended wheat production procedure was followed [16].

When all the plants turned brown and matured properly then the crop was harvested. The harvesting for the collection of yield data was completed on 26 March, 2016. Data were collected on the following characters: thousand grain weight (g), number of grains per spike, number of spikelet's per spike, days to anthesis, heading days, plant height (cm), days to maturity and grain yield (g/plot).

Table 1. Name and sources of the wheat genotypes

Sl. No.	Name	Source	Sl. No.	Name	Source
1	Aghrani	WRC, BARI	13	PYT-6	WRC, BARI
2	Protiva	WRC, BARI	14	PYT-11	WRC, BARI
3	Sawrav	WRC, BARI	15	PYT-12	WRC, BARI
4	Gourav	WRC, BARI	16	PYT-13	WRC, BARI
5	Shatabdi	WRC, BARI	17	PYT-14	WRC, BARI
6	Sufi	WRC, BARI	18	PYT-15	WRC, BARI

9	BARI Gom 25	WRC, BARI	21	PYT-19	WRC, BARI
10	BARI Gom 26	WRC, BARI	22	PYT-20	WRC, BARI
11	BARI Gom 27	WRC, BARI	23	PYT-21	WRC, BARI
12	BARI Gom 28	WRC, BARI	24	BAW-1135	WRC, BARI

75

76 **2.1 Data analysis**

77 R- Program version 3.2.2. was used to prepare analysis of variance and to test the differences
78 among genotypes Duncan's Multiple Range Test (DMRT).

79 **3. RESULTS AND DISCUSSION**

80 **3.1 Performance of the traits on the basis of analysis of variance**

81 The analysis of variance and mean performance of the traits viz. 1000-grain weight (g),
82 number of grains per spike, number of spikelets per spike, days to anthesis, heading days,
83 plant height (cm), days to maturity and grain yield (g/plot) are presented in the table 2 and 3
84 respectively. The results exhibit that there was significant variation among the varieties for
85 almost all the traits indicating considerable amount of genetic variation in the experimental
86 materials. So, there has a great scope for the improvement of such traits through selection.
87 The co-efficient of variation was low for most of the traits but differed from the lowest value
88 (2.03%) in plant height (cm) to the highest 7.54% in 1000-grains weight (g).

89 **3.2 Mean performance of the wheat genotypes**

90 The mean performance of the 24 wheat varieties for yield and yield related traits showed
91 significant variation. These are described below-

92 **1000-grain weight**

93 The 1000-grain weight (g) is one of the most important characters for choosing the wheat
94 cultivar. It was ranged from 60.46-39.09 (g). The highest thousand grain weight was recorded
95 in the genotype PYT-12 (60.46 g) and lowest was found in Sawrav (39.09 g) followed by the
96 genotypes Sufi, Aghrani and PYT-18 (Table 3). Also found the similar observation [17].

97 **Number of grains per spike**

98 The number of grains per spike ranged was from 54.43-40.33. The maximum number of
99 grains per spike was recorded in the genotypes BARI Gom 26 (54.43) followed by the
100 genotype PYT-15 (53.43). Sawrav (52.07) while the lowest from the Protiva (40.33) followed

104 grains per spike [19] and [20].

105 **Number of spikelet per spike**

106 A wide range of variation was found among the genotypes in relation to number of spikelet
107 per spike. It was ranged from 22.50-16.80. PYT-16 (22.50) produced the highest spikelets per
108 spike which was statistically similar to Protiva. On the contrary, Gourav (16.80) followed by
109 PYT-19 and BARI Gom 26 produced the lowest spikelets per spike (Table 3).

110 **Days to anthesis**

111 Significant variation in respect of days to anthesis was observed among the selected varieties
112 indicating the presence of wide variability. Days to anthesis of genotypes ranged from 84.00-
113 74.33. The highest anthesis day was recorded in the variety BARI Gom 25 (84.00)
114 statistically similar with BARI Gom 26, Shatabdi, Sufi, Bijoy, Protiva, Sawrav, Gourav,
115 PYT-6, PYT-11, PYT-12, PYT-13, PYT-14, PYT-16 and PYT-21. On the other hand, lowest
116 days to anthesis was found in the genotype PYT-19 (74.33) which also statistically similar
117 with PYT-21, PYT-15, BAW-1135 and Aghrani (Table 3). The Bangladesh Agriculture
118 Research Institute (BARI) developed the wheat genotypes which are taking maximum days to
119 anthesis [17].

120 **Days to heading**

121 Heading days is an important character in Bangladesh condition. The genotypes which head
122 later are exposed to high temperature and grains become shriveled. So early heading is
123 important. Days to heading showed significant variation among the genotypes. It ranged from
124 76.00-62.33 among the genotypes. The highest heading days was recorded in the genotype
125 Shatabdi (76.00) followed by the genotype PYT-28 (75.67) and PYT-20 (75.00) those are
126 statistically similar (Table 3). On contrast, the lowest heading days was found in the genotype
127 PYT-19 (62.33). Revealed the same observation and reported that delay head is related to
128 shriveling of wheat grain [21].

129 **Plant height**

130 Plant height is a yield contributing trait that related to the food source of wheat plant. Semi
131 dwarf plant height in wheat is preferred because they suffer from source limitation and a tall
132 type suffers from lodging. BARI Gom 25, BARI Gom 28, PYT-6, PYT-12, PYT-13 and

137 plant types are the desirable one [22].

138 **Days to maturity**

139 For identification of the early maturing genotypes, days to maturity are important. The
140 variation in days to maturity among the different genotypes was found to statistically
141 significant (Table 2). A wider range of variation was observed among the genotypes those
142 ranked from 100.7-111.00 days (Table 3). Among the genotypes BAW-1135 (100.7) matured
143 earlier followed by PYT-19 (101.0), PYT-18 (102.7), Aghrani (103.3), Gourav (103.7) and
144 PYT-18 (104.3) those are statistically different from other genotypes for the highest value.
145 Therefore, these genotypes could be considered as promising for breeding early matured
146 wheat genotypes. Narrated that early mature genotypes are escaper from different
147 environmental stresses especially in south Asian countries where short winter season prevails
148 [23].

149 **Yield per plot**

150 Grain yield per plot is the ultimate goal for a breeding programme. Wide range of variation
151 was found among the genotypes for yield per plot and ranged from 4460.0-5813.0 (g). The
152 genotype PYT-15 was the best performer considering yield per plot (5813.0 g) and
153 statistically similar with PYT-20, PYT-18, PYT-21, BARI Gom 26, PYT-13 and Shatabdi
154 were also the high yielding genotypes those are statistically likewise with the highest
155 performer genotypes. On the other hand, PYT-16 (4460.0 g) was lower performer which
156 statistically alike with Sawrav, PYT-6, Prodip, Aghrani, Bijoy, Gourav, BARI Gom 27,
157 BARI Gom 25 and BAW-1135. Stated variation for yield and yield contributing traits in
158 wheat [22]. Found remarkable variation in wheat yield per plant [23].

159 **4. CONCLUSION**

160 In the present experiment, the variety PYT-15, BARI Gom 25 and PYT-12 performed better
161 in relation to yield and yield contributing traits. So, that it is recommended for further
162 popularization of these genotypes in Bangladesh especially in northern parts. Since the
163 experiment is one site one season experiment, to generate more reliable information on
164 performance of genotypes across location and year further studies using combination of
165

Items	df	grams weight (g)	grains per spike	spikelet per spike	Days to anthesis	Days to heading	height (cm)	Days to maturity	Yield per plot (g)
Replication	3	0.38 ^{NS}	25.45**	1.78 ^{NS}	6.11**	11.62**	51.14**	38.15**	8.19**
Genotypes	24	4.77**	4.38**	5.82**	2.27**	13.65**	21.96**	1.99 ^{NS}	3.88*
Error	72	13.766	7.704	0.794	13.007	2.070	4.300	10.536	93532.548
Coefficient of Variation		7.54%	5.82%	4.59%	4.56%	2.05%	2.03%	3.06%	6.04%

167

168 **** and * indicates significant at 0.01 and 0.05 level of probability, respectively; NS**

169 **means not significant**

Table 3: Mean performance of different traits of wheat genotypes

Genotype	1000-grain weight(g)	Number of grains per spike	Number of spikelet per spike	Days to anthesis
Aghrani	43.38 efg	51.40 abcd	18.50 f-h	75.67 c-e
Protiva	48.96 bcde	40.33 i	21.37 ab	78.00 a-e
Sawrav	39.09 g	52.07 abc	20.20 b-f	77.00 a-e
Gourav	46.54 c-f	48.20 b-g	16.80 i	77.33 a-e
Shatabdi	47.39 b-f	47.27 c-h	19.67 c-g	81.67 a-d
Sufi	41.54 fg	48.60 b-g	19.60 c-g	76.00 c-e
Bijoy	50.12 b-e	42.77 hi	19.13 c-h	80.67 a-e
Prodip	52.47 bc	49.57 a-f	20.57 bc	81.33 a-e
BARI Gom 25	53.26 bc	44.87 e-i	20.63 bc	84.00 a
BARI Gom 26	47.36 b-f	54.43 a	17.97 g-i	79.33 a-e
BARI Gom 27	52.25 bc	46.60 c-h	18.43 gh	76.67 b-e
BARI Gom 28	46.04 c-f	47.07 c-h	19.27 c-h	75.33 de
PYT-6	51.20 b-d	46.00 d-h	20.40 b-d	82.00 a-d
PYT-11	46.45 c-f	48.07 b-h	19.13 c-h	83.67 ab
PYT-12	60.46 a	46.77 c-h	19.30 c-h	82.00 a-d
PYT-13	54.30 b	48.40 b-g	18.40 gh	82.67 a-c
PYT-14	54.31 b	49.87 a-e	20.27 b-e	81.67 a-d
PYT-15	48.90 b-e	53.43 ab	18.60 e-h	75.00 de
PYT-16	51.97 b-d	44.57 e-i	22.50 a	81.67 a-d
PYT-18	44.90 d-g	44.33 f-i	19.07 c-h	79.33 a-e
PYT-19	52.17 b-d	48.47 b-g	17.83 hi	74.33 e
PYT-20	47.45 b-f	49.43 a-f	19.37 c-h	81.67 a-d
PYT-21	53.15 bc	48.63 b-g	20.43 bc	75.33 de
BAW-1135	47.45 b-f	43.20 g-i	18.67 d-h	75.33 de
LSD (0.05)	6.098	4.562	1.464	5.927
Min	39.09	40.33	16.80	74.33
Max	60.46	54.43	22.50	84.00

Table 3: Mean performance of different traits of wheat genotypes

Genotype	Days to heading	Plant height(cm)	Days to maturity	Yield per plot(g)
Aghrani	67.67 f-h	97.58 gh	103.3 b-e	4633.0 d-f
Protiva	71.00 c-e	116.1 a	108.7 a-c	5073.0 b-e
Sawrav	72.67 bc	100.7 fg	108.0 a-c	4583.0 ef
Gourav	67.33 gh	97.17 gh	102.7 c-e	4863.0 c-f
Shatabdi	76.00 a	107.6 b-d	111.0 a	5283.0 a-c
Sufi	70.00 c-g	109.5 b	106.0 a-e	5227.0 b-d
Bijoy	70.67 c-e	108.4 bc	108.7 a-c	4853.0 c-f
Prodip	71.00 c-e	105.0 c-e	109.7 ab	4603.0 ef
BARI Gom 25	69.00 e-h	104.1 d-f	107.0 a-e	4920.0 c-f
BARI Gom 26	69.67 d-g	98.32 gh	109.3 ab	5393.0 a-c
BARI Gom 27	67.67 f-h	96.05 h	106.3 a-e	4903.0 c-f
BARI Gom 28	71.33 c-e	103.0 ef	105.0 a-e	5143.0 b-e
PYT-6	70.00 c-g	102.7 ef	105.3 a-e	4583.0 ef
PYT-11	72.33 cd	97.43 gh	107.0 a-e	5133.0 b-e
PYT-12	69.67 d-g	104.1 d-f	106.3 a-e	5117.0 b-e
PYT-13	69.33 e-g	102.7 ef	104.7 a-e	5283.0 a-c
PYT-14	70.67 c-e	98.57 gh	105.7 a-e	5167.0 b-e
PYT-15	67.33 gh	106.5 b-e	107.3 a-d	5813.0 a
PYT-16	70.33 c-f	96.50 h	107.0 a-e	4460.0 f
PYT-18	75.00 ab	104.1 d-f	104.3 b-e	5440.0 a-c
PYT-19	62.33 i	97.03 gh	103.7 b-e	5063.0 b-e
PYT-20	75.67 a	107.2 b-d	108.3 a-c	5637.0 ab
PYT-21	67.33 gh	95.23 hi	101.0 de	5433.0 a-c
BAW-1135	66.33 h	92.02 i	100.7 e	5003.0 c-f
LSD (0.05)	2.365	3.408	5.335	502.6
Min	62.33	92.02	100.7	4460.0
Max	76.00	116.1	111.0	5813.0

REFERENCES

1. Rauf M, Munir M, Hassan M, Ahmad M, Afzal M. Performance of wheat genotypes under osmotic stress at germination and early seedling growth stage. *African. J. Agric. Res.*, 2007; 6: 971-975.
2. Abd-El-Haleem SHM, Reham MA, Mohamed SMS, Abdel-Aal ESM, Sosulski FW, Hucl P. Origins, characteristics and potentials of ancient wheat's. *Cereal Foods World*. 1998; 43: 708–715.
3. Hamam, KA, Sabour A, Khaled GA. Stability of wheat genotypes under different environments and their evaluation under sowing dates and nitrogen fertilizer levels, *Australian. J. Bas. Appl. Sci.*, 2009; 3(1): 206-217.
4. Shewry PR. The HEALTHGRAIN programme opens new opportunities for improving wheat for nutrition and health. *Nutrition Bulletin*. 2009; 34(2): 225–231.
5. Karim MA, Awal MA, Akhter M. Forecasting of wheat production in Bangladesh. *Bangladesh J. Agric. Res.*, 2010; 35(1): 17-28.
6. Yagdi K. A Research on Determination of Stability Parameters of Bread Wheat (*Triticum aestivum*L.) Cultivars and Lines Grown in Bursa Province. *J. Agric. Faculty Uludag Univ.*, 2002; 16:51-57.
7. Agrawal PK, Gupta HS. Enhancement of nutritional quality of cereals using biotechnological options. In P. S. Kendurkar, G. P. Srivastava, M. Mohan & Vajpeyi (Eds.), *Proceeding of ICPHT*; 2006.
8. Shewry PR. Improving the protein content and composition of cereal grain. *J. Cereal Sci.* 2007; 46: 239–250.
9. Simmonds DH. Inherent Quality Factors in Wheat. *Wheat and Wheat Quality in Australia*. Australia Wheat Board, Melbourne. 1989; pp. 31–61.
10. Adams ML, Lombi E, Zhao FJ, McGrath SP. Evidence of low selenium concentrations in UK bread-making wheat grain. *J. Sci. Food and Agric.* 2002; 82: 1160–1165.
11. Siahpoosh MR, Assad MT, Emam Y, Saidi A. Implication of four selection indices in wheat cultivars (*Triticum aestivum* L.) for increasing the grain yield, *Indian J. Genetics*. 2001; 32: 219-236.
12. Asif M, Mujahid I, Ahmad NS, Kisna M, Mustafa SZ. Determining the direct selection criteria for identification of high yielding lines in bread wheat (*Triticum aestivum* L.). *Pak J. Biol. Sci.*, 2001; 6: 48-50.
13. Singh T, Balyan HS. Relative efficiency of various single plant selection criteria and F3 generation yield testing in wheat (*Triticum aestivum* L.), *Indian J. Genetics*. 2001; 63: 24-29.
14. Chandra S, Nigam SN, Cruickshank AW, Bandopadhyaya A, Harikrishna S. Selection index for identifying high yielding groundnut genotypes in irrigated and rainfed environments, *Annual Appl. Biol.*, 2003; 143: 303-310.

15. Nadeem M. Growth, radiation use efficiency and yield of some new wheat cultivars under variable nitrogen rates. M.Sc. thesis. Department of Agronomy, Univ. Agri. Faisalabad; 2001.
16. Bangladesh Agricultural Research Council (BARC), Fertilizer Recommendation Guide. Bangladesh Agricultural Research Council, Dhaka, Bangladesh. 2005; 61-78.
17. Subhra M, Gupta S, Bhowmik N. Genetic variability of important quantitative characters in modern wheat (*Triticum aestivum* L.) cultivars in Tarai Region of West Bengal. Environ. and Ecol., 2008; 26(2): 683-686.
18. Khan K, Naqvi FN. Heritability of Morphological Traits in Bread Wheat Advance Lines under Irrigated and Non-Irrigated Conditions. Asian J. Agric. Sci., 2011; 3(3): 215-222.
19. Akmal M, Shah SM, Asim M. Yield to flag leaf area. Pakistan J. Bio. Sci. 2000; 3(12): 2072-2074.
20. Heidari B, Saeidi G, Sayed TBE. Factor analysis for quantitative traits and path analysis for grain yield in wheat. J. Sci. Tec. Agric. Nat. Res., 2008; 42: 135-144.
21. Sharma SN, Sharma Y. Estimates of variation and heritability of some quantitative and quality characters in durum wheat (*Triticum turgidum* L.). Acta Agron. Hungarica. 2007; 55:261-264.
22. Biju S, Malik SK. Variability studies in wheat. Int. J. Agric. Sci., 2007; 3(1):142-144.