

## The Effects of Vermicompost Application on the Yield of Lettuce Plant (*Lactuca sativa* L. var. *crispa*)

**ABSTRACT:** This research was done to determine the effect of increasing Vermicompost application on yield of lettuce (*Lactuca sativa* L. var. *crispa*) plant. For this purpose *Lactuca sativa* L. var. *crispa* cv. *Bellafigesta* lettuce kind and Riverm Company Vermicompost were used in this research. Four Vermicompost doses (I. dose: 0 kg/da II. dose: 400 kg/da, III. dose: 800 kg/da, IV. dose: 1200 kg/da) were applied to lettuce plant. According to the results, important increases of fresh weight, plant size, number of leaf, length of leaf and width for each plant and dry matter yield of plants were determined with increasing Vermicompost applications. But the effects of Vermicompost applications on some macro element (N, P, K, Ca and Mg) contents of plant were not found significant statistically.

**Key words:** Vermicompost, biological property, macro nutrient element, lettuce.

### 1. INTRODUCTION

Today an increase in the products gained from per unit area has become a necessity to satisfy mankind's need for food. On the other hand, the necessity of increase in the products brings another necessity to use more inorganic fertilizers per unit area. As a result of the excessive application of inorganic fertilizers, natural sources such as soil and water are being polluted and serious health problems are occurring.

Mostly in vegetable farming, in order to reach a high degree of productivity and maximum growth, the amount of inorganic fertilizer as the main source of nutrient is emphasized, however, usually exceeded [1,2].

It has been a well-known fact that green plants respond positively to inorganic fertilizer with nitrogen content however, nitrogen application has a limited positive effect on the crop yield. Increasing nitrogen fertilization can affect plant's agronomy, macro and micro nutrient element contents, and the quality of the product negatively. Nitrogen has an important role in the plant's vegetative development and crop yield. However, the excessive use of nitrogen fertilizer in order to increase productivity might cause the risk of nitrate accumulation [3,4].

While excessive inorganic nitrogen fertilizer causes soil pollution [5,6,7], it also causes the accumulation of harmful compounds for human health in vegetables [8]. Nevertheless, according to FAO/WHO [9] nitrate generates toxic effect if it exceeds 5 mg for each kg of human body. For this reason, in vegetable farming the management of nitrogen fertilization should carefully be programmed.

The scientists who are seeking a solution for this problem put forward that organic fertilizer applications should be increased in following years. Because while organic fertilizers are manure materials and nutrient sources, they can reform the degenerations in soil and water caused by inorganic fertilizers. In recent years, the use of organic fertilizers is increasing especially in vegetable farming.

Organic fertilizers are quite popular in farming recently. The use of these fertilizers provides significant rise in the crop yield and quality of the plant. In a research [10], 50, 100, 200, 400 and 600 mg/kg humic acid were applied to corn and the effects of this application on fresh and dry weight of the plant and on the amount of

protein were examined. A significant increase in biological characteristics of corn was observed upon the application of humic acid doses above 200 mg/kg.

In a research conducted to identify the effects of increasing doses of vermicompost application to potato plant on the crop yield of the plant, 0, 4.5, 9 and 12 tones/da of vermicompost were applied. As a result, it was concluded that maximum plant height, dry weight of leaf and plant stem, dry and wet weight of tuber, total tuber weight, the number of tubers, tuber diameter, tuber nitrogen percentage, tuber potassium percentage and such biological parameters were obtained from the application of 12 tones/da of vermicompost [11].

A research by Alam et al., [12] examines the crop yield of potato plant and its characteristics upon the application of increasing doses of vermicompost. Certain increases in some agronomic features such as potato's tuber diameter, tuber weight, the crop yield gained from per area, and leaf width index were observed once vermicompost was applied.

In a study held in Azerbaijan the effects of the application of 2, 4, 6 tones/da of vermicompost on red onion plant (*Allium cepa* L.) were analyzed. The highest crop yield of the onion plant in terms of protein and ascorbic acid content was determined from the fields on which 6 tones/ha of vermicompost were applied [13].

In Bangladesh the effects of various doses of vermicompost on cauliflower farming have been investigated. 0, 1.5, 3 and 6 tones/ha doses of vermicompost were applied to the plants, and certain biological features were measured such as maximum plant height, the number of leaves, fruit width, fruit height, total weight, commercial weight, and crop yield of stem. According to the results, the maximum yield was obtained from the field on which 6 tones/ha of vermicompost was applied [14].

The effects of increasing doses vermicompost application on lettuce (*Lactuca sativa* L. var. *crispa*) plant, it's some nutrient element content and some agronomic properties of the plant was investigated in this research.

## 2. MATERIALS AND METHODS

In the research, *Lactuca sativa* L. var. *crispa* cv. *Bellafiesta* was used, which is a type of lettuce plant. Two seeds for each pot were planted and peat was used as the production ground (Klasmann-Deilmann, potground H, Germany). When they gained 3-4 leaves 30 days after the plantation and they were transmitted to their permanent pots. Randomized blocks were designed as 3 replications on the experimental design, and there were 108 plants in total, 9 in each parcel. Total experimental area was 75 m<sup>2</sup>. The vermicompost was applied to the plants (1<sup>st</sup> dose: 0 g/m<sup>2</sup>, 2<sup>nd</sup> dose: 400 g/m<sup>2</sup>, 3<sup>rd</sup> dose: 800 g/m<sup>2</sup>, 4<sup>th</sup> dose: 1200 g/m<sup>2</sup>) right after the plantation. Some chemical properties of the vermicompost used in the experiment were presented in Table 1 below.

**Table 1. Some chemical properties of Vermicompost**

pH	7.60
Org. Matter, %	51.80
Total hümic+ fulvic acid, %	46.10
Org. C %	27.80

<b>Total N, %</b>	1.50
<b>Soluble P<sub>2</sub>O<sub>5</sub>, %</b>	0.20
<b>Soluble K<sub>2</sub>O, %</b>	1.10
<b>Soluble CaO, %</b>	0.26
<b>Soluble MgO, %</b>	0.13

30 days after the plantation the plants were harvested and plant height (cm), plant diameter (cm), leaf size (cm), leaf width (cm), root height (cm), plant weight (gr), and number of leaves were measured. Dry material content of the plants was obtained by washing them with pure water and drying them in 65 °C for 48 hours. After the dry weight was determined necessary elemental analyses were conducted on grained samples by using ICP-OES device [15].

### 3. RESULTS AND DISCUSSION

#### 3.1. The Effects of Vermicompost Application on Some Biological Properties of Lettuce Plant (*Lactuca Sativa* L. var. *crispa* cv. *Bellafigesta*)

The effects of increasing doses of vermicompost application on the height, diameter, number of leaves, leaf size, leaf width and the weight of lettuce plant (*Lactuca sativa* L. var. *crispa* cv. *Bellafigesta*) were presented on Table 2. The effects on the biological properties of the lettuce plant vary depending on the amount of doses (Table 2).

**Table 2. The effect of Vermicompost on some biological properties of lettuce plant.**

Dose	Height (cm)	Diameter (cm)	Root length (cm)	Num. of leaf	Length of leaf (cm)			Leaf width (cm)			Plant weight (gr)
					Int.	Med	Ext.	Int.	Med	Ext.	
I	35.5	28.8	16.6	19.7	9.0	13.6	14.8	6.9	12.6	13.2	108.1
II	34.9	31.0	16.7	23.6	9.3	15.3	16.8	7.3	14.4	15.7	130.1
III	34.6	31.4	16.0	22.8	10.0	15.2	17.3	8.3	14.3	16.9	133.2
IV	32.7	31.7	14.8	21.8	8.4	15.0	17.0	7.3	13.6	16.8	125.4

Significant rise in the plant width, the number of leaves, leaf size, leaf width, and the plant height was observed upon the increasing doses of vermicompost application. On the other hand, with the vermicompost application decrease in the height and root size of the lettuce plant was determined comparing to the control group. These results are concordant to the research indications by Yourtchi et al. [11]. In the statistical evaluation, the increases were statistically significant at the level of 5 %.

#### 3.2. The Effects of Vermicompost Application on Some Macro Nutrient Element (N, P, K, Ca, Mg) Contents of The Lettuce Plant (*Lactuca sativa* L. var. *crispa* cv. *Bellafigesta*)

The effects of vermicompost application on the some macro nutrient element contents of the lettuce plant were presented in average of three replications in Table 3.

**Table 3. The effect of Vermicompost application on some macro nutrient element (N, P, K, Ca, Mg) contents of lettuce plant, %.**

Doses	N	P	K	Ca	Mg
I	5.43	0.92	10.95	1.42	0.21
II	5.30	0.81	9.24	1.25	0.18
III	5.09	0.71	9.86	1.24	0.18
VI	5.11	0.67	9.61	1.42	0.20

As it can be seen on Table 3, no significant mutation was discovered in N, P, K, Ca and Mg contents of the plant upon the increasing doses of vermicompost application. These effects were not considered as significant. The short probation period which is 30 days and plants' inability to obtain enough nutrient elements from soil could be demonstrated as the reason.

In as research held in agricultural fields, a relationship was determined between protein and ascorbic acid content of red onion plant (*Allium cepa* L.) and increasing doses of vermicompost [13].

A study by Yourtchi et al. [11] reveals that a significant increase in nitrogen, phosphorus, and potassium contents of the potato plant were determined with the application of increasing doses of vermicompost.

#### 4. CONCLUSION

This study exposes that the increasing doses of vermicompost application multiply the crop yield, wet weight and diameter, number of leaves, size and width of leaves of the lettuce plant. However, the plant's nitrogen, phosphorus, potassium, calcium, and magnesium contents do not receive an important fluctuation.

Most of the agricultural fields in Turkey have inadequacy of organic matter amount. For this reason the increase in organic matter amount of agricultural fields has become a necessity. Thus, the use of vermicompost in agriculture will increase the organic matter amount of soils and constitute a significant input in some quality parameters of plants.

The common use of vermicompost in agriculture is expected to reform the pollution and deterioration of nutrient balance of soils caused by chemical fertilizers. Vermicompost will also be able to be used for different purposes and outputs in agricultural production on different regions, plants, soils and climate conditions.

This study proves that vermicompost can be used in agricultural production by exhibiting the example of lettuce farming. It has been a well-known fact that there is an inadequacy of organic matter amount in most of the agricultural fields of Turkey. With this study, it has been revealed that vermicompost application could be an alternative source of organic matter in eliminating the inadequacy of organic matter in

agricultural fields.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

- [1] Adediran, A.J., Taiwo B.L., Akande O.M., Sobule A.R., Idowu J.O. Application of organic and inorganic fertilizer for sustainable maize and cowpea yields in Nigeria. *J. Plant Nutrition*, 2004, 27: 1163– 1181.
- [2] Naeem, M., Iqbal J., Bakhsh M.A.A. Comparative study of inorganic fertilizers and organic manures on yield and yield components of Mungbean (*Vigna radiat* L.). *J. Agric. Soc. Sci.*, 2006, 2: 227– 229.
- [3] Addiscott, T.M. Nitrate. *Agriculture and Environment*. 2005. Wallingford, Oxfordshire, UK, CABI Publishing.
- [4] Lemaire, G., Gastal, F.F. Quantifying crop responses to nitrogen deficiency and avenues to improve nitrogen use efficiency. In: Sadras V, Calderini D, (Eds.), *Crop Physiology*, Academic Press, USA, 2009, pp: 171– 211.
- [5] Gollany, H.T., Molina J.E., Clapp C.E., Allmaras R.R., Layese M.F., Baker J.M., Cheng, H.H. Nitrogen leaching and denitrification in continuous corn as related to residue management and nitrogen fertilization. *Environmental Management*, 2004, 33: 289– 298.
- [6] Beman, J.M., Arrigo K., Matson, P.M. Agricultural runoff fuels large phytoplankton blooms in vulnerable areas of the ocean, *Nature*, 2005, 434: 211– 214.
- [7] Zand-Parsa, S., Sepaskhah, A.R., Ronaghi, A. Development and evaluation of integrated water and nitrogen model for maize. *Agricultural Water Management*, 2006, 81: 227– 256.
- [8] Ruiz, J.M., Romero, L. Cucumber yield and nitrogen metabolism in response to nitrogen supply. *Scientia Horticulturae*, 1999, 82: 309– 316.
- [9] FAO/WHO, Toxicological evaluation of certain food additives and contaminants. Geneva, World Health Organization, Joint FAO/WHO Expert Committee on Food Additives, 1996, WHO Food Additives Series No. 35.

- [10] Doğru, A., Darçın, E.S., Tutar A., Dizman, M., Koç, Y. Potasyum humatın mısır (*Zea mays* L.) bitkisinin büyümesi üzerine etkileri. SAÜ Fen Ed. Dergisi, 2012. 14 (1): 83- 93.
- [11] Yourtchi, M.S., Hadii M.H.S., Darzi, M.T. Effect of nitrogen fertilizer and vermin-compost on vegetative growth, yield and NPK uptake by tuber of potato (*Agriacv.*). Int. J. Agric. Crop Sci. 2013, 5(18): 2033-2040.
- [12] Alam, M.N., Jahan M.S., Ali M.K., Ashraf, M.A., Islam, M.K. Effect of vermicompost and chemical fertilizers on growth, yield and yield components of potato in barind soils of Bangladesh. J. Appl. Sci. Res., 2007, 3 (12): 1879-1888.
- [13] Bai, B.A., Malakout, M.J. The Effect of different organic manures on some yield and yield quality parameters in onion. Iran Soil and Water Sciences Journal, 2007, 21 (1): 43- 53.
- [14] Jahan, F.N., Shahjalal A.T.M., Paul A.K., Mehraj, H., Uddin, A.F.M.J. Efficiency of vermicompost and conventional compost on growth and yield of cauliflower. Bangladesh Research Publications Journal, 2014, 10 (1): 33- 38.
- [15] Kacar, B., İnal, A. Bitki Analizleri. Nobel Yayın, No: 849, 2010, 659s, Ankara.