

# Evaluation of anaerobic digestate potential as organic fertilizer in improving wheat production and soil properties

## Abstract

Integrated use of synthetic and organic fertilizers is crucial to sustainable crop production and stabilization of depleting soil fertility. Keeping in view these certainties, a three year field study was undertaken to evaluate the potential of anaerobic digestate alone or in integration with chemical fertilizer for improving wheat production and soil fertility. Six treatments viz; control (with no amendment), recommended dose (RD) of chemical fertilizers (CF), anaerobic digestate (AD) on the basis of RD of N, farm manure (FM) on the basis of RD of N,  $\frac{1}{2}$  N from CF and  $\frac{1}{2}$  N from AD,  $\frac{1}{2}$  N from CF and  $\frac{1}{2}$  N from FM were applied in Randomized Complete Block Design (RCBD) with three replications. The obtained results revealed that the highest yield (grain and straw), N uptake, NUE, NAE and NRE were acquired through the utilization of application of chemical fertilizers which was statistically at par with combined application of anaerobic digestate and chemical fertilizer in all years of study while the minimum was found in control. It was concluded from the outcomes that by the utilization of anaerobic digestate alongside chemical fertilizer, half of nitrogenous fertilizer (urea) can be saved.

**Keywords:** Anaerobic digestate, farm manure, fertilizers, yield, NUE

## 1. Introduction

The sustainability of agricultural productivity is of utmost significance keeping in view the pace of total population growth. It is evaluated that the production of food should expand 70% to sustain flourishing populace, which is relied upon to reach up to 9

27 billion by 2050 (FAO, 2010). Frequent utilization of chemical fertilizers is practiced all  
28 through the world to amplify crop production with a specific end goal to satisfy the food  
29 needs of growing population (Bos et al., 2005). The prolonged utilization of chemical  
30 fertilizers and intensive agriculture have prompted deterioration of soil fertility and  
31 additionally caused environmental hazards like ground and surface water pollution  
32 from nitrate leaching (Pimentel, 1996). Lack of appropriate crop management, decrease  
33 in addition of fertility restoring inputs and unbalance nutrients application have paved  
34 the path for soils to become fragile (Ajayi et al., 2007; Mbah and Onweremadu, 2009).

35 Under current scenario, it is indispensable and dire to take suitable measures to check  
36 decrease in soil fertility and profitability. The circumstance immovably emphasis the  
37 selection of ecofriendly agricultural practices for keeping up soil fertility and getting  
38 crop production on sustainable basis. For this reason, a sustainable approach is  
39 prescribed regarding organic agriculture (Oyewole et al., 2012). Utilization of organic  
40 manures won't just be productive in diminishing unfavorable impacts of synthetic  
41 fertilizers, yet in addition will support soil fertility and productivity (Aksoy, 2001;  
42 Chowdhury, 2004). Organic manures upgrade food quantity and quality by fulfilling crop  
43 nutritional requirement with the provision of essential nutrients in a way similar to  
44 synthetic fertilizers (Liu et al., 2007; Tonfack et al., 2009; Maske et al., 2015).

45 Moreover, the costs of chemical fertilizers are going past the purchasing capacity of  
46 normal land holding farmers and furthermore sole use of either organic or chemical  
47 fertilizer isn't appropriate (Wakene et al., 2007). In this way the present circumstance  
48 requests mix of organic manures with synthetic fertilizers (Jayathilake et al., 2006). To  
49 get feasible crop production without declining and weakening soil fertility, appropriate  
50 mix of both mineral and organic manure ought to be adopted (Rekhi et al., 2000).

51 Integration of organic and mineral fertilizers will enhance absorption, distribution as  
52 well as nutrient and fertilizer use efficiency (*Orkaido, 2004; Jayathilake et al., 2006*).

53 The digestates generated by anaerobic processing of domesticated animals excrements  
54 amid biogas generation are commonly rich in macronutrients, for example, N, P, and K,  
55 and micronutrients, for example, Zn, Fe, Mo, and Mn (*De La Fuente et al., 2013*). In like  
56 manner, these digestates can possibly be utilized as organic manures and soil  
57 amendment in the agricultural land. The advantages of land utilization of digestates  
58 include change in seedling development, crop yield, and fruit/vegetable quality (*Feng et*  
59 *al., 2011; Zhang et al., 2015*). Besides, the physical, chemical, and biological properties of  
60 soil can likewise be improved (*Riva et al., 2016; Zerzghi et al., 2010*). Digestate holds  
61 significant amounts of organic matter (20 to 30%), which is necessary for our soils with  
62 low organic matter (<1%). In this way, digestate can be a best choice to be integrated  
63 with chemical fertilizers for getting optimum crop yield and to recharge soil fertility.

64 Keeping in view the significance of integrated utilization of organic and inorganic  
65 manures, the present examination was conducted to study the capability of digestate as  
66 organic manure alone and in combination with mineral fertilizers for enhancing wheat  
67 yield and soil fertility.

## 68 **2. Materials and Methods**

69 The current field study was conducted at the farm area of Soil Chemistry Section,  
70 Institute of Soil Chemistry and Environmental Sciences, Ayub Agricultural Research  
71 Institute Faisalabad, Pakistan for three consecutive years. The investigation was  
72 performed by utilizing Randomized Complete Block configuration having plot size of 5m  
73 × 7m and was replicated thrice. Nitrogen was applied at the rate of 120 kg ha<sup>-1</sup> as urea,

74 phosphorus 90 kg ha<sup>-1</sup> as single super phosphate (SSP) and 60 kg ha<sup>-1</sup> potash as sulfate  
75 of potash (SOP). Following six treatments were used in the study;

76 T<sub>1</sub> = Control (without any fertilizer)

77 T<sub>2</sub> = Recommended dose (RD) of chemical fertilizer (CF)

78 T<sub>3</sub> = Anaerobic digestate (AD) on the basis of RD of nitrogen

79 T<sub>4</sub> = Farm manure (FM) on the basis of RD of nitrogen

80 T<sub>5</sub> = ½ N from anaerobic digestate (AD) and ½ N from chemical fertilizer (CF)

81 T<sub>6</sub> = ½ N from farm manure (FM) and ½ N from chemical fertilizer (CF)

82 Calculated amount of farm manure was well mixed in the soil at seed bed preparation  
83 whereas digestate was applied through fertigation with first irrigation. Wheat cultivar  
84 Punjab-2011 was sown following recommended methods with seed rate of 124 kg ha<sup>-1</sup>  
85 and row to row distance of 22.5cm × 22.5cm.

## 86 **2.1. Soil Sampling and analysis**

87 For evaluation of initial fertility status of the field, a soil composite sample was  
88 collected. The collected soil samples were air dried, crushed and sieved through a 2 mm  
89 stainless steel sieve and before analyzing for physical and chemical characteristics  
90 (Table 1). Soil particle distribution was measured by hydrometer method (*Blake and*  
91 *Hartge*, 1986). About 250 g soil was saturated with distilled water for determining pH of  
92 soil. The paste was allowed to stand for one hour and pH was recorded by pH meter  
93 with glass electrodes using buffer of pH 4.0 and 9.0 as standard (*Mclean*, 1982). For  
94 determining E<sub>ce</sub>, extract of each soil paste was obtained by using vacuum pump and  
95 E<sub>ce</sub> was noted with conductivity meter (Corning 220). Soil organic carbon (SOC)  
96 content was estimated following the method as described by *Ryan et al.* (2001), and  
97 available phosphorus was estimated by Olsen's method (*Jackson*, 1962) using

spectrophotometer. For potassium content, soil extraction was done with ammonium acetate (1 N of pH 7.0) and potassium was determined by using PFP-7 Janway Flame photometer (Rowell, 1994). After each crop harvest, soil samples were taken and analyzed to evaluate improvement in soil physicochemical properties by organic amendments supplementation. Meteorological data for each year of the study is given in Figure 1.

## 2.2. Characterization of organic materials

The anaerobic digestate used was collected each year of the study from a biogas plant located at Chak No. 254 RB, Faisalabad. Whereas farm manure was taken from dairy farm of Ayub Agricultural Research Institute, Faisalabad. Before application, the samples of both fresh slurry and farm manure were collected and analyzed for chemical constituents by following standard methods (Table 2).

## 2.3. Plant sampling and analysis

At harvest, data regarding grain yield ( $\text{Mg ha}^{-1}$ ) and straw yield ( $\text{Mg ha}^{-1}$ ) was collected in each year of study. Area of  $9 \text{ m}^2$  was harvested from each experimental unit. The harvest of each experimental unit was labeled, sun dried and threshed separately. Grain samples were collected and dried at  $70^\circ\text{C}$  for the determination nitrogen. The dry grain samples were ground and 0.5g sample was digested with Tri-acid mixture of  $\text{HNO}_3$ - $\text{H}_2\text{SO}_4$ - $\text{HClO}_4$  for the determination of total nitrogen by Kjeldhal method (Jackson, 1962). Nitrogen uptake, nitrogen use efficiency (NUE), nitrogen recovery efficiency (NRE) and nitrogen agronomic efficiency (NAE) were calculated by using following equations as mentioned by Javid et al. (2015).

$$N \text{ uptake } (\text{kg ha}^{-1}) = \frac{N \text{ content } (\%) \text{ grain (dry matter)} \times \text{Yield } (\text{kg ha}^{-1})}{100}$$

$$NUE = \frac{\text{Wheat grain yield (kg ha}^{-1}\text{)}}{\text{N fertilizer applied (kg ha}^{-1}\text{)}}$$

$$NRE = \frac{\text{N uptake (kg ha}^{-1}\text{) in fertilized plot} - \text{N uptake in control (kg ha}^{-1}\text{)}}{\text{N fertilizer applied (kg ha}^{-1}\text{)}}$$

$$NAE = \frac{\text{Yield in fertilized plot} - \text{yield in control (kg ha}^{-1}\text{)}}{\text{N fertilizer applied (kg ha}^{-1}\text{)}}$$

## 2.4. Statistical and economic analysis

The data regarding various traits were subjected to analysis of variance to test the significance of treatments using *Statistix 8.1*<sup>®</sup> (Analytical Software, Tallahassee, USA) and treatment means were compared using least significant difference (LSD) (Steel et al., 1997). A benefit-cost analysis was conducted to estimate the economic feasibility of different organic amendments to increase vegetable production and net economic returns as described by CIMMYT (1988).

## 3. Results and Discussion

The present three year investigation was conducted for the evaluation of anaerobic digestate potential as fertilizer alone and in mix with chemical fertilizers. The three year pool data in regards to wheat yield (Figure 2) portrayed that highest yield (4.33 and 7.01 Mg ha<sup>-1</sup> for grain and straw individually) was acquired with the use of prescribed quantity of chemical fertilizers which was at par with the yield got in plots accepting digestate in combination with chemical fertilizers in 1:1 premise (4.16 and 6.46 Mg ha<sup>-1</sup> for grain and straw separately). Both treatments displayed similar performance in each year of study period (Table 3, 4). The treatments with sole application of either farm manure or digestate on the basis of recommended dose of N were statistically at par with each other. The highest yield with the application of chemical fertilizer might be attributed to the delivery of nutrients in soluble form in the soil solution, which become available promptly for plants to take up and flourish (Aziz et al., 2010). The improved

nutrient and soil moisture availability could be the reason of noteworthy increase in yield in response to integration of organic and inorganic fertilizers. The propensity of organic fertilizers for improving soil physicochemical properties as well as nutrient supplying capability might also be the reason for better yield. The lowest yield in case of farm manure or digestate alone compared to that of chemical fertilizers and their integration might be due to slow release of nutrients needed by plants for their growth and development (Powon et al., 2004). The findings of the current study are in agreement to the observations of Shaheen et al. (2017) who found higher soybean dry matter yield with the application of chemical fertilizers followed by integrated use of inorganic and organic fertilizer, whereas Noreen and Noreen (2012) found non-significant wheat grain and straw yield with the supplementation of chemical fertilizers alone and combined application of chemical fertilizers (75%) and farm manure (25%). Similarly Muhammad et al., (2009) and Ayoola and Makinde (2008) found highest corn cob yield with the application of chemical fertilizers followed by combination of synthetic fertilizers and organic amendments.

### 3.1. Nitrogen uptake and use efficiency

The data regarding nitrogen uptake and use efficiency (Figure 3) illustrated that maximum nitrogen uptake ( $85.7 \text{ kg ha}^{-1}$ ) was obtained in treatment receiving nitrogen from chemical source followed by treatment with integration of chemical fertilizer and digestate ( $77.2 \text{ kg ha}^{-1}$ ). Similar trend was obtained for nitrogen use efficiency, nitrogen agronomic efficiency and nitrogen recovery efficiency with values of 36.1, 19.5 and 57.4  $\text{kg kg}^{-1}$  correspondingly. Chemical fertilizers contain nutrients in readily available form which release instantly upon application for plant uptake (Aziz et al., 2010). This is the reason why the nitrogen uptake as well as use efficiencies were maximum in case of

application of nitrogen from synthetic source. The greater nitrogen uptake and use efficiency in case of integrated use of organic and inorganic fertilizer than sole application of organic amendments may be ascribed to the sustainable and prolonged supply of nutrients (*Singh and Singh, 2000; Aziz et al., 2010*) more so the application of organic fertilizer alone results in the slow release from their decomposition by soil microbes (*Mahajan et al., 2008; Ghemam and Mourad, 2013*).

The findings of the present study are consistent with the results obtained by *Haile et al., (2012)* who found increase in application rate of N resulted in significant improvement in N uptake by wheat crop. Similar results were also found by *Shaheen et al., (2017)* and *Islam and Munda (2012)*. However, findings that differ from this study were obtained by *Naing et al., (2010)* who reported significantly higher agro-physiological N and P use efficiencies for rice with organic-inorganic mixed fertilizers compared to chemical fertilizers or organic fertilizer alone. Whereas *Hossain et al., (2010)* found higher nutrient uptake with the application of farm yard manure alone compared to that of no fertilizer and inorganic fertilizers application.

### **3.2. Improvement in soil chemical properties**

The findings regarding post-harvest soil chemical analysis (Table 5) depicted that there is a little variation in soil pH and E<sub>c</sub> during the studied period. The contents of organic matter improved significantly with the addition of organic fertilizer either alone or in combination with chemical fertilizers. The maximum contents (0.96%) were seen in the treatment receiving combined application of organic and inorganic fertilizer while minimum was observed in control without any amendment. Similarly, significant improvements were observed for phosphorus and potassium contents in the current study. The highest amount of P and K (14.63 and 245 mg kg<sup>-1</sup> respectively) were obtained with the application of digestate and chemical fertilizer together. The non-



significant change in soil pH and ECe in this study might be due to highest buffering capacity of the soil. Similar to the current study, *Yadav et al.*, (2002) found no appraisable change in pH and ECe in response to the application of organic amendments. The improved contents of organic matter, phosphorus and potassium contents could be attributed to the sustained supplementation of organic amendments over the studied period and their subsequent residual effects on the soil properties that last for several years after their application due to their slow mineralization. The findings of our study are in accordance with the results obtained by *Enujeke et al.*, (2013) in a study undertaken for evaluation of residual effect of organic manure and chemical fertilizer on soil properties.

### **3.3. Economic analysis**

The economic analysis (Table 6) depicted that the maximum cost benefit ratio was obtained with the combined application of chemical fertilizer and digestate followed by the integration of chemical fertilizers and farm manure while minimum was in case of recommended dose of chemical fertilizer among all treatments except control.

## **4. Conclusions**

The discoveries of the present investigation uncovered that anaerobic digestate can possibly enhance crop production and soil fertility. Wheat yield, nutrient uptake and soil chemical properties portrayed critical contrasts with the application of various blends of organic and inorganic fertilizers. In any case, the best mix was anaerobic digestate alongside inorganic fertilizers. It not only brought about a significant increase in nutrient uptake and yield of wheat but also proved to be cost effective and monetary feasible. Furthermore, it was concluded that about half of nitrogenous fertilizer can be saved with the adoption of chemical fertilizer and digestate integration. Besides, it was

inferred that about half of nitrogenous fertilizer can be spared with the appropriation of chemical fertilizers e and digestate integration.

## References

*Ajayi, O.C., Akinnifesi, F.K., Sileshi, G., Chakeredza, S.* (2007). Adoption of renewable soil fertility replenishment technologies in the southern African region: Lessons learnt and the way forward. *Nat. Resour. Forum.* 31, 306-317.

*Aksoy, U.* (2001). Ecological farming. II. In Proceedings of the Ecological Farming Symposium, Antalya, Turkey, 14–16. *Asian J. Agri. Biol.* 5(2), 60-69.

*Ayoola, O.T., Makinde, E.A.* (2007). Complementary organic and inorganic fertilizer application: influence on growth and yield of cassava/maize/melon intercrop with a relayed cowpea. *AJBAS.* 1(3), 187-192.

*Aziz, T., Ullah, S., Sattar, A., Nasim, M., Farooq, M., Khan, M.M.* (2010). Nutrient availability and maize (*Zea mays* L.) growth in soil amended with organic manures. *Int. J. Agric. Biol.* 12, 621–624.

*Blake, G.R., Hartge, K.H.* (1986). Bulk density. In: A. Klute (ed.), *Methods of Soil Analysis, Part 1: Physical and Mineralogical Methods.* American Society of Agronomy, Madison, WI, pp. 363-375.

*Bos, C., Juillet, B., Fouillet, H., Turlan, L., Dare, S., Luengo, C., N'tounda, R., Benamouzig, R., Gausseres, N., Tome, D., Gaudichon, C.* (2005). Postprandial metabolic utilization of wheat protein in humans. *Am. J. Clin. Nutr.* 81, 87-94.

*Chowdhury, R.* (2004). Effects of chemical fertilizers on the surrounding environment and the alternative to the chemical fertilizers. *IES-ENVIS News.* 7, 4–5.

*CIMMYT,* (1988). *An Economic Training Handbook,* Economic Program, CIMMYT, Mexico.

- 238 *De La Fuente, C., Alburquerque, J.A., Clemente, R., Bernal, M.P.* (2013). Soil C and N  
239 mineralization and agricultural value of the products of an anaerobic digestion  
240 system. *Biol. Fert. Soils.* 49, 313-322.
- 241 *Enujeke, E.C., Ojeifo, I.M., Nnaji, G.U.* (2013). Residual effects of organic manure and  
242 inorganic fertilizer on maize grain weight and some soil properties in Asaba area  
243 of delta state. *IJABBR.* 3(3), 433-442.
- 244 *Feng, H., Qu, G., Ning, P., Xiong, X., Jia, L., Shi, Y., Zhang, J.* (2011). The Resource Utilization  
245 of Anaerobic Fermentation Residue. *Procedia Environ. Sci.* 11, 1092-1099.
- 246 *FAO.* (2010). Production Year Book; Food and Agriculture Organization of the United  
247 Nations: Rome, Italy.53, pp. 132–133.
- 248 *Haile, D., Dechassa, N., Ayana, A.* (2012). Nitrogen use efficiency of bread wheat: Effects  
249 of nitrogen rate and time of application. *J. Soil Sci. Plant Nutr.* 12(3), 389-410.
- 250 *Hossain, A.T., Rahman, F., Saha, P.K., Solaiman, A.R.M.* (2010). Effects of different aged  
251 poultry litter on the yield and nutrient balance in boro rice cultivation. *BJAR.*  
252 35(3): 497-505.
- 253 *Hossain, N., Kibria, M.G., Osman, K.T.* (2012). Effects of poultry manure, household waste  
254 compost and inorganic fertilizers on growth and yield. *J. Pharm. Biol. Sci.* 3(2),  
255 38-43.
- 256 *Islam, M., Munda, G.C.* (2012). Effect of organic and inorganic fertilizer on growth,  
257 productivity, nutrient uptake and economics of maize (*Zea mays* L.) and toria  
258 (*Brassica campestris* L.). *Agric. Sci. Res. J.* 2(8), 470-479.
- 259 *Jackson, M.L.* (1962). Soil chemical analysis. Prentice Hall: Inc. Englewood Cliffs, New  
260 Jersey, USA.

- 261 Javid, S., Majeed, A., Sial, R.A., Ahmad, Z.A., Niaz, A., Muhmood, A. (2015). Effect of  
262 phosphorus fertigation on grain yield and phosphorus use efficiency by maize  
263 (*Zea mays* L.). *J. Agric. Res.* 53(1), 37-47.
- 264 Jayathilake, P.S., Reddy, I.P., Srihari, D., Reddy, K.R. (2006). Productivity and soil fertility  
265 status as influenced by integrated use of N-fixing biofertilizers, organic manures  
266 and inorganic fertilizers in onion. *J. Agric. Sci.* 2(1), 46 - 58.
- 267 Liu, B., Gumpertz, M.L., Hu, S., Ristaino, J.B. (2007). Long-term effects of organic and  
268 synthetic soil fertility amendments on soil microbial communities and the  
269 development of southern blight. *Soil Biol. Biochem.* 39, 2302-2316.
- 270 Mahajan, A., Bhagat, R.M., Gupta, R.D. (2008). Integrated nutrient management in  
271 sustainable rice-wheat cropping system for food security in India. *SAARC. J. Agric.*  
272 6(2), 29-32.
- 273 Maske, S.N., Munde, G.R., Maske, N.M. (2015). Effect of manures and fertilizer on brinjal  
274 (*Solanum melongena* L.) C.V. Krishna. *Bioinfolet.* 12, 678-679.
- 275 Mbah, C.N., Onweremadu, E.U. (2009). Effect of organic and mineral fertilizer inputs on  
276 soil and maize grain yield in an acid ultisol in Abakaliki-South Eastern Nigeria.  
277 *Am. Eurasian J. Agron.* 2(1), 7-12.
- 278 Mclean, E.O. (1982). Soil pH and lime requirement, In: A.L. Page, R.H. Miller and D.R.  
279 Keeney (eds.) *Methods of Soil Analysis part 2: Chemical and Microbiological*  
280 *Properties*, 2nd Ed. American Society of Agronomy 9, Madison, WI, USA, pp.199-  
281 209.
- 282 Muhammad, D., Khattak, R.A. (2009). Growth and nutrient concentrations of maize in  
283 press mud treated saline-sodic soils. *Soil Environ.* 28 (2), 145-155.
- 284 Noreen, F., Noreen, S. (2014). Effect of different fertilizers on yield of wheat. *IJSR.* 3(11):  
285 1596-1599.

- 286 *Orkaido, O. (2004).* Effects of nitrogen and phosphorus fertilizers on yield and yield  
287 components of Maize (*Zea Mays L.*) on black soils of Regede, Konso, MSc. Thesis,  
288 Alemaya University, Alemaya, Ethiopia.
- 289 *Oyewole, C.I., Opaluwa, H., Omale, R. (2012).* Response of Tomato (*Lycopersicon*  
290 *esculentum*): Growth and yield, to rates of mineral and poultry manure  
291 application in the guinea savanna agro-ecological zone in Nigeria. *J. Biol. Agric.*  
292 *Health Care.* 2, 44-56.
- 293 *Pimentel, D. (1996).* Green revolution and chemical hazards. *Sci. Total Environ.* 188, 86-  
294 98.
- 295 *Powon, M.P., Mwaja, V., Aguyo, J.N. (2004).* The effect of potassium, phosphorus and  
296 farmyard manure on growth and yield of potato (*Solanum tuberosum L.*). Paper  
297 Presented At the 9th Triennial Symposium of International Society For Tropical  
298 Root- Africa Branch (ISTRC-AB).
- 299 *Rekhi, R.S., Benbi, D.K., Bhajan, S. (2000).* Effect of fertilizers and organic manures on  
300 crop yields and soil properties in rice-wheat cropping system: In Long-term Soil  
301 Fertility Experiments in Rice-Wheat Cropping Systems, pp.1-6.
- 302 *Riva, C., Orzi, V., Carozzi, M., Acutis, M., Boccasile, G., Lonati, S., Tambone, F., D'Imporzano,*  
303 *G., Adani, F. (2016).* Short-term experiments in using digestate products as  
304 substitutes for mineral (N) fertilizer: Agronomic performance, odours, and  
305 ammonia emission impacts. *Sci. Total Environ.* 547, 206-214.
- 306 *Rowell, D.L. (1994).* Soil Science: Methods and application. Longman Scientific &  
307 Technical, UK.
- 308 *Ryan, J., Estefan, G., Rashid, A. (2001).* Soil and Plant Analysis Laboratory Manual, 2nd Ed.  
309 International Center for Agricultural Research in Dry Areas, Aleppo, Syria.

- 310 Shaheen, A., Tariq, R., Khaliq, A. (2017). Comparative and Interactive Effects of Organic and  
 311 Inorganic Amendments on Soybean Growth, Yield and Selected Soil properties. *Asian J*  
 312 *Agri & Biol.* 5(2), 60-69.
- 313 Singh, V.P., Singh, R.K. (2000). Rainfed Rice: A Sourcebook of Best Practices and  
 314 Strategies in Eastern India. International Rice Research Institute, p. 292.
- 315 Steel, R.G.D., Torrie, J.H., Dickey, D.A. (1997). Principles and Procedures of Statistics: A  
 316 biometrical approach, 3rd Ed. McGraw Hill Book Co., New York, USA.
- 317 Tonfack, L.B., Bernadac, A., Youmbi, E., Mbouapouognigni, V.P., Nguiguim, M., Akoa, M.  
 318 (2009). Impact of organic and inorganic fertilizers on tomato vigor, yield and  
 319 fruit composition under tropical and soil conditions. *Fruits.* 64, 167-177.
- 320 Wakene, N., Getahun, F., Deressa, A., Dinsa, B. (2007). Integrated Use of Organic and  
 321 Inorganic Fertilizers for Maize Production. Utilization of diversity in land use  
 322 systems: Sustainable and organic approaches to meet human needs. University of  
 323 Rostock, Institute of Land Use, Justus-von-Liebig Weg 6, 18059, Rostock, German,  
 324 pp.1-18.
- 325 Zerzghi, H., Gerba, C.P., Brooks, J.P., Pepper, I.L. (2010). Long-term effects of land  
 326 application of class B biosolids on the soil microbial populations, pathogens, and  
 327 activity. *J. Environ. Qual.* 39, 402-408.
- 328 Zhang, J., Wang, M., Cao, Y., Liang, P., Wu, S., Leung, A.O.W., Christie, P. (2015).  
 329 Replacement of mineral fertilizers with anaerobically digested pig slurry in  
 330 paddy fields: assessment of plant growth and grain quality. *Environ. Sci. Pollut. R.*  
 331 24 (10): 8916-8923.

332

333 **Table 1. Soil physicochemical properties**

Characteristics	Unit	Value
pH	-	8.29

Ece	dSm <sup>-1</sup>	1.83
Organic Matter	%	0.76
Total Nitrogen	%	0.038
Available Phosphorus	mg kg <sup>-1</sup>	8.83
Extractable Potassium	mg kg <sup>-1</sup>	200
Sand	%	52.6
Silt	%	21.4
Clay	%	26
		Sandy clay loam

334

335

336 **Table 2. Chemical composition of organic materials**

Organic Material	Anaerobic Digestate			Farm Manure		
	N	P	K	N	P	K
	-----%-----			-----%-----		
Year 1	0.96	0.63	1.10	0.57	0.42	0.98
Year 2	1.01	0.58	0.92	0.62	0.44	0.86
Year 3	0.94	0.62	0.94	0.58	0.46	0.91

337

338 **Table 3. Effect of integrated use of organic and inorganic fertilizer on wheat grain**  
 339 **yield (Mg ha<sup>-1</sup>)**

Treatments	Year I	Year II	Year II	Pool
Control	1.64 d	1.68 d	1.75 d	1.69 e
RD of NPK	4.30 a	4.42 a	4.28 a	4.33 a
FS on the basis of RD of N	3.17 c	3.11 c	3.30 c	3.19 d
FYM on the basis of RD of N	3.08 c	2.96 c	3.27 c	3.10 d
½ N from FS + ½ N from CF	4.12 ab	4.28 ab	4.04 ab	4.16 ab
½ N from FYM + ½ N from CF	3.87 b	3.96 b	3.61 bc	3.81 c
LSD	0.36	0.39	0.47	0.22

340

341 **Table 4. Effect of integrated use of organic and inorganic fertilizer on wheat straw**  
 342 **yield (Mg ha<sup>-1</sup>)**

Treatments	Year I	Year II	Year II	Pool
Control	3.85 d	3.93 d	3.21 c	3.66 e
RD of NPK	6.22 a	8.12 a	6.68 a	7.01 a
FS on the basis of RD of N	4.86 c	5.65 c	5.23 b	5.25 cd
FYM on the basis of RD of N	4.67 c	4.95 c	5.19 b	4.64 d
½ N from FS + ½ N from CF	5.72 ab	7.63 ab	6.02 a	6.46 ab
½ N from FYM + ½ N from CF	5.52 b	6.89 b	5.67 ab	6.03 bc
LSD	0.63	0.72	0.73	0.84

343

344

345 **Table 5. Post-harvest soil properties**


Treatment	pH	ECe	O.M	P	K
		dSm <sup>-1</sup>	%	-----mg kg <sup>-1</sup> -----	
Control	8.13 <sup>NS</sup>	1.74 <sup>NS</sup>	0.51 e	7.17 d	170 d
RD of NPK	8.18	1.77	0.68 d	11.06 c	196.7 bc
FS on the basis of RD of N	8.19	1.76	0.82 c	12.87 bc	230 ab

FYM on the basis of RD of N	8.19	1.80	0.86 bc	13.63 bc	215 bc
½ N from FS + ½ N from CF	8.19	1.71	0.96 a	14.93 a	245 a
½ N from FYM + ½ N from CF	8.18	1.74	0.93 ab	14.63 bc	220 b
LSD	0.10	0.08	0.07	0.84	19.6


**Table 6. Economic analysis of the study (Benefit cost ratio; BCR)**

Treatments	Total Expenditure(Rs.)	Gross income (Rs.)	Net income (Rs.)	Benefit-cost ratio (BCR)
Control	24595	50700	26105	2.06
RD of NPK	55986	129967	73981	2.32
FS on the basis of RD of N	35595	95800	60205	2.69
FYM on the basis of RD of N	35595	93100	57505	2.62
½ N from FS + ½ N from CF	40290	122367	82077	3.04
½ N from FYM + ½ N from CF	40290	114400	74110	2.84






**Figure 1. Meteorological data showing mean monthly temperature and total rainfall during wheat growing period (2014, 2015, 2016)**



**Figure 2. Three year pool grain and straw yield of wheat in response to digestate and chemical fertilizers application**



**Figure 3. N uptake and use efficiencies as affected by digestate and chemical fertilizer application**