

Symbiotic effectiveness of indigenous rhizobial strains on biological nitrogen fixation of Lablab (*Lablab purpureus*) in the derived savanna of Nigeria

Abstract

Field and pot experiments were conducted in the derived savanna of Nigeria to determine the effectiveness of three indigenous rhizobial strains on the Nitrogen (N) fixation of lablab (*Lablab purpureus*). Soils were collected from two locations, Idi Ayunre and University of Ibadan Teaching and Research Farm (UITRF) for pot experiment in a completely randomised design with a factorial arrangement of $2 \times 2 \times 6$. The treatments were soil type (Idi Ayunre and UITRF soils), sterilisation (partially-sterile and unsterile) and six rhizobial strains inoculation {three indigenous strains, IDC8, OISa-6e and TRC; two exotic strains, IRj 2180A and R25B; and the control (resident native rhizobial strains)}. A field experiment was further conducted at UITRF using five rhizobial inoculations, the three indigenous strains, a combination of R25B and IRj 2180A (R25B+IRj 2180A) and control. Data were collected on biomass dry weight, a number of nodules, nodule dry weight, N derived from the atmosphere (Ndfa), N and P uptake and total N fixed. No significant difference was observed in the Ndfa (%) among the strains. However, indigenous strains IDC8 and OISa-6e performed better than the resident native rhizobia regarding N fixed, dry biomass weight, nodule formation and N uptake. N fixed by lablab inoculated with IDC8 was more than 200% higher than that fixed by resident native bacteria. Lablab N fixation efficiency can be improved using effective indigenous rhizobial strains. Further screening of indigenous strain for N fixation efficiency is recommended.

Keywords: Rhizobium N fixation Forage legume; Lablab

Introduction

The soil improvement potential of lablab (*Lablab purpureus*) has continued to generate research attention in the derived and moist, humid savannah of West Africa where soil fertility has continued to decline due to the pressure caused by increasing population. Lablab is widely cultivated in sub-Saharan Africa, and like other food and forage legumes, it possesses qualities that could prove exceptionally valuable for Africa's rural development and environmental stability (Adebisi and Bosch 2004). These conditions include wide adaptation to different environment, ease of planting and management under subsistence production typical of African agriculture, high yield, drought tolerance and soil improvement potentials (Adebisi and Bosch 2004; Cook et al. 2005; Nworgu and Ajayi 2005, National Research Council 2006; Maass et al. 2010). Sanginga et al. (1996) reported that *Lablab purpureus* had a high N₂-fixing capability and could also adapt to low-P soils in the Guinea savanna of West-Africa.

The high cost and environmental risks associated with the use of chemical fertiliser has made lablab cultivation one of the alternatives for low cost and sustainable N supply in the soil. Lablab can produce about six tonnes of total biomass containing about 140 kg N which can be returned to the soil after decay (Valenzuela and Smith 2002; FAO 2012). McDonald et al. (2001) estimated N₂ – fixed by lablab to be 177 kg N ha⁻¹ which to an extent is substantial for soil fertility improvement. Lablab inclusion into cereal-legume based cropping systems especially maize-based production system as an intercrop or as fallow crop has yielded beneficial results concerning crop yields and soil fertility improvement (MacColl 1990; Tarawali 1991; Devkota and Rerkasem, 2000). Apart from the N-fixation potential, lablab as a cover crop is able to conserve soil and compete with weeds (Schaaffhausen 1963) thereby improving the soil quality for crop production.

50 Nitrogen fixation in all legumes is due to their symbiosis with the soil bacteria
51 Rhizobium. However, the amount of atmospheric N that is fixed is partly hinged on the
52 effectiveness of the rhizobial strains that infect the plant for nodulation and fixation.
53 Biological nitrogen fixation is a natural process, which can be improved by introducing or
54 inoculating legumes with an efficient rhizobia strain for effective nitrogen fixation. FAO
55 (2012) suggested that the seed of lablab should be inoculated with a cowpea-type,
56 Bradyrhizobium strain as it does not easily nodulate with native rhizobia. However,
57 indigenous or native rhizobia have been found to be effective for some legumes like cowpea
58 in some parts of West Africa (N2Africa 2013). Efficient rhizobial inoculants could be
59 obtained by isolating, purifying and screening the native or indigenous rhizobia with standard
60 rhizobia under controlled conditions (Kumar et al. 1997). In Nigeria, Ojo et al (2015) isolated
61 three rhizobial strains from derived savanna soils that are comparable in infectivity and
62 nodulation with some introduced/exotic strains. Thies et al. (1991) reported that the success
63 of any rhizobial inoculation starts with the ability of the inoculant strain to survive and
64 nodulate the host plant.

65 Legume-rhizobial symbiosis is an efficient source of soil nitrogen for sustainable
66 agriculture. Since lablab is among the promising forage legumes that have been identified to
67 improve agricultural sustainability in the moist savannah (Tarawali 1991), this study
68 investigated the effectiveness of three native rhizobial strains on biological N fixation of
69 lablab.

70 **Materials and Methods**

71 **Pot experiment**

72 A pot experiment was conducted at the International Institute of Tropical Agriculture
73 screenhouse, Ibadan, Nigeria using topsoil (0-15 cm) collected from two locations in the
74 rainforest-savanna transition zone of Nigeria, Idi-Ayunre (latitude 7°26'N and longitude

75 3°54'E) and the University of Ibadan Teaching and Research Farm (UITRF, latitude 7°30'N
76 and longitude 3°45'). Idi-Ayunre has soil classified as Nitosol and that of the University of
77 Ibadan Teaching and Research Farm as Alfisol (USDA, 2006). The samples were analysed
78 for pH in water (1:1) (IITA, 1982), soil organic matter using wet dichromate acid oxidation
79 method (Nelson and Sommers, 1982), total nitrogen using Kjeldahl analytical method
80 (Bremner and Mulvaney, 1982), available phosphorus using Bray-1 method (Bray and Kurtz,
81 1945), particle size using Bouyoucus hydrometer method (Okalebo et al., 1993),
82 exchangeable Mg, Ca, K and Na extracted using neutral 1M ammonium acetate and
83 determined with spectrophotometer (Okalebo et al., 1993).

84 Soil samples for each location were divided into two parts, one part was sterilized using a
85 direct flaming method with the aid of a Terraforce sterilizing machine (IITA fabricated
86 sterilizing machine). Two kilograms soil was weighed into each pot for the planting

87 **Rhizobial population count**

88 The rhizobial population count of the two locations was determined using the Most
89 Probable Number (MPN) as outlined by Somasegaran and Hoben (1994). Two soyabean
90 varieties, TGx1448-2E and TGx 1456-2E, and one cowpea variety IT89KD-288 seeds were
91 sterilised, pre-germinated and transplanted into sterilised growth pouches containing
92 modified Jensen's N-free nutrient solution (Roughley, 1984). A 5-fold dilution series (5^{-1} – 5^{-6})
93 of the soils of the location with four replicates was used to inoculate each plant in the
94 growth pouch one week after planting. The diluent contained 0.250 g K_2HPO_4 and 0.10 g Mg
95 $SO_4 \cdot 7H_2O$ dissolved in 1 L distilled water. Nodule formation on plants was observed and
96 recorded for thirty days. The presence (+) or absence (-) of nodules on each plant were scored
97 and MPN values were calculated with MPN table.

98 Experimental design and data collection

The experimental design was a $2 \times 2 \times 6$ factorial arrangements in a completely randomised design with three replicates. The treatments were location (Idi-Ayunre and UITRF), soil sterilization at two levels (sterile and unsterile) and rhizobial strains at six levels (control, Isolate 1 – OISa-6e, Isolate 2 – IDC8, Isolate 3 – TRC2, IITA strain 1 – R25B, IITA strain 2 – IRj 2180A). Two seeds of lablab were planted per pot and inoculation was done one week after planting using dispenser to directly inoculate the soil just below the seedling with 2 ml broth culture of the rhizobial strains. Data were collected at harvest to determine the shoot dry weight, number of nodules per pot, nodule dry weight per pot and percentage nitrogen derived from atmosphere [Ndfa (%)]. N_2 fixed was determined by taking samples from stem + petioles for tissue extraction (Hot water extract) for ureide – N and NO_3^- –N in the laboratory. The procedure was followed as outlined by Herridge (1982).

Description of rhizobial strain

The five strains that were used for the experiment were IDC8, TRC, OISa-6e, IRj 2180A and R25B. The IDC8, TRC and OISa-6e were indigenous strains isolated from cowpea planted on Idi-Ayunre and UITRF soils and soybean planted at Orile Ilugun soil (Ojo et al 2015). The IRj 2180A (soybean isolate) and R25B (promiscuous isolate) were rhizobial collections from International Institute of Tropical Agriculture, Ibadan, Nigeria (Sanginga et al 2000).

Field experiment

Field experiment was set up at UITRF in a randomized complete block design with three replicates. The rhizobial strains IDC8, TRC2, OISa-6e, R25B+IRj 2180A and the control were used as treatments. R25B and IRj 2180A were exotic strains used separately in the greenhouse experiment but mixed together for the field experiment.

One kilogramme of seeds of lablab was inoculated with 10 g peat culture of each rhizobial strain using the method of Somasegaran and Hoben (1994). The Yeast Mannitol Broth (YMB) cultures of each strain were aseptically injected into different peat package carriers at

ratio 1: 1 (ml/wt in g). Inoculated peats were incubated for 2 weeks at 28°C to gain excess of $10^8 - 10^9$ cells g^{-1} . Planting was done between July and early August using a late maturing variety of lablab NAFR14. The seeds were planted at a spacing of 75 cm × 25 cm.

Data collection

Plants were uprooted systematically at three points per plot using a 30 cm × 30 cm quadrant. Root of five plants and soil under the quadrant area were removed to a depth of 15 cm to determine number of nodules. The nodules on the root and those detached in the soil were counted and weighed to get the fresh nodule weight and oven-dried at 78°C to a constant dry weight. The root and the shoot were separated, air-dried for 72 hours before oven dried at 78°C to constant dry weights. The stem + petioles of the five sampled plants were and ground. Ureide – N and $NO_3 - N$ were determined from the stem + petioles samples using tissue extraction (hot water extract) as described by Herridge (1982). Shoot biomass N and P contents were also analysed using the method of Herridge (1982). Harvesting was done at physiological maturity of each variety.

Statistical analyses: Data obtained were subjected to analysis of variance using PROC GLM of SAS (SAS 2003). Data for number of nodule were transformed using square root transformation. Means were separated at a significant level $P = 0.05$ using Least Significant Differences (LSD).

Results

Physical, chemical and biological properties of the experimental soils

The Idi Ayunre and UITRF soils were slightly acidic with UITRF being more acidic. The total N, available P, Ca, Mg and Na were higher in Idi-Ayunre soil compared to UITRF soil. In fact, total N and available P were more than two-fold higher in Idi-ayunre soil compared to those of UITRF soil (Table 1). The soil physical properties of Idi-Ayunre soil revealed a

sandy loam soil while that of UITRF revealed a loamy sand soil. The number of rhizobial cell in the UITRF soil was significantly higher ($P=0.05$) than that of Idi Ayunre (Fig. 1). The number of rhizobial cell ($14.2 \times 10^6 \text{ g}^{-1} \text{ soil}$) in the UITRF was about two-fold higher than that of Idi Ayunre ($7.3 \times 10^6 \text{ g}^{-1} \text{ soil}$).

Growth and nodulation of lablab in pot experiment

The soils used for the pot experiment significantly ($P=0.05$) affected biomass dry weight, number of nodules, nodule dry weight and Ndfa (%). The biomass, nodule dry weights and Ndfa (%) were significantly higher ($P=0.05$) in Idi Ayunre soil than UITRF soil (Table 2). However, the numbers of nodules were higher in UITRF soil than Idi Ayunre soil. The effect of soil sterilization was significant on nodulation as partially sterile soil had significantly lower ($P=0.05$) number of nodules and nodule dry weight compared to unsterile soil. However, biomass dry weight was significantly higher ($P=0.05$) in sterile soil than unsterile soil. The rhizobial strains used for the inoculation of lablab in the pots significantly affected nodulation of the plants. The numbers of nodules formed by lablab in the pot were significantly higher in inoculated plants than the control (Table 2). Similarly the nodule dry weight was higher in inoculated plants than the uninoculated (control). The indigenous strain IDC8 isolated from Idi Ayunre soil had the highest number of nodules followed by the exotic strain R25. The effect of the rhizobial strains on nodulation did not reflect in the Ndfa (%) as there was no significant difference between inoculated plants and uninoculated plants (Table 2).

The effect of interaction of the soil, sterilization and inoculants on nodulation revealed that numbers of nodules of lablab in UITRF unsterile soil was significantly higher than the number of nodules in the sterile soils and unsterile of Idi-Ayunre (Table 3). In both sterile and unsterile soil of UITRF, the number of nodules in inoculated plants was significantly higher than the control plants except the plants inoculated with IRJ 2180A and

OISa-6e in sterile soil. The number of nodules of plants under unsterilized soil inoculated with R25 and TRC2 in Idi-Ayunre and IDC8 and IRJ 2180A in UITRF were significantly higher than their corresponding treatment in sterile soil. In sterile soil at Idi-ayunre and UITRF, IDC8 inoculated plants had significantly higher number of nodules than the uninoculated plants. The nodule dry weight of the unsterile soil in Idi-Ayunre was significantly higher than that of sterile soils in the two soil type and that of the unsterile soil in UITRF (Table 3). Inoculated plants with bacterial strains OISa-6e, R25 and TRC2 in Idi Ayunre had significantly higher nodule dry weight than the control plant. In UITRF, only IDC8 had higher nodule dry weight than the control treatment. In spite of significantly higher number of nodules in UITRF when inoculated plants in sterile were compared to the control, the nodule dry weights were not significantly difference (Table 3).

Growth, nutrient uptake and N fixation of lablab at UITRF (field experiment), Ibadan

The biomass dry weights of lablab in the plots inoculated with the strains IDC8 and R25 were significantly higher than those of the strain TRC3 and the control. In fact, there was more than two-fold increase in the biomass dry weight of lablab inoculated with IDC8 than that of the control (Fig 2). The number of nodule in lablab inoculated with indigenous rhizobial OISA-6E and IDC8 were significantly higher than the other treatments (Table 4). The control plant had the lowest number of nodules but was not significantly lower than plants inoculated with TRC2. There was no significant difference in the Ndfa (%) when the rhizobial strains were compared even with the control. However, the total N fixed (kg/ha) in the IDC8 inoculated plants was significantly higher than other strains except R25B+ IRj2180A inoculated plants. The control treatment had significantly lower total N fixed when compared with other strains except TRC2. The N fixed (kg/ha) in IDC8 and R25B plots was about 144% and 93% higher than control plot respectively (Table 4). The N and P contents of plant and P uptake were not significantly affected by the inoculation of rhizobial strains. However,

the N uptake was significantly increased in IDC8 inoculated plants compared to the control and TRC2 inoculated plants (Table 5) .

Discussion

The biological analyses of the bacteria cells in the two soils revealed a very low population of rhizobial cells with UITRF having higher population which almost double that of Idi Ayunre. The rhizobial count observed was within the range of 2×10^0 to 3.2×10^3 reported by Abaidoo et al. (2007) for twelve sites in Nigeria. Soils that have never been inoculated or cultivated with legumes were selected for the study and this probably contributed to the low rhizobial cell count. Slattery and Pearce (2002) also observed a range varying from less than 10 to 10^6 in Australian soils and reported that considerable variation exist between bacteria population in soils and could be due to several factors which include field history, location of sampling, soil characteristics and the presence of a host plant. The two locations had different soil characteristics and field history. Idi Ayunre soil was under bush fallow of over ten years unlike UITRF soil which was collected from a research farm cultivated with crops others than legumes. The detection of rhizobia in the fallow field was consistent with other previous work by Abaidoo et al. (2007). Higher fertility status of Idi Ayunre soil compared to UITRF soil can also be linked to the fallow and cropping history of the two locations and it reflected in the higher biomass dry weight of lablab in Idi Ayunre soil compared to UITRF soil in the pot experiment.

The inoculation of lablab by the indigenous isolated strains yielded more nodules and N fixation compared to where plants were not inoculated. In the pot experiment, it was clearly shown that the introduced indigenous and exotic strains increased nodulation of lablab more than the native indigenous rhizobia. This shows that inoculant strains were highly compatible and were able to establish a functional symbiotic relationship without much interference of soil native rhizobia or any soil microbes. In other words, the inoculated strains

were able to demonstrate their nodulation potential and out-competed the soil native rhizobia for nodule occupancy. The competitive ability of these introduced strains was also displayed in the field experiment at UITRF where introduced indigenous strains like IDC8 and OISA-6E were able to increase the number of nodules by two fold compared to the indigenous rhizobia in the control treatment. The competitive ability of the introduced strains was probably enhanced by the low rhizobial cell in the two soils. It has been reported that the introduced rhizobia are able to perform better than the native one when the population is $<10^6$ cell g⁻¹ soil (Brockwell et al 1995; Slattery and Pearce, 2002).

The nodulation of lablab by the three introduced indigenous strains demonstrated a broad specificity of the three strains for certain crops. These strains were either isolated from soybean or cowpea but were able to cross-inoculate with the two crops (Ojo et al 2015) and also with lablab in this study. Lablab, soybean and cowpea belong to the genera *Phaseolae* (Andrew and Andrew 2017) and all the three have been found to form nodules with *bradyrhizobium* (Andrew and Andrew 2017; Steenkamp et al. 2008). *Bradyrhizobium lablabi* that nodulate lablab were also isolated in peanut *Arachis hypogaea* (Chang 2011). Symbiosis specificity is complex, involving fine-tuned signal communication between the symbiotic partners, which can occur at multiple phases of the interaction, ranging from initial bacterial attachment and infection to late nodule development associated with nitrogen fixation (Wang et al 2012). The rhizobium species of the three isolates were not determined in this study but it can be concluded that the three isolates had specificity the three crops.

The N derived from the atmosphere by the plant with and without the introduced rhizobia on the field was between 51-56 % which was within the range reported for other legumes (Abaidoo et al 2017). Although there was no significance difference in the Ndfa (%) through inoculation, the N fixed in the soil was significantly influenced by rhizobial strains. The introduced indigenous rhizobial strains IDC8 and OISA-6E performed better than the

resident indigenous rhizobia. This is an indication that the use of this isolated indigenous bacteria for inoculation of lablab in the derived savanna of Nigeria could provide more N for the soil especially where the residue are returned back to the soil. Biomass dry weight and N uptake was greatly improved by IDC8 compared to the resident native rhizobia. Yield increase were also observed in some legumes inoculated with isolated strains of indigenous rhizobia even in soil with high population of resident indigenous rhizobia (Mostasso et al., 2002, Mrabet et al, 2005, Mulas et al., 2011). The isolated indigenous strain IDC8 performed better than the combination of IRj 2180A and R25B, the exotic strains, in terms of biomass dry weight, nodule formation and N uptake. Of the three indigenous that were tested, it is only IDC8 that was better than the exotic strains. This is an indication that some indigenous rhizobial strains may compete favourably or out-performed some of the exotic strains that are in use for farmers.

Conclusion

The **introduced** indigenous rhizobial strains could effectively fixed N in lablab in the derived savanna of Nigeria. The extent of the performance of the indigenous rhizobia can be influenced by the strain type coupled with other plant and soil factors. Screening for the effectiveness of more indigenous rhizobia can provide more effective strains than the exotic ones. With better biological N fixation by lablab using effective rhizobial strains such as IDC8, the crop can serve a dual purpose of enriching the soil and providing forage for the farmers.

References

Abaidoo RC, Keyser HH, Singleton PW, Dashiell KE, Sanginga N. Population size, distribution, and symbiotic characteristics of indigenous Bradyrhizobium spp. that nodulate TGx soybean genotypes in Africa. *Applied Soil Ecology*. 2007; 35: 57–67.

273 Abaidoo RC, Dare MO, Killani S, Opoku A. Evaluation of early maturing cowpea (*Vigna*
274 *unguiculata*) germplasm for variation in phosphorus use efficiency and biological
275 nitrogen fixation potential with indigenous rhizobial populations. *J. Agric. Sci*
276 *(Camb)* 2017; 155: 102 – 116.

277 Adebisi AA, Bosch CH. *Lablab purpureus* (L.) Sweet. In: Grubben GJH, Denton OA, editors.
278 Plant Resources of Tropical Africa (PROTA), no. 2, vegetables. PROTA Foundation,
279 Wageningen, The Netherlands / Backhuys, Leiden, The Netherlands / CTA,
280 Wageningen, The Netherlands, pp 343–348. 2004.

281 Andrews M, Andrews ME. Specificity in legume-rhizobium symbioses. *Int. J of Mol. Sci.*
282 2017; 18: 705.

283 Bray RH, Kurtz LT. Determination of total organic and available form of phosphorus in soils.
284 *Soil Science Journal*. 1945; 59: 39-45.

285 Bremmer JM, Mulvaney CS. Nitrogen-Total. *Methods of soil analysis II*. A.L. Page., R. H.
286 Miller and D. R. Keeney. Eds. *Agronomy Journal*. 1982; 9: 595-624.

287 Brockwell J, Botteonley PJ, Thies JE. Manipulation of rhizobia microflora for improving
288 legume productivity and soil fertility: a critical assessment. *Plant and Soil* 174: 1995;
289 143 – 150.

290 Chang YL, Wang JY, Wang ET, Liu HC, Sui XH, Chen WX. *Bradyrhizobium lablabi* sp.
291 nov., isolated from effective nodules of *Lablab purpureus* and *Arachis hypogaea*. *Int.*
292 *J. Syst. Evol. Microbiol.* 2011; 61: 2496–2502.

293 Cook BG, Pengelly BC, Brown SD, Donnelly JL, Eagles DA, FrancoMA, Hanson J, Mullen
294 BF, Partridge IJ, Peters M, Schultze-Kraft R. Tropical forages: an interactive selection
295 tool [CD-ROM], CSIRO, DPI&F(Qld), CIAT and ILRI, Brisbane, Australia. 2005.
296 Accessed Feb 15 2016 Available: <http://www.tropical-forages.org>

297 Eaglesham, RJ. Comparison of nodulation promiscuity of US and Asian-type soyabean.
 298 Tropical Agriculture. (Trinidad) 1985; 62: 105 – 109.

299 FAO. Grassland species index. Lablab purpureus. 2012. Accessed June 6 2012. Available:
 300 <http://www.fao.org/ag/AGP/AGPC/doc/Gbase/DATA/Pf000047.HTM>

301 Herridge, DF. Legume Nitrogen fixation. In Methods to measure legume nitrogen fixation.
 302 Workshop Handbook 1994.NWFP. Agriculture University Peshawar, Pakistan. 1982.

303 IITA. International Institute of Tropical Agriculture, Manual series 1982; 7:15-17.

304 Kumar TKD, Rao ARM, Reddy SM, Srivastava HP. Ram-Reddy S. Effect of host plant on
 305 rhizobial populations. Microbial Biotechnology (“Prof. K.S. Bilgramm.
 306 Commemoration) 1997; 11: 79 – 82.

307 Maass BL, Knox MR, Venkatesha SC, Angessa TT, Ramme S, Pengelly BC. *Lablab*
 308 *purpureus*-a crop lost for Africa? Trop. Plant Biol. 2010; 3(3):123–135.

309 MacColl D. Studies in maize (*Zea mays*) at Bunda, Malawi. III. Yield in rotation with pasture
 310 legumes. Experimental Agriculture. 1990; 26: 263–271.

311 McDonald LM, Wright P, MacLeod DA, Nitrogen fixation by lablab (*Lablab purpureus*) and
 312 lucerne (*Medicago sativa*) rotation crops in an irrigated cotton farming system.
 313 Australian Journal of Experimental Agriculture. 2001; 41(2): 219–225.

314 N2Africa. Establishment of rhizobial inoculant production factories in Africa (Internet).
 315 Wageningen UR. 2013. Accessed Jan 3 2014. Available:
 316 [http://www.n2africa.org/content/establishment-rhizobial-inoculant-production-](http://www.n2africa.org/content/establishment-rhizobial-inoculant-production-factories-africa/)
 317 [factories-africa/](http://www.n2africa.org/content/establishment-rhizobial-inoculant-production-factories-africa/)

318 National Research Council Lost Crops of Africa: Volume II: Vegetables. Washington, DC:
 319 The National Academies Press,. 2006. doi:10.17226/11763

320 Nelson DW, Sommers LE. Total Carbon, Organic matter. In: Page AL, editor. Methods of
 321 soil analysis Part 2. Agronomy Monograph 9 2nd ed. Madison, Wisconsin: American
 322 Society of Agronomy and Soil Science Society of America; 1982. 539-579.

323 Nworgu FC, Ajayi FT. Biomass, dry matter yield, proximate and mineral composition of
 324 forage legumes grown as early dry season feeds. Livestock Research for Rural
 325 Development 2005. 17(11) Accessed Nov 14 2014. Available:
 326 <http://cipav.org.co/lrrd/lrrd17/11/nwor17121.htm>

327 Odunze AC, Tarawali SA, de Haan NC, Akouegnon GE, Amadji AF, Schultze-Kraft R,
 328 Bawa GS. Forage legumes for soil productivity enhancement and quality fodder
 329 production. J Food Agric Environ. 2004; 2(2):201–209.

330 Okalebo JR, Gathua KW, Woomer PL. Laboratory methods of soil and plant analysis: A
 331 working manual. Tropical Soil Biology and Fertility Programme, UNESCO. Nairobi,
 332 Kenya: EPZ Publishers. 1993.

333 Roughley RJ, Gemell LG, Thompson JA, Brockwell J. The number of Bradyrhizobium
 334 species (Lupinus) applied to seed and its effect on rhizosphere, colonization,
 335 nodulation and yield of Lupin. Soil Biology and Biochemistry. 1993; 25: 1453 –
 336 1458.

337 Sanginga, N. Role of biological nitrogen fixation in legume based cropping system; a case
 338 study of West Africa farming systems. Plant and Soil. 2003; 252: 25 – 39.

339 Sanginga N, Thottappilly G, Dashiell K. Effectiveness of rhizobia nodulating recent
 340 promiscuous soyabean selection in the moist savanna of Nigeria. Soil Biology and
 341 Biochemistry 2000; 32(1): 127 – 133.

342 Sanginga N, Okogun JA, Akobundu IO, Kang BT. Phosphorus requirement and nodulation of
 343 herbaceous and shrub legumes in low P soils of a Guinea savannah in Nigeria.
 344 Applied Soil Ecology. 1996; 3: 247-255.

345 SAS. SAS/STAT.Guide for personal computers, Version 9.1 Edition,1028 pp Cary, NC,SAS.
 346 Institute Inc. 2003.

347 Schaaffhausen RV. Dolichos lablab or hyacinth bean: its uses for feed, food and soil
 348 improvement. Econ Bot. 1963; 17:146–153.

349 Sheahan CM. Plant guide for lablab (*Lablab purpureus*). USDA-Natural Resources
 350 Conservation Service, Cape May Plant Materials Center. Cape May, NJ. 2012.

351 Slattery JF, Pearce DJ. Effects of resident rhizobia communities and soil type on the effective
 352 nodulation of pulse legumes. Soil Biology and Biochemistry 2004; 36: 1339 – 1346.

353 Somasegaran P, Hohen HJ. Handbook for Rhizobia: Method in Legume-Rhizobium
 354 Technology. University of Hawaii NifTAL Project 1000 Holomua Road Paia, HI
 355 USA. 1994.

356 Steenkamp ET, Stepkowski T, Przymusiak A, Botha WJ, Law IJ. Cowpea and peanut in
 357 southern Africa are nodulated by diverse Bradyrhizobium strains harboring
 358 nodulation genes that belong to the large pantropical clade common in Africa. Mol.
 359 Phylogenet. Evol. 2008; 48: 1131–1144.

360 USDA. A basic system of soil classification for making and interpreting soil surveys.
 361 Agriculture Handbook 436, Natural Resources Conservation Service, 2006. USDA
 362 Accessed Nov. 14 2012. Available:
 363 www.intersciencewiley.com/journal/18981858/abstract.

364 Tarawali G. Residual effect of Stylosanthes fodder banks on grain yield of maize. Tropical
 365 Grasslands. 1991; 25: 26–31.

366 Thies JE, Singleton PW, Bohlool SB. Influence of the size of indigenous rhizobia population
 367 on establishment and symbiotic performance of introduced rhizobia on field legumes.
 368 Applied and Environmental Microbiology 1991; 57(1): 19 – 28.

Wang D, Yang S, Tang F, Zhu H. Symbiotic specificity in the legume: rhizobium mutualism. Cell Microbiol. 2012; 14(3): 334-342.

Yousef AN, Al-Nassir AS, Al-Azawi SK, Abdull-Hussain N. Abundance of peanut rhizobia as affected by environmental conditions in Iraq. Soil Biol. Biochem. 1987; 19: 391–396.

Zaharan IH. Rhizobium-legume symbiosis and nitrogen fixation under severe conditions and in an arid climate. Microbiology and molecular Biology Reviews 1999; 63: 968 – 989.

Table 1. Soil physical and chemical properties of the experimental locations

Soil properties	Idi Ayunre	UITRF
pH (KCl)	6.55	5.76
Total N (g kg ⁻¹)	0.23	0.08
Available P (mgkg ⁻¹)	0.42	0.13
Ca (cmolkg ⁻¹)	7.87	4.84
Mg (cmol kg ⁻¹)	2.59	1.65
K (cmolkg ⁻¹)	0.83	0.85
Na (cmol kg)	0.44	0.43
Fe (mg/kg ⁻¹)	25.34	26.29
Mn (mg/ kg ⁻¹)	14.74	12.38
Sand (gkg ⁻¹)	645.0	812.5
Clay (gkg ⁻¹)	185.0	100.0
Silt (gkg ⁻¹)	170.0	87.5
Textural class	Sandy loam	Loamy sand




Figure 1: Rhizobial population in the studied locations (Error bar represents LSD at P=0.05)

Table 2. Biomass dry weight, nodulation and N derived from atmosphere of lablab as affected by soil type, sterilization and rhizobial inoculation in pot experiment

Treatment		Biomass	Number	Nodule	
		dry weight	of	dry weight	NDFA
		(g/pot)	nodules	(g/pot)	(%)
Soil type					
	Idi Ayunre	12.95a	27.28b	2.13a	59.75a
	UITRF	8.77b	40.28a	1.60b	43.57b

	<i>P value</i>	0.0001	0.0001	0.004	0.003
Sterilization					
	Partially sterile	11.36a	28.69b	1.20b	50.48
	Unsterile	10.36b	38.86a	2.53a	52.84
	<i>P value</i>	0.05	0.0001	0.0001	0.63
Rhizobial Inoculation					
	OISa-6e	9.88	30.58b	2.08a	49.96
	IDC8	11.78	41.33a	1.97a	52.29
	IRJ 2180A	11.37	37.25ab	1.98a	53.16
	R25	10.72	39.83a	2.02a	54.40
	TRC2	10.03	36.25ab	2.07a	51.39
	Control	11.39	17.42c	1.09b	48.76
	<i>P value</i>	0.16	0.0001	0.04	0.73

Values followed by the same alphabet are not significantly different (P=0.05)

Table 3. Interactive effect of soil type, sterilization and rhizobial inoculation on biomass dry weight and nodulation in pot experiment

Soil type	Rhizobial inoculation	Biomass dry weight (g/pot)		Number of nodule		Nodule dry weight g/pot	
		Sterile	Unsterile	Sterile	Unsterile	Sterile	Unsterile
Idi Ayu nre	OISa-6e	10.58	11.78	20.33	18.00	0.89	4.08
	IDC8	15.20	12.99	36.67	28.33	1.65	2.43
	IRj 2180A	15.71	11.94	41.00	29.33	1.89	2.55
	R25	11.22	12.99	12.33	56.00	0.66	4.10
	TRC2	12.73	11.06	16.67	37.00	1.47	3.55
	Control	17.90	11.37	10.00	21.67	0.40	1.94
UITRF	OISa-6e	9.06	8.10	38.67	45.33	1.36	1.98
	IDC8	8.61	10.34	40.00	60.33	1.08	2.71
	IRj 2180A	9.04	8.80	28.67	50.00	1.18	2.32

R25	10.26	8.40	45.33	45.67	1.57	1.74
TRC2	7.99	8.34	42.67	48.67	1.42	1.86
Control	8.06	8.24	12.00	26.00	0.86	1.15
<i>SE</i>	4.57		8.92		0.34	

393

394 Table 4. Nodulation and N fixation as affected by rhizobial inoculation at UITRF field
395 experiment, Ibadan

Rhizobial inoculation	Number of		N fixed (kg/ha)
	nodule	Ndfa (%)	
OISa-6e	76.0a	54.37	83.6bc
IDC8	76.0a	56.16	119.0a
R25B+IRj2180A	54.7b	56.0	94.3ab
TRC2	50.7bc	51.24	58.8cd
Control	34.0c	55.7	48.7d
<i>P value</i>	0.001	0.36	0.001

396 Means within column followed by the same alphabet are not significantly different (P=0.05)

397 Table 5. Nitrogen and P uptake and content as affected by rhizobial inoculation at UITRF
398 field experiment

Rhizobial inoculation	N content (% dm ¹)	P content (% dm)	N uptake (kg/ha)	P uptake (kg/ha)
OISa-6e	3.12	0.37	153.1ab	18.5
IDC8	3.09	0.38	216.0a	26.1
R25B	2.53	0.28	159.9ab	17.6
TRC2	2.88	0.42	110.3b	17.3
Control	3.02	0.29	104.0b	12.9

<i>P value</i>	0.14	0.09	0.05	0.18
----------------	------	------	------	------

Means within columns followed by the same alphabet are not significantly different (P=0.05);
¹ dry matter;




Figure 2: Biomass dry weights of lablab treated with different strains of rhizobium (Error bar represents LSD at P=0.05).