

Integrated Nitrogen Management on Nutrient Contents, Uptake and Use Efficiency of BRRI Dhan 29

ABSTRACT

A field experiment was performed with BRRI Dhan 29 at Field Laboratory of Soil Science, Hajee Mohammad Danesh Science and Technology University, Dinajpur, Bangladesh. The effects of organic (cowdung) and inorganic (urea) amended N fertilizers were evaluated for NPKS contents and uptakes in grain and straw, and also for observing N use efficiency. Randomized complete block design was set for seven treatments based on recommended dose of N (RDN) @ 150 kg ha⁻¹ using cowdung and/or urea alone or their combinations. The NPK contents as well as their uptake in grain and straw were significantly affected due to different treatments while S content was insignificant. The application of recommended doses of N from urea (T₁) showed highest N content in grain (1.11%) and straw (0.71%) which was closely followed by the treatment T₆ (20% RDN from cowdung + 80% RDN from urea). The maximum P content was found from the grain and straw of T₆ treatment whereas a significant increase in the P, K and S contents were noted due to combined application of N from cowdung and urea. The NPKS uptake of grain and straw as well as total uptake (107.60 kg ha⁻¹ N, 27.84 kg ha⁻¹ P, 71.36 kg ha⁻¹ K and 25.63 kg ha⁻¹ S) were recorded maximum in T₁, followed by T₆. The treatment T₁ yielded maximum apparent N recovery efficiency (45.06%) and agronomic N use efficiency (19.60 kg kg⁻¹) while the maximum physiological N use efficiency (43.55 kg kg⁻¹) was found in T₆. Results also suggested that the application of lower doses of urea N with higher doses of cowdung N were not useful for the N recovery due to low N supplying potentiality of manures in a single cropping season. However, the incremental rates of urea N upto 80% along with 20% cowdung N effectively increased the recovery of N in all the parameters of N use efficiency and should be applicable to optimize the need for N requirement and build up a good soil health.

Keywords: Cowdung, urea, NPKS, uptake, nitrogen use efficiency, rice etc.

1. INTRODUCTION

Rice is the principal carbohydrate supplying food crop for the people of Bangladesh. The country is the fourth-largest rice producer in terms of area and production among the rice growing countries [1]. The agriculture in Bangladesh is mainly dominated by intensive rice cultivation in favor of its geographical and agroclimatic conditions but the soils experience multiple nutrient deficiencies over the years. In intensive cropping system, continuous use of high levels of chemical fertilizers usually with high N inputs lead to nutritional imbalance in soil, decline crop productivity and reduced N use efficiency as well as increased N loss to the environment [2]. The farmers of Bangladesh use only about 172 kg nutrients ha⁻¹ annually (132 kg N, 17 kg P₂O₅, 17 kg K₂O, 4 kg S, 2 kg Zn + B), as against the crop removal of about 250 kg ha⁻¹ [3] and they are mainly concerned about the widely used urea N fertilizer for rice cultivation. A recent estimate also showed that rice (HYV) uptake about 108 kg N, 18 kg P, 102 kg K and 11 kg S ha⁻¹ from soils [4]. Considering the ecological and environment concerns over the increased and indiscriminate uses of inorganic fertilizers have continued to stimulate research on uses of organic materials as sources of nutrients [5]. Use of organic matter as a source of plant nutrients increases the fertilizers use efficiency and makes soil living. Nitrogen is characteristically the nutrient of most concern because of its enormous impact on

cereal crop yields including rice. Manure is one of the most important N sources in paddy rice systems although it poses a countless challenge in meeting rice N requirement. Generally, manure amended rice systems historically **has** been used for its N use efficiency improvement and animal waste recycling. Nitrogen release from manure is relatively slow compared to chemical N fertilizer like urea, and may mismatch the N requirement for rice growth, especially during the mid or late rice growth period. Many agroecologists have focused that manure should be applied as basal fertilizer and combined with urea or other fast released chemical N fertilizers as topdressing [6,7]. The split application of N might provide a compromise between traditional and modern production systems which would improve both soil micro environment and N use efficiency [8]. So, selection of adequate amounts of N from organic and inorganic sources is one of the best solutions for sustainable rice cultivation. Cowdung is a potential source of organic manure in Bangladesh and extensively used in the vegetables cultivation. The application of cowdung in rice fields as a nutrient source may reduce the requirement of chemical nitrogenous fertilizers, but the question has not been examined sufficiently for a wide array of soil and variety. It is also important to look beyond the immediate crop needs of highly demanding N nutrition during growth in order to optimize the uptake and use efficiency from organic and inorganic sources. Therefore, the present study was carried out to evaluate the effects of nitrogen either from urea or cowdung along with their combinations on the changes of nutrient contents, uptake and use efficiency by BRRI Dhan 29.

2. MATERIAL AND METHODS

2.1 Experimental site and soil

The experiment was set up at the Soil Science Field Laboratory, Hajee Mohammad Danesh Science and Technology University (HSTU), Dinajpur during the boro season (January to May 2015). The site was under the Old Himalayan Piedmont Plain (AEZ 1) and the soils belong to Ranisankail series having sandy loam texture. The soil was characterized by slightly acidic in nature (pH = 6.56) with OC (0.34%), total N (0.02%), available P (11.24 ppm), exchangeable K (0.13 me 100⁻¹ g), exchangeable Ca (1.26 me 100⁻¹ g), exchangeable Mg (0.51 me 100⁻¹ g), available S (35.47 ppm), available Zn (0.74 ppm), available B (0.23 ppm), and CEC (5.5 me 100⁻¹ g).

2.2 Treatments and design

There were seven treatments consisting different combinations of recommended dose of N (N₁₅₀) either from organic and/or inorganic sources (cowdung and urea used as organic and inorganic sources, respectively). The treatment combinations were T₀: Control (no nitrogen), T₁: 100 % RDN from urea, T₂: 100% RDN from cowdung, T₃: 80% RDN from cowdung + 20% RDN from urea, T₄: 60% RDN from cowdung + 40% RDN from urea, T₅: 40% RDN from cowdung + 60% RDN from urea, and T₆: 20% RDN from cowdung + 80% RDN from urea. All the treatments also received recommended doses of other nutrients on soil test basis (P₂₀K₆₅S₁₈Zn_{1.3}) through TSP, MOP, gypsum and ZnSO₄, respectively. The treatment wise required nitrogen from cowdung was calculated on the basis of 0.78% N content of well decomposed dried cowdung. The experiment was laid out in the randomized complete block design (RCBD) with four replications following the net plot size of 15 sq. m (5m x 3m). The full doses of cowdung as per treatments were added 15 days before transplanting. The full doses of TSP, MP, Gypsum, Zinc sulphate were applied during the final land preparation. Urea was applied in three equal splits: the first split after 7 days of transplanting, the second split as top dressing after 30 days of transplanting while third one after 60 days of transplanting (before panicle initiation stage). The standard procedure was followed for transplanted as well as other intercultural operations [9]. After recording the yield, grains and straw samples from each unit plot were collected for analysis of nutrient contents.

2.4 Analyses of nutrient contents in plant samples

2.4.1 Preparation of plant samples

Both the grain and straw samples were dried in an oven at 60°C for 24 hours and then grounded by a grinding mill. The prepared samples were kept in desiccators until analyses.

2.4.2 Determination of total nitrogen

Well ground 0.1 g oven dry samples were taken in kjeldahl flasks containing 1.1 g catalyst mixture (K_2SO_4 : $CuSO_4 \cdot 5H_2O$: $Se = 100: 10: 1$), 3 ml 30% H_2O_2 and 5 ml conc. H_2SO_4 . The flasks were swirled and allowed to stand for 10 minutes and heated at 380 °C until the digest became clear and colourless. After cooling, the content was diluted with water and made the volume of 100 ml. Then, 40% NaOH was added with the digests for distillation and the evolved ammonia was trapped in 4% H_3BO_3 solution having 5 drops of the mixed indicator [bromocressol green ($C_{21}H_{14}O_5Br_4S$) and methyl red ($C_{10}H_{10}N_3O_3$) solution]. Finally, the distillates were titrated with the standard 0.01 N H_2SO_4 until the colour changed from green to pink [10]. A reagent blank was also prepared in the same way for accuracy in analysis.

2.4.3 Determination of P, K and S

Plant samples (0.5 g) were digested by using 10 ml of diacid mixture (HNO_3 : $HClO_4 = 2: 1$) into 100 ml kjeldahl flasks. After leaving for overnight, the flasks were heated slowly upto 200 °C until the contents became sufficiently clear and colourless. After cooling, the digests were diluted by distilled water to make 50 ml in a volumetric flask. The digests were then filtrated and used for P, K and S determination. 1 ml digest from grain samples and 2 ml digests from straw samples were used for P determination followed by developing blue colour of phosphomolybdate complex using $SnCl_2$. The absorbance was measured at 660 nm wave length in spectrophotometer and available P was calculated with the help of a standard curve [11]. In case of K determination, 5 ml digest for grain and 2 ml for straw were taken and diluted to 50 ml volume for getting the desired concentration because of the absorbance of samples could be measured within the range of standard solutions. The absorbances were finally measured by flamephotometer. The content of S in the digest was determined by adding acid solution followed by forming turbidity using $BaCl_2$. The intensity was measured by spectrophotometer at 420 nm wave length [12].

2.5 Nutrient uptake

Nutrient uptake was calculated from the measured yield and nutrient contents (grain and straw), and expressed by the formula stated below:

$$\text{Nutrient uptake (kg ha}^{-1}\text{)} = \frac{\text{Yield (kg ha}^{-1}\text{)} \times \text{Nutrient content (\%)}}{100}$$

2.6 Nitrogen use efficiency

Nitrogen use efficiency (NUE) generally accounts for the quantity of N accumulated in the plant, showing the N uptake efficiency and the quantity of N utilized in grain production or the N utilization efficiency of the plant to applied N. The NUE components: apparent N recovery efficiency (ANRE), physiological N use efficiency (PNUE), and agronomic N use efficiency (ANUE) were calculated with the following expressions [13]:

$$\text{Apparent N recovery efficiency (\%)} = \frac{(N_f - N_c) \times 100}{NA}$$

$$\text{Physiological N use efficiency (kg kg}^{-1}\text{)} = \frac{G_f - G_c}{N_f - N_c}$$

$$\text{Agronomic N use efficiency (kg kg}^{-1}\text{)} = \frac{G_f - G_c}{NA}$$

where, N_f and N_c refer to the total above ground plant dry matter-N content ($kg\ ha^{-1}$) in the fertilized and control (nonfertilized) plots, G_f and G_c refer to grain yield ($kg\ ha^{-1}$) in the fertilized and control plots, and NA is the amount of fertilizer-N in $kg\ ha^{-1}$ applied.

2.7 Statistical analyses

All the collected data were analyzed for ANOVA with the help of the computer package program MSTAT. The differences among the treatment means were evaluated by the Duncan's New Multiple Range Test (DMRT) as outlined by [14].

3. RESULTS AND DISCUSSION

3.1 Nutrient contents in grain and straw of BRRI Dhan 29

3.1.1 Nitrogen(N)

The N contents in rice grain and straw were significantly influenced by the application of organic and inorganic sources of N over control, ranged from 0.74 to 1.10% in grain and 0.41 to 0.71% in straw (Table 1). The highest N content (1.11%) in grain was observed in the treatment T₁ (application of N₁₅₀ through urea) which was statistically parallel to treatment T₅ and T₆. The lowest (0.83%) grain N was noted in the treatment T₀ (control) where no N fertilizer was applied. So, use of inorganic fertilizers increased the N content in the rice grain markedly. It was noticed that the influence of T₁ on the straw N contents was statistically superior to the other treatments. An increasing tendency of N contents both in grain and straw was noted from the treatments receiving incremental doses of urea N along with cowdung amended N compared to sole cowdung treated plot. The effect of 80% urea N along with 20% cowdung N was more pronounced in both grain and straw N contents than other combinations. It was also conceivable that the N content was comparatively higher than that of straw. Application of S fertilizer increased the N content in straw. A significant increase in N content in rice grain and straw due to application of organic manure and fertilizers have also been reported by many investigators [15,16].

3.1.2 Phosphorus(P)

The P contents in grain and straw of BRRI Dhan 29 were significantly varied by different treatments under the study (Table 1). In case of grain, the maximum P content (0.25%) was recorded in the treatment T₆ and the minimum was found in the treatment T₀ (control). From the Table 1, it was clear that the treatment T₆ was statistically different from only control treatment. The results pinpointed that P supplied to all the treatments in same amounts but the increased amount of P content was found in the T₆ than that of the T₁ treatment. This might be due to the utilization of cowdung N on positive governance on the P content in grain. On the other hand, straw P content was almost lower than grain and varied from 0.12 to 0.21%. The highest P content (0.214%) was observed in the treatment T₆ and was statistically similar to those measured in the treatments T₁, T₂, T₃, T₄ and T₅ having the values 0.21, 0.16, 0.18, 0.19 and 0.20%, respectively. Application of organic N as cowdung either alone or in association with decremental rates of the advocated inorganic N as urea caused pronounced effect in increasing the straw P content. Increase in P contents both in rice grain and straw increased due to application of cowdung, poultry manures and chemical fertilizers were reported by other researches [17,18].

3.1.3 Potassium(K)

The contents of K in grain and straw were governed profoundly by the different treatments (Table 1). The highest content of K in grain (0.31%) was obtained in the treatment T₅, which was statistically similar to that observed in the treatment T₁, T₄ and T₆. The treatment T₀ (control) produced the lowest (0.28%) grain K content, which was statistically dissimilar to all other treatments. The highest K content (0.88%) in straw was obtained in the treatment T₁ which was statistically similar to all other treatments except control. It was also observed that the K content in straw was higher than that of grain in all the treatments. The incorporation of increased doses of cowdung amended organic N combined with reduced doses of inorganic urea N showed better impact in increasing K contents both in grain and straw. K contents in rice grain increased considerably due to application of sulphur fertilizer. [19] reported that K contents in grain and straw were increased due to beneficial effects from combined application of organic and inorganic fertilizers.

3.1.4 Sulphur(S)

Results in the Table 1 indicated that S contents in both grain and straw were statistically insignificant ($p > 0.05$) due to different treatments. The highest value of S contents in grain (0.07%) was obtained from both T₁ and T₆ treatments while the lowest was noted in T₀ (control). All the treatments caused an increasing effect of sulphur content of rice grain. In straw, S contents ranged from 0.03 to 0.05%. The treatment T₀ (control) had the lowest value of S content (0.03%). It was also revealed that the S contents in grain were higher than that of straw in all the treatments. The incorporation of organic N

combined with reduced doses of inorganic urea N showed better performance in increasing S contents both in grain and straw over the control. [20] reported that application of manures and fertilizers increased the sulphur content both in grain and straw of rice.

3.2 Nutrient uptake by grain and straw of BRRi Dhan 29

3.2.1 Nitrogen(N)

Significant effects on N uptake by BRRi Dhan 29 were found in rice grain and straw (Table 2). The N uptake by grain ranged from 21.87 to 64.40 kg ha⁻¹. The effect of T₁ treatment receiving 100% N from urea on N uptake by grain was statistically superior to all other treatments but alliance with the treatment T₆. From Table 2, it was clear that the uptake of N by grain by using 80% N from urea with 80% cowdung amended N exerted greater effect compared to cowdung N alone or other combinations. Application of S increased the N uptake by grain considerably. A significant linear relationship was observed between grain yield and grain N uptake (Figure 1a). The N uptake in grain from different treatments ranged from 18.14 to 43.24 kg ha⁻¹. The uptake of N was found maximum (43.24 kg ha⁻¹) in the T₁ treatment and was statistically superior to the rest of treatments. It was noted that N uptake by grain was higher than that of straw. The total N uptake varied distinctly and ranged from 40.10 to 107.60 kg ha⁻¹. The highest total N uptake (107.60 kg ha⁻¹) was found in the treatment T₁, which was statistically similar to T₆ treatment with total N uptakes of 103.82 kg ha⁻¹. The lowest total N uptake (40.10 kg ha⁻¹) was manifested in the treatment T₀ (control) that was statistically inferior to all other treatments and followed by the T₂ treatment. The total N uptake for the treatments ranked in the order of T₁ > T₆ > T₅ > T₄ > T₃ > T₂ > T₀. [18] and [21] reported that application of N from manures and fertilizers significantly increased the N uptake both in grain and straw of rice.

3.2.2 Phosphorus(P)

There was a significant variation in P uptakes by rice grain and straw due to different treatments. P uptake ranged from 4.32 to 14.85 kg ha⁻¹ in grain and 5.21 to 12.99 kg ha⁻¹ in straw. The highest uptake in rice grain was found in the treatment T₁ and was significantly similar to T₆ with the value of 12.80 kg ha⁻¹. The lowest uptake of P was noted in the control (T₀). From the Figure 1b, it was observed that grain yield was significantly linked with grain P uptake. The highest P accumulation by straw (12.99 kg ha⁻¹) was manifested in the treatment T₁ whereas the lowest (5.21 kg ha⁻¹) was recorded in the treatment T₀. The treatment T₁ was closely succeeded by the treatments T₄, T₅ and T₆. However, all the treatments significantly increased the P uptake over control (T₀). The total uptake was also different due to different treatments (9.52 kg ha⁻¹ in T₀ and 27.84 kg ha⁻¹ in T₁ (Table 2). The treatment T₁ was statistically different from all other treatments on total P uptake but followed by those recorded in the treatments T₅ and T₆. The application of incremental doses of inorganic N performed better in increasing P uptake compared to organic source alone. Similar result was also experienced by other studies [15,17,18].

3.2.3 Potassium(K)

The K uptake ranged from 8.54 to 17.84 kg ha⁻¹ in grain and 30.26 to 53.53 kg ha⁻¹ in straw (Table 3). The highest uptake by grain (17.84 kg ha⁻¹) was found in the treatment T₁, which was statistically dissimilar with the treatments T₄, T₅ and T₆ that gave 12.23, 12.90 and 14.87 kg ha⁻¹ K, respectively. Nonetheless, it was worthwhile to mention that the grain yield was significantly correlated with grain K uptake (Figure 1c). In case of straw, the highest value (53.53 kg ha⁻¹) was appeared in the treatment T₁ and succeeded by that observed in the treatments T₄, T₅ and T₆. However, all the treatments significantly enhanced the uptake over control (T₀). The total uptake was also shaped significantly due to different treatments and ranged from 38.81 to 71.36 kg ha⁻¹ (Table 3). The treatment T₁ had the highest total uptake of K (71.36 kg ha⁻¹) which was statistically identical to T₄, T₅ and T₆ treatments. The least total uptake (38.81 kg ha⁻¹) was noted in the treatment T₀ (control). It was obvious that K uptake by grain was much less than that of straw and the results were in agreement with [22].

3.2.4 Sulphur(S)

Table 3 showed significant effects on S uptake in rice grain and straw as well as in total uptake. All the applied treatments significantly influenced on S uptake by grain over the control treatment (T₀) but the effect of T₁ treatment on S uptake by grain was statistically superior and different from all other

treatments. A linear relationship between grain yield and grain S uptake was observed and presented in the Figure 1d. The S uptake in straw from different treatments ranged from 0.86 to 1.68 kg ha⁻¹ and found maximum (43.24 kg ha⁻¹) in the T₁ treatment that was statistically parallel to T₄, T₅ and T₆ treatments. It was noted that S uptake by grain was higher than that of straw. The highest total S uptake (25.63 kg ha⁻¹) was noted in T₁ treatment, which was statistically different from all other treatments and followed by T₆ treatment with the value of 20.96 kg ha⁻¹. The lowest total S uptake (9.57 kg ha⁻¹) was observed in the treatment T₀ that was statistically inferior to all other treatments. Similar findings were also reported by many researchers [21,23].

3.3 Nitrogen use efficiency (NUE) of BRRi Dhan 29

The response of the N use efficiency (NUE) components due to organic and inorganic sources of N varied as presented in Figure 2. The apparent N recovery efficiency (ANRE) by BRRi Dhan 29 ranged from 8.33% to 45.06% (Figure 2a). The data clearly indicated that the maximum apparent N recovery (45.06%) was obtained with the 100% recommended doses of N from urea (T₁). However, similar results of ANUE (42.54%) was also found in T₆. The reasons for high recovery of applied N could be the split application of urea in rice field that resulted in continuous supply of readily available N from urea throughout the growth period of rice. It was also found that lower rates of urea N with higher rates of cowdung amended N were not useful for the N recovery but the incremental rates of inorganic N with lower rates of organic N improve the recovery of N. Similar observation was also reported in other literatures [24,25]. On the contrary, the physiological N use efficiency (PNUE) varied from 2.40 to 43.55 kg kg⁻¹ (Figure 2b). The peaked value in respect of PNUE was noted in the treatment T₆ (43.55 kg kg⁻¹) followed by the treatment T₁ with the value of 43.50 kg kg⁻¹. Agronomic N use efficiency (ANUE) is a term used to represent the response of rice plant in terms of grain yield to N fertilizer. The range of ANUE varied from 0.20 to 19.60 kg kg⁻¹ with highest value in T₁ treatment and lowest in T₂. This result suggested that application of recommended doses of N through inorganic sources lead to efficient uptake and utilization of applied N. It was also clear that the sole application of N through organic sources had the lowest ANUE. However, the ANUE increased when reduced doses of organic N sources were applied along with incremental doses of inorganic N from urea. These results were also in agreement with the other researchers [18,26].

Table 1. Effects of organic and inorganic sources of N on the nutrient contents in grain and straw of BRRi Dhan 29

Treatments	% N		% P		%K		% S	
	Grain	Straw	Grain	Straw	Grain	Straw	Grain	Straw
T ₀	0.74e	0.41d	0.15b	0.12b	0.28d	0.70b	0.05	0.03
T ₁	1.11a	0.71a	0.24a	0.21a	0.30ab	0.88a	0.07	0.05
T ₂	0.91d	0.51c	0.24a	0.16ab	0.29cd	0.80a	0.06	0.03
T ₃	0.98c	0.59b	0.24a	0.18a	0.29bcd	0.83a	0.06	0.04
T ₄	1.04b	0.60b	0.24a	0.19a	0.30abc	0.82a	0.07	0.04
T ₅	1.06ab	0.62b	0.24a	0.20a	0.31a	0.86a	0.07	0.04
T ₆	1.09a	0.64b	0.25a	0.21a	0.30abc	0.86a	0.07	0.04
CV (%)	2.83	6.66	7.45	7.88	7.43	6.45	2.82	6.89

In a column figure(s) having different letter(s) differed significantly at 5% level of significance by DMRT ($P = .05$)

Table 2: Effects of organic and inorganic sources of N on the N and P uptakes by grain and straw of BRRi Dhan 29

Treatments	N uptake (kg ha ⁻¹)			P uptake (kg ha ⁻¹)		
	Grain	Straw	Total	Grain	Straw	Total
T ₀	21.87d	18.14e	40.01f	4.32d	5.21d	9.52 e
T ₁	64.40a	43.24a	107.60a	14.85a	12.99a	27.84a
T ₂	27.13cd	25.37d	52.50e	6.49cd	8.15c	14.63d
T ₃	33.24c	31.07c	64.30d	7.52bc	9.49bc	17.01d
T ₄	41.78b	34.56bc	76.33c	9.39b	11.04b	20.43c
T ₅	43.89b	36.31b	80.21c	9.83b	11.23b	21.06bc
T ₆	61.21a	42.61a	103.82a	12.80a	11.30b	24.10b
CV (%)	2.52	9.13	8.50	7.53	5.36	3.37

In a column figure(s) having different letter(s) differed significantly at 5% level of significance by DMRT ($P = .05$)

Table 3: Effects of organic and inorganic sources of N on the K and S uptakes by grain and straw of BRR1 Dhan 29

Treatments	K uptake (kg ha ⁻¹)			S uptake (kg ha ⁻¹)		
	Grain	Straw	Total	Grain	Straw	Total
T ₀	8.54d	30.26d	38.81e	8.70e	0.86d	9.57e
T ₁	17.84a	53.53a	71.36a	23.95a	1.68a	25.63a
T ₂	8.48d	39.98a	48.47d	10.07de	1.12cd	11.19de
T ₃	9.85cd	43.89bc	53.74cd	12.04d	1.25bc	13.29d
T ₄	12.23bc	46.80ab	59.03bc	15.45c	1.46ab	16.91c
T ₅	12.90b	49.67ab	62.58b	16.19c	1.49ab	17.68c
T ₆	14.87b	48.24ab	63.11b	19.56b	1.41abc	20.96b
CV (%)	4.77	9.61	7.51	3.37	8.96	2.92

In a column figure(s) having different letter(s) differed significantly at 5% level of significance by DMRT ($P = .05$)




Figure 1: Relationship between the grain yield of BRR1 Dhan 29 and grain nutrient uptake; grain N uptake (a), grain P uptake (b), grain K uptake (c) and grain S uptake (d).




Figure 2: Effects of organic and inorganic sources of nitrogen on apparent nitrogen recovery efficiency (a), physiological nitrogen use efficiency (b), and agronomic nitrogen use efficiency (c) of BRRI Dhan 29. Data are means \pm S.E.M. ($n = 4$).

4. CONCLUSION

The present study was conducted to investigate the effects of different ratios of N supplied either from cowdung and/or urea or from their combinations on the changes in nutrient contents, uptake and use efficiency by BRRI Dhan 29. The results indicated that nutrient contents and uptake by the grain and straw of the crops ranked top when applied with sole application of inorganic N (urea), followed by the application of N in mixture (80% N from urea + 20% N from cowdung), while no nitrogen application (control) ranked in the bottom. The performance of 40, 60, 80, 100% manure amended N level showed a relatively little comparable effect on nutrient contents and corresponding uptake. However, there was an obvious contribution of N supplied from cowdung on N use efficiency components, though the application of recommended dose of inorganic N from urea performed better in this case. Further research should be focused on multiple locations with different paddy soils and climate to scale up the optimum requirement.

REFERENCES

1. FAO. 2013. FAOSTAT agriculture data [online]. Rome, Italy: Food and Agriculture Organization (FAO).
2. Zhao X, Wang SQ, Xing GX. Maintaining rice yield and reducing N pollution by substituting winter legume for wheat in a heavily-fertilized rice-based cropping system of southeast China. *Agric. Ecosyst. Environ.* 2015; 202: 79–89.
3. Islam MR. Effects of different levels of chemical and organic fertilizers on growth, yield and protein content of wheat. *J. Biol. Sci.* 2002; 2(5): 304-306.
4. Islam MB, Salam MA, Hossain MF, Shil NC. Impact of maize based cropping patterns on soil fertility and future research needs. Presented in the Workshop on Soil Fertility, fertilizer Management and Future research Strategy, held at BARC, during 18-19 January, 2010.
5. Ayoub AT. Fertilizer and the Environment. *Nutr. Cycl. Agroecosys.* 1999; 55(2): 117 - 121.
6. Hao XH, Liu SL, Wu JS, Hu RG, Tong CL, Su YY. Effect of long-term application of inorganic fertilizer and organic amendments on soil organic matter and microbial biomass in three subtropical paddy soils. *Nutr. Cycl. Agroecosys.* 2007, 81(1): 17 - 24.
7. Liu MQ, Hu F, Chen XY, Huang QR, Jiao JG, Zhang B. Organic amendments with reduced chemical fertilizer promote soil microbial development and nutrient availability in a subtropical paddy field: the influence of quantity, type and application time of organic amendments. *Appl. Soil Ecol.* 2009, 42(2): 166 -175.
8. Ahrens TD, Lobell DB, Ortiz-Monasterio JI, Li Y, Matson PA. Narrowing the agronomic yield gap with improved nitrogen use efficiency: a modeling approach. *Ecol. Appl.* 2010, 20(1): 91- 100.
9. BRRI. *Adhunik Dhaner Chas* (in Bengali). Bangladesh Rice Research Institute. Gazipur 1701, Bangladesh.
10. Bremner JM, Mulvaney CS. Nitrogen-Total. In *Methods of Soil Analysis Part 2*. Edited by A. L. Page, R. H. Miller and D. R. Keeney. American Society of Agronomy Incorporation and Soil Science Society. Amer., Inc., Madison, Wisconsin, 1982, 595-624 pp.
11. Page AL, Miller RH, Keeney DR. *Methods of Soil Analysis Part II, Chemical and Microbiological Properties*, 2nd edn., American Society of Agronomy, Inc., Madison, Wisconsin, US. 1982.

- 364 12. Hunter AH. Soil Fertility Analytical Service in Bangladesh. Consultancy Report. Bangladesh
365 Agricultural Research Project – Phase II, BARC, Dhaka, Bangladesh. 1982.
- 366 13. Craswell ET, Godwin DC. The efficiency of nitrogen fertilizers applied to cereals in different
367 climates. Adv. Plant Nutr. 1984, 1: 1-55.
- 368 14. Gomez KA, Gomez AA. Statistical Procedures for Agricultural Research, 2nd edn., John Wiley
369 and Sons, New York, 1984, 680 p.
- 370 15. Mamun MAA, Rahman MM, Hossain AKMM, Hoque MM. Effect of organic and inorganic fertilizers
371 on the yield and nutrient uptake of BRRI Dhan 28. Bangladesh J. Prog. Sci. Technol. 2009, 7(2):
372 245-248.
- 373 16. Bhaskaram U, Krishna D. Effect of organic farming on soil fertility, yield and quality of crops in the
374 tropics. XVIth International Plant Nutrition Colloquium. 2009, Sacramento, California, 89-90
375 pp.
- 376 17. Sengar SS, Wade LJ, Baghel SS, Singh RK, Singh G. Effect of nutrient management on rice
377 (*Oryza sativa* L.) in rainfed lowland of Southeast Madhya Pradesh. Indian J. Agron. 2000,
378 45(2): 315 - 322.
- 379 18. Akter S, Islam MR, Rahman MM, Hoque MM. Influences of nitrogen supplied from inorganic and
380 organic sources on the yield, nutrient uptake and nitrogen use efficiency of BRRI dhan29.
381 Bangladesh J. Crop Sci. 2012, 22-23: 151-158.
- 382 19. Rahman MS, Islam MR, Rahman MM, Hossain MI. Effect of cowdung, Poultry manure and Urea-N
383 on the yield and nutrient uptake of BRRI Dhan-29. Bangladesh Res. Publ. J. 2009, 2(2): 552-
384 558.
- 385 20. Mann KK, Brar BS, Dhillon NS. Influence of long-term use of farmyard manure and inorganic
386 fertilizers on nutrient availability in a Typic Ustochrept. Indian J. Agril. Sci. 2006, 76(8): 72-77.
- 387 21. Parvez MS, Islam MR, Begum MS, Rahman MS and Abedin Miah MJ. Integrated use of manure
388 and fertilizers for maximizing the yield of BRRI dhan 30. J. Bangladesh Soc. Agril. Sci. Technol.
389 2008, 5(1&2): 257-260.
- 390 22. Meena SL, Singh S, Shivay YS. Response of rice (*Oryza sativa*) to N and K application in sandy
391 clay loam soils. Indian J. Agril. Sci. 2003, 73(1): 443-447.
- 392 23. Bhuvaneswari R, Sriramachandrasekharan MV, Ravichandran M. Effect of organic sulphur and
393 graded levels of sulphur on rice yield and sulphur use efficiency. J. Intl. Academia, 2007, 2(2):
394 51-54.
- 395 24. Quanbao Y, Hongcheng Z, Haiyan W, Ying Z, Benfo W, Ke X, Zhongyang H, Qigen D, Ke X.
396 Effect of nitrogen fertilizers on nitrogen use efficiency and yield of rice under different soil
397 conditions. Frontiers of Agric. China. 2007, 1(1): 30-36.
- 398 25. Shiferaw N, Heluf G, Sharma JJ, Tareke B. Effects of nitrogen sources and application on yield,
399 yield attributes, and grain protein of rainfed NERICA-3 rice in Gambella, Ethiopia. Intl. J.
400 Agron. Agril. Res. 2012, 2(9):14-32.
- 401 26. Jahan N, Islam MR, Siddique AB, Islam MR, Hasan MM, Shamsuzzaman SM, Samsuri AW.
402 Effects of integrated use of prilled urea, urea super granule and poultry manure on yield of
403 transplant Aus rice and field water quality. Life Sci. J. 2014, 11(8): 101-108.