

Evaluation of Anaerobic Digestate Potential as Organic Fertilizer in Improving Wheat Production and Soil Properties

Original Research Article

ABSTRACT

Integrated use of synthetic and organic fertilizers is crucial to sustainable crop production and stabilization of depleting soil fertility. Keeping in view these certainties, a three-year field study was undertaken to evaluate the potential of anaerobic digestate alone or in integration with chemical fertilizer for improving wheat production and soil fertility. Six treatments viz; control (with no amendment), recommended dose (RD) of chemical fertilizers (CF), anaerobic digestate (AD) on the basis of RD of N, farm manure (FM) on the basis of RD of N, $\frac{1}{2}$ N from CF and $\frac{1}{2}$ N from AD, $\frac{1}{2}$ N from CF and $\frac{1}{2}$ N from FM were applied in Randomized Complete Block Design (RCBD) with three replications. The obtained results revealed that the highest yield (grain and straw), N uptake, NUE, NAE, and NRE were acquired through the utilization of chemical fertilizers which was statistically at par with combined application of anaerobic digestate and chemical fertilizer in all years of study while the minimum was found in control. The integration of digestate not only increase yield but also found to be a monitory feasible strategy. Besides, it was inferred that about half of nitrogenous fertilizer (urea) can be spared with the appropriation of chemical fertilizers and digestate integration.

Keywords: Anaerobic digestate; farm manure; fertilizers; yield; NUE.

1. INTRODUCTION

The sustainability of agricultural productivity is of utmost significance keeping in view the pace of total population growth. It is evaluated that the production of food should expand 70% to sustain flourishing populace, which is relied upon to reach up to 9 billion by 2050 (FAO, 2010). Frequent utilization of chemical fertilizers is practised all through the world to amplify crop production with a specific end goal to satisfy the food needs of growing population (Bos et al., 2005). The prolonged utilization of chemical fertilizers and intensive agriculture have prompted deterioration of soil fertility and additionally caused environmental hazards like ground and surface water pollution from nitrate leaching (Pimentel, 1996). Lack of appropriate crop management, decrease in addition of fertility restoring inputs and unbalance nutrients

application have paved the path for soils to become fragile (Ajayi et al., 2007; Mbah and Onweremadu, 2009).

Under the current scenario, it is indispensable and dire to take suitable measures to check the decrease in soil fertility and profitability. The circumstance immovably emphasis the selection of eco-friendly agricultural practices for keeping up soil fertility and getting crop production on a sustainable basis. For this reason, a sustainable approach is prescribed regarding organic agriculture (Oyewole et al., 2012). Utilization of organic manures would not just be productive in diminishing the unfavourable impacts of synthetic fertilizers, yet in addition will support soil fertility and productivity (Aksoy, 2001; Chowdhury, 2004). Organic manures upgrade food quantity and quality by fulfilling crop nutritional requirement with the provision of essential

*Corresponding author: E-mail: atif_1534@yahoo.com;

nutrients in a way similar to synthetic fertilizers (Liu et al., 2007; Tonfack et al., 2009; Maske et al., 2015).

Moreover, the costs of chemical fertilizers are going past the purchasing capacity of normal landholding farmers and furthermore, sole use of either organic or chemical fertilizer is not appropriate (Wakene et al., 2007). In this way, the present circumstance requests a mix of organic manures with synthetic fertilizers (Jayathilake et al., 2006). To get feasible crop production without declining and weakening soil fertility, an appropriate mix of both mineral and organic manure ought to be adopted (Rekhi et al., 2000). Integration of organic and mineral fertilizers will enhance absorption, distribution as well as nutrient and fertilizer use efficiency (Orkaido, 2004; Jayathilake et al., 2006).

The digestates generated by anaerobic processing of domesticated animals excrements amid biogas generation are commonly rich in macronutrients, for example, N, P, and K, and micronutrients, for example, Zn, Fe, Mo, and Mn (De La Fuente et al., 2013). In like manner, these digestates can possibly be utilized as organic manures and soil amendment in the agricultural land. The advantages of land utilization of digestates include a change in seedling development, crop yield, and fruit/vegetable quality (Feng et al., 2011; Zhang et al., 2015). Besides, the physical, chemical and biological properties of soil can likewise be improved (Riva et al., 2016; Zerkghi et al., 2010). Digestate holds significant amounts of organic matter (20 to 30%), which is necessary for our soils with low organic matter (<1%). In this way, digestate can be a good choice to be integrated with chemical fertilizers for getting optimum crop yield and to recharge soil fertility.

Keeping in view the significance of integrated utilization of organic and inorganic manures, the present examination was conducted to study the potential of digestate as organic manure alone and in combination with mineral fertilizers for enhancing wheat yield and soil fertility.

2. MATERIALS AND METHODS

The current field study was conducted at the farm area of Soil Chemistry Section, Institute of Soil Chemistry and Environmental Sciences, Ayub Agricultural Research Institute Faisalabad, Pakistan for three consecutive years. The investigation was performed by utilizing Randomized Complete Block Design having a

plot size of 5m × 7m and was replicated thrice. Nitrogen was applied at the rate of 120 kg ha⁻¹ as urea, phosphorus 90 kg ha⁻¹ as single super phosphate (SSP) and 60 kg ha⁻¹ potash as sulfate of potash (SOP). Following six treatments were used in the study;

- T₁ = Control (without any fertilizer)
- T₂ = Recommended dose (RD) of chemical fertilizer (CF)
- T₃ = Anaerobic digestate (AD) on the basis of RD of nitrogen
- T₄ = Farm manure (FM) on the basis of RD of nitrogen
- T₅ = ½ N from anaerobic digestate (AD) and ½ N from chemical fertilizer (CF)
- T₆ = ½ N from farm manure (FM) and ½ N from chemical fertilizer (CF)

A calculated amount of farm manure (on the basis of N contents) was well mixed in the soil at seedbed preparation whereas digestate was applied through fertigation with first irrigation. Wheat cultivar Punjab-2011 was sown following recommended methods with the seed rate of 124 kg ha⁻¹ and row to row distance of 22.5cm × 22.5cm.

2.1 Soil Sampling and Analysis

For evaluation of the initial fertility status of the field, a soil composite sample was collected. The collected soil samples were air dried, crushed and sieved through a 2 mm stainless steel sieve and before analyzing for physical and chemical characteristics (Table 1). Soil particle distribution was measured by hydrometer method (Blake and Hartge, 1986). About 250 g of soil was saturated with distilled water for determining the pH of soil. The paste was allowed to stand for one hour and pH was recorded by pH meter with glass electrodes using a buffer of pH 4.0 and 9.0 as standard (McLean, 1982). For determining E_{Ce} (Electrical conductivity of extract), extract of each soil paste was obtained by using a vacuum pump and E_{Ce} was noted with the conductivity meter (Corning 220). Soil organic carbon (SOC) content was estimated TOC analyzer (Shimadzu TOC analyzer; TOC-VCSH) (APHA, 2005) and available phosphorus was estimated by Olsen's method (Jackson, 1962) using the spectrophotometer. For potassium content, soil extraction was done with ammonium acetate (1 N of pH 7.0) and potassium was determined by using PFP-7 Janway Flame photometer (Rowell, 1994). After each crop harvest, soil samples were taken and analyzed to evaluate improvement in soil physicochemical properties

by organic amendments supplementation. Meteorological data for each year of the study is given in Fig. 1.

Table 1. Soil physicochemical properties

Characteristics	Unit	Value
pH	-	8.29
ECe	dSm ⁻¹	1.83
Organic matter	%	0.76
Total nitrogen	%	0.038
Available phosphorus	mg kg ⁻¹	8.83
Extractable potassium	mg kg ⁻¹	200
Sand	%	52.6
Silt	%	21.4
Clay	%	26
Texture textural class	-	Sandy clay loam

2.2 Characterization of Organic Materials

The anaerobic digestate used was collected each year of the study from a biogas plant located at Chak No. 254 RB, Faisalabad, Pakistan. Whereas farm manure was taken from

the dairy farm of Ayub Agricultural Research Institute, Faisalabad, Pakistan. Before application, the samples of both fresh slurry and farm manure were collected and analyzed for chemical constituents by following standard methods (Table 2).

2.3 Plant Sampling and Analysis

At harvest, data regarding grain yield (Mg ha⁻¹) and straw yield (Mg ha⁻¹) was collected in each year of study. Area of 9 m² was harvested from each experimental unit. The harvest of each experimental unit was labelled, sundried and threshed separately. Grain samples were collected and dried at 70 °C for the determination of nitrogen. The dry grain samples were ground and 0.5g sample was digested with the tri-acid mixture of HNO₃-H₂SO₄-HClO₄ for the determination of total nitrogen by Kjeldahl method (Jackson, 1962). Nitrogen uptake, nitrogen use efficiency (NUE), nitrogen recovery efficiency (NRE) and nitrogen agronomic efficiency (NAE) were calculated by using

Table 2. Chemical composition of organic materials

Year	Anaerobic Digestate			Farm Manure		
	N	P	K	N	P	K
	-----%-----			-----%-----		
Year 1	0.96±0.09	0.63±0.07	1.10±0.13	0.57±0.05	0.42±0.03	0.98±0.11
Year 2	1.01±0.11	0.58±0.04	0.92±0.11	0.62±0.07	0.44±0.05	0.86±0.09
Year 3	0.94±0.13	0.62±0.10	0.94±0.08	0.58±0.06	0.46±0.03	0.91±0.12




Fig. 1. Meteorological data showing mean monthly temperature and total rainfall during wheat growing period (2014, 2015, 2016)

following equations as mentioned by Javid et al. (2015).

$$N \text{ uptake (kg ha}^{-1}\text{)} = \frac{N \text{ content (\% grain (dry matter))} \times \text{Yield (kg ha}^{-1}\text{)}}{100}$$

$$NUE = \frac{\text{Wheat grain yield (kg ha}^{-1}\text{)}}{N \text{ fertilizer applied (kg ha}^{-1}\text{)}}$$

$$NRE = \frac{N \text{ uptake (kg ha}^{-1}\text{) in fertilized plot} - N \text{ uptake in control (kg ha}^{-1}\text{)}}{N \text{ fertilizer applied (kg ha}^{-1}\text{)}}$$

$$NAE = \frac{\text{Yield in fertilized plot} - \text{yield in control (kg ha}^{-1}\text{)}}{N \text{ fertilizer applied (kg ha}^{-1}\text{)}}$$

2.4 Statistical and Economic Analysis

The data regarding various traits were subjected to analysis of variance to test the significance of treatments using *Statistix 8.1*[®] (Analytical Software, Tallahassee, USA) and treatment means were compared using least significant difference (LSD) (Steel et al., 1997). A benefit-cost analysis was conducted to estimate the economic feasibility of different organic amendments to increase vegetable production and net economic returns as described by CIMMYT (1988).

3. RESULTS AND DISCUSSION

Table 3. Effect of integrated use of organic and inorganic fertilizer on wheat grain yield (Mg ha⁻¹)

Treatments	Year I	Year II	Year II	Pool
Control	1.64 d	1.68 d	1.75 d	1.69 e
RD of NPK	4.30 a	4.42 a	4.28 a	4.33 a
FS on the basis of RD of N	3.17 c	3.11 c	3.30 c	3.19 d
FYM on the basis of RD of N	3.08 c	2.96 c	3.27 c	3.10 d
½ N from FS + ½ N from CF	4.12 ab	4.28 ab	4.04 ab	4.16 ab
½ N from FYM + ½ N from CF	3.87 b	3.96 b	3.61 bc	3.81 c
LSD	0.36	0.39	0.47	0.22
CV	5.81	5.58	7.70	5.89

* $p < 0.05$

Table 4. Effect of integrated use of organic and inorganic fertilizer on wheat straw yield (Mg ha⁻¹)

Treatments	Year I	Year II	Year II	Pool
Control	3.85 d	3.93 d	3.21 c	3.66 e
RD of NPK	6.22 a	8.12 a	6.68 a	7.01 a
FS on the basis of RD of N	4.86 c	5.65 c	5.23 b	5.25 cd
FYM on the basis of RD of N	4.67 c	4.95 c	5.19 b	4.64 d
½ N from FS + ½ N from CF	5.72 ab	7.63 ab	6.02 a	6.46 ab
½ N from FYM + ½ N from CF	5.52 b	6.89 b	5.67 ab	6.03 bc
LSD	0.63	0.72	0.73	0.84
CV	6.89	7.89	7.67	6.89

* $p < 0.05$

The present three-year investigation was conducted for the evaluation of anaerobic digestate potential as fertilizer alone and in a mix with chemical fertilizers. The three year pool data in regards to wheat yield (Fig. 2) portrayed that highest yield (4.33 and 7.01 Mg ha⁻¹ for grain and straw respectively) was acquired with the use of prescribed quantity of chemical fertilizers which was at par with the yield got in plots accepting digestate in combination with chemical fertilizers in 1:1 ratio (4.16 and 6.46 Mg ha⁻¹ for grain and straw separately). Both treatments displayed similar performance in each year of the study period (Table 3, 4). The treatments with the sole application of either farm manure or digestate on the basis of the recommended dose of N were statistically at par with each other. The highest yield with the application of chemical fertilizer might be attributed to the delivery of nutrients in soluble form in the soil solution, which become available promptly for plants to take up and flourish (Aziz et al., 2010). The improved

nutrient and soil moisture availability could be the reason for a noteworthy increase in yield in response to integration of organic and inorganic fertilizers. The propensity of organic fertilizers for improving soil physicochemical properties as well

as nutrient supplying capability might also be the reason for better yield. The lowest yield in case of farm manure or digestate alone compared to that of chemical fertilizers and their integration might be due to slow release of nutrients needed

by plants for their growth and development (Powon et al., 2004). The findings of the current study are in agreement to the observations of Shaheen et al. (2017) who found higher soybean dry matter yield with the application of chemical fertilizers followed by integrated use of inorganic and organic fertilizer, whereas Noreen and Noreen (2012) found non-significant wheat grain and straw yield with the supplementation of chemical fertilizers alone and combined application of chemical fertilizers (75%) and farm manure (25%). Similarly, Muhammad et al., (2009) and Ayoola and Makinde (2008) found the highest corn cob yield with the application of chemical fertilizers followed by a combination of synthetic fertilizers and organic amendments.

3.1 Nitrogen Uptake and Use Efficiency

The data regarding nitrogen uptake and use efficiency (Figure 3) illustrated that maximum nitrogen uptake (85.7 kg ha^{-1}) was obtained in

treatment receiving nitrogen from chemical source followed by treatment with an integration of chemical fertilizer and digestate (77.2 kg ha^{-1}). A similar trend was obtained for nitrogen use efficiency, nitrogen agronomic efficiency and nitrogen recovery efficiency with values of 36.1, 19.5 and 57.4 kg kg^{-1} correspondingly. Chemical fertilizers contain nutrients in a readily available form which release instantly upon application for plant uptake (Aziz et al., 2010). This is the reason why the nitrogen uptakes, as well as use efficiencies, were maximum in case of application of nitrogen from the synthetic source. The greater nitrogen uptake and use efficiency in case of integrated use of organic and inorganic fertilizer than the sole application of organic amendments may be ascribed to the sustainable and prolonged supply of nutrients (Singh and Singh, 2000; Aziz et al., 2010). More so, the application of organic fertilizer alone results in the slow release from their decomposition by soil microbes (Mahajan et al., 2008; Ghemam and Mourad, 2013).




Fig. 2. Three-year pool grain and straw yield (Mg ha^{-1}) of wheat in response to digestate and chemical fertilizers application




Fig. 3. N uptake and use efficiencies as affected by digestate and chemical fertilizer application

The findings of the present study are consistent with the results obtained by *Haile et al.*, (2012) who found an increase in the application rate of N resulted in significant improvement in N uptake by wheat crop. Similar results were also found by *Shaheen et al.*, (2017) and *Islam and Munda* (2012). However, findings that differ from this study were obtained by *Naing et al.*, (2010) who reported significantly higher agro-physiological N and P use efficiencies for rice with organic-inorganic mixed fertilizers compared to chemical fertilizers or organic fertilizer alone. Whereas *Hossain et al.*, (2010) found higher nutrient uptake with the application of farmyard manure alone compared to that of no fertilizer and inorganic fertilizers application.

3.2 Improvement in Soil Chemical Properties

The findings regarding post-harvest soil chemical analysis (Table 5) depicted that there is a little variation in soil pH and ECe during the studied period. The contents of organic matter improved significantly with the addition of organic fertilizer either alone or in combination with chemical fertilizers. The maximum contents (0.96%) were seen in the treatment receiving the combined application of organic and inorganic fertilizer while the minimum was observed in control without any amendment. Similarly, significant improvements were observed for phosphorus and potassium contents in the current study. The

highest amount of P and K (14.63 and 245 mg kg⁻¹ respectively) were obtained with the application of digestate and chemical fertilizer together.

The non-significant change in soil pH and ECe in this study might be due to the highest buffering capacity of the soil. Similar to the current study, *Yadav et al.*, (2002) found no appraisable change in pH and ECe in response to the application of organic amendments. The improved contents of organic matter, phosphorus and potassium contents could be attributed to the sustained supplementation of organic amendments over the studied period and their subsequent residual effects on the soil properties that last for several years after their application due to their slow mineralization. The findings of our study are in accordance with the results obtained by *Enujeke et al.*, (2013) in a study undertaken for evaluation of the residual effect of organic manure and chemical fertilizer on soil properties. Likewise, *Zahariev et al.* (2014) and *Aladjadjiyan et al.* (2016) also concluded that the application of composted and anaerobically digested manures in the field could improve soil physicochemical characteristics.

3.3 Economic Analysis

The economic analysis (Table 6) depicted that the maximum cost-benefit ratio was obtained with the combined application of chemical

fertilizer and digestate followed by the integration of chemical fertilizers and farm manure while the minimum was in case of the recommended dose

of chemical fertilizer among all treatments except control.

Table 5. Post-harvest soil properties

Treatment	pH	ECe dSm ⁻¹	O.M %	P -----mg kg ⁻¹ -----	K
Control	8.13 ^{NS}	1.74 ^{NS}	0.51 e	7.17 d	170 d
RD of NPK	8.18	1.77	0.68 d	11.06 c	196.7 bc
FS on the basis of RD of N	8.19	1.76	0.82 c	12.87 bc	230 ab
FYM on the basis of RD of N	8.19	1.80	0.86 bc	13.63 bc	215 bc
½ N from FS + ½ N from CF	8.19	1.71	0.96 a	14.93 a	245 a
½ N from FYM + ½ N from CF	8.18	1.74	0.93 ab	14.63 bc	220 b
LSD	0.10	0.08	0.07	0.84	19.6

* $p < 0.05$

Table 6. Economic analysis of the study (Benefit-cost ratio; BCR)

Treatments	Total Expenditure(Rs.)	Gross income (Rs.)	Net income (Rs.)	Benefit-cost ratio (BCR)
Control	24595	50700	26105	2.06
RD of NPK	55986	129967	73981	2.32
FS on the basis of RD of N	35595	95800	60205	2.69
FYM on the basis of RD of N	35595	93100	57505	2.62
½ N from FS + ½ N from CF	40290	122367	82077	3.04
½ N from FYM + ½ N from CF	40290	114400	74110	2.84

4. CONCLUSIONS

The discoveries of the present investigation uncovered that anaerobic digestate can possibly enhance crop production and soil fertility. Wheat yield, nutrient uptake, and soil chemical properties portrayed critical contrasts with the application of various blends of organic and inorganic fertilizers. In any case, the best mix was anaerobic digestate alongside inorganic fertilizers. It not only brought about a significant increase in nutrient uptake and yield of wheat but also proved to be cost-effective and monetary feasible. Furthermore, it was concluded that about half of nitrogenous fertilizer can be saved with the adoption of chemical fertilizer and digestate integration.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

Ajayi, O.C., Akinnifesi, F.K., Sileshi, G., Chakeredza, S. (2007). Adoption of renewable soil fertility replenishment

technologies in the southern African region: Lessons learnt and the way forward. *Nat. Resour. Forum.* 31, 306-317.

Aksoy, U. (2001). Ecological farming. II. In Proceedings of the Ecological Farming Symposium, Antalya, Turkey, 14–16. *Asian J. Agri. Biol.* 5(2), 60-69.

Aladjadjiyan, A., Penkov, D., Verspecht, Ann, Zahariev, A., Kakanakov, N. (2016). Biobased Fertilizers - Comparison of Nutrient Content of Digestate/Compost. *JAERI.* 8(1), 1-7.

APHA, (2005). Standard Methods for the Examination of Water and Wastewater, 21st edn. Total organic carbon (TOC): high temperature combustion method (5310 A and 5310 B)

Ayoola, O.T., Makinde, E.A. (2007). Complementary organic and inorganic fertilizer application: influence on growth and yield of cassava/maize/melon intercrop with a relayed cowpea. *AJBAS.* 1(3), 187-192.

Aziz, T., Ullah, S., Sattar, A., Nasim, M., Farooq, M., Khan, M.M. (2010). Nutrient availability and maize (*Zea mays* L.) growth in soil amended with organic manures. *Int. J. Agric. Biol.* 12, 621–624.

- Blake, G.R., Hartge, K.H. (1986). Bulk density. In: A. Klute (ed.), *Methods of Soil Analysis, Part 1: Physical and Mineralogical Methods*. American Society of Agronomy, Madison, WI, pp. 363-375.
- Bos, C., Juillet, B., Fouillet, H., Turlan, L., Dare, S., Luengo, C., N'tounda, R., Benamouzig, R., Gausseres, N., Tome, D., Gaudichon, C. (2005). Postprandial metabolic utilization of wheat protein in humans. *Am. J. Clin. Nutr.* 81, 87-94.
- Chowdhury, R. (2004). Effects of chemical fertilizers on the surrounding environment and the alternative to the chemical fertilizers. *IES-ENVIS News*. 7, 4-5.
- CIMMYT, (1988). *An Economic Training Handbook*, Economic Program, CIMMYT, Mexico.
- De La Fuente, C., Albuquerque, J.A., Clemente, R., Bernal, M.P. (2013). Soil C and N mineralization and agricultural value of the products of an anaerobic digestion system. *Biol. Fert. Soils*. 49, 313-322.
- Enujeke, E.C., Ojeifo, I.M., Nnaji, G.U. (2013). Residual effects of organic manure and inorganic fertilizer on maize grain weight and some soil properties in Asaba area of delta state. *IJABBR*. 3(3), 433-442.
- Feng, H., Qu, G., Ning, P., Xiong, X., Jia, L., Shi, Y., Zhang, J. (2011). The Resource Utilization of Anaerobic Fermentation Residue. *Procedia Environ. Sci.* 11, 1092-1099.
- FAO. (2010). *Production Year Book; Food and Agriculture Organization of the United Nations: Rome, Italy*. 53, pp. 132-133.
- Haile, D., Dechassa, N., Ayana, A. (2012). Nitrogen use efficiency of bread wheat: Effects of nitrogen rate and time of application. *J. Soil Sci. Plant Nutr.* 12(3), 389-410.
- Hossain, A.T., Rahman, F., Saha, P.K., Solaiman, A.R.M. (2010). Effects of different aged poultry litter on the yield and nutrient balance in boro rice cultivation. *BJAR*. 35(3): 497-505.
- Islam, M., Munda, G.C. (2012). Effect of organic and inorganic fertilizer on growth, productivity, nutrient uptake and economics of maize (*Zea mays* L.) and toria (*Brassica campestris* L.). *Agric. Sci. Res. J.* 2(8), 470-479.
- Jackson, M.L. (1962). *Soil chemical analysis*. Prentice Hall: Inc. Englewood Cliffs, New Jersey, USA.
- Javid, S., Majeed, A., Sial, R.A., Ahmad, Z.A., Niaz, A., Muhmood, A. (2015). Effect of phosphorus fertigation on grain yield and phosphorus use efficiency by maize (*Zea mays* L.). *J. Agric. Res.* 53(1), 37-47.
- Jayathilake, P.S., Reddy, I.P., Srihari, D., Reddy, K.R. (2006). Productivity and soil fertility status as influenced by integrated use of N-fixing biofertilizers, organic manures and inorganic fertilizers in onion. *J. Agric. Sci.* 2(1), 46 - 58.
- Liu, B., Gumpertz, M.L., Hu, S., Ristaino, J.B. (2007). Long-term effects of organic and synthetic soil fertility amendments on soil microbial communities and the development of southern blight. *Soil Biol. Biochem.* 39, 2302-2316.
- Mahajan, A., Bhagat, R.M., Gupta, R.D. (2008). Integrated nutrient management in sustainable rice-wheat cropping system for food security in India. *SAARC. J. Agric.* 6(2), 29-32.
- Maske, S.N., Munde, G.R., Maske, N.M. (2015). Effect of manures and fertilizer on brinjal (*Solanum melongena* L.) C.V. Krishna. *Bioinfolet.* 12, 678-679.
- Mbah, C.N., Onweremadu, E.U. (2009). Effect of organic and mineral fertilizer inputs on soil and maize grain yield in an acid ultisol in Abakaliki-South Eastern Nigeria. *Am. Eurasian J. Agron.* 2(1), 7-12.
- McLean, E.O. (1982). Soil pH and lime requirement, In: A.L. Page, R.H. Miller and D.R. Keeney (eds.) *Methods of Soil Analysis part 2: Chemical and Microbiological Properties*, 2nd Ed. American Society of Agronomy 9, Madison, WI, USA, pp.199-209.
- Muhammad, D., Khattak, R.A. (2009). Growth and nutrient concentrations of maize in press mud treated saline-sodic soils. *Soil Environ.* 28 (2), 145-155.
- Noreen, F., Noreen, S. (2012). Effect of different fertilizers on yield of wheat. *IJSR*. 3(11), 1596-1599.
- Orkaido, O. (2004). Effects of nitrogen and phosphorus fertilizers on yield and yield components of Maize (*Zea Mays* L.) on black soils of Regede, Konso, MSc. Thesis, Alemaya University, Alemaya, Ethiopia.
- Oyewole, C.I., Opaluwa, H., Omale, R. (2012). Response of Tomato (*Lycopersicon esculentum*): Growth and yield, to rates of mineral and poultry manure application in the guinea savanna agro-ecological zone in Nigeria. *J. Biol. Agric. Health Care.* 2, 44-56.
- Pimentel, D. (1996). Green revolution and chemical hazards. *Sci. Total Environ.* 188, 86-98.

- Powon, M.P., Mwaja, V., Aguyo, J.N. (2004). The effect of potassium, phosphorus and farmyard manure on growth and yield of potato (*Solanum tuberosum* L.). Paper Presented At the 9th Triennial Symposium of International Society For Tropical Root-Africa Branch (ISTRC-AB).
- Rekhi, R.S., Benbi, D.K., Bhajan, S. (2000). Effect of fertilizers and organic manures on crop yields and soil properties in rice-wheat cropping system: In Long-term Soil Fertility Experiments in Rice-Wheat Cropping Systems, pp.1-6.
- Riva, C., Orzi, V., Carozzi, M., Acutis, M., Boccasile, G., Lonati, S., Tambone, F., D'Imporzano, G., Adani, F. (2016). Short-term experiments in using digestate products as substitutes for mineral (N) fertilizer: Agronomic performance, odours, and ammonia emission impacts. *Sci. Total Environ.* 547, 206-214.
- Rowell, D.L. (1994). Soil Science: Methods and application. Longman Scientific & Technical, UK.
- Shaheen, A., Tariq, R., Khaliq, A. (2017). Comparative and Interactive Effects of Organic and Inorganic Amendments on Soybean Growth, Yield and Selected Soil properties. *Asian J Agri & Biol.* 5(2), 60-69.
- Singh, V.P., Singh, R.K. (2000). Rainfed Rice: A Sourcebook of Best Practices and Strategies in Eastern India. International Rice Research Institute, p. 292.
- Steel, R.G.D., Torrie, J.H., Dickey, D.A. (1997). Principles and Procedures of Statistics: A biometrical approach, 3rd Ed. McGraw Hill Book Co., New York, USA.
- Tonfack, L.B., Bernadac, A., Youmbi, E., Mbouapouognigni, V.P., Nguenguim, M., Akoa, M. (2009). Impact of organic and inorganic fertilizers on tomato vigor, yield and fruit composition under tropical and soil conditions. *Fruits.* 64, 167-177.
- Wakene, N., Getahun, F., Deressa, A., Dinsa, B. (2007). Integrated use of organic and inorganic fertilizers for maize production. Utilization of diversity in land use systems: Sustainable and organic approaches to meet human needs. University of Rostock, Institute of Land Use, Justus-von-Liebig Weg 6, 18059, Rostock, German, pp.1-18.
- Zahariev A., Kostadinova Sv. Aladjadjian A. (2014). Composting municipal waste for soil recultivation in Bulgaria. *IJPSS* 3(2), 178-185.
- Zerzghi, H., Gerba, C.P., Brooks, J.P., Pepper, I.L. (2010). Long-term effects of land application of class B biosolids on the soil microbial populations, pathogens, and activity. *J. Environ. Qual.* 39, 402-408.
- Zhang, J., Wang, M., Cao, Y., Liang, P., Wu, S., Leung, A.O.W., Christie, P. (2015). Replacement of mineral fertilizers with anaerobically digested pig slurry in paddy fields: assessment of plant growth and grain quality. *Environ. Sci. Pollut. R.* 24 (10), 8916-8923.