

Impact of Insect Infestation on Plant Damage and Yield of Roselle [*Hibiscus sabdariffa* L.] in Benue State, Nigeria

ABSTRACT

Impact of insect infestation on growth and yield of Roselle (*Hibiscus sabdariffa* L.) was evaluated at the Teaching and Research Farm of University of Agriculture, Makurdi, in the 2016 cropping season. An early and late crop (as main plot), of the red (*H. sabdariffa sabdariffa*) and green (*H. sabdariffa altissima*) types (as subplot) were planted in randomized complete block design. Four weekly application of 100 g a.i/ha of cypermethrin + 400 g a. i./ha of dimethoate constituted sub-subplot treatments. All treatments were replicated three times. Insects were visually counted in 1 m² area in two rows of each plot. The dominant insect pests included *Monolepta thomsoni*, *Nisotra sjostedti*, *Dysdercus volkeri* and *Oxycarenus hyalinipennis*. The early crop differed significantly (having 9 % wider stem girth, 2x more branches/stem, and 1.5x more leaf damage) from the late crop. The green Roselle had more pod (2.5x) and seed (1.1x) damage and gave from 1.3 – 1.5x lower calyx, pod and seed yield. Plant growth and productivity were significantly higher in sprayed than in the unsprayed plots. Plants sprayed at both vegetative and reproductive stages were the most productive having significantly more fresh leaf biomass (2.5-103.6x), calyx yield (2.6-2.8x), pod yield (2.2-7.4x), seed yield (3.1-11.0x) sequel to more vigorous growth and less pod damage (2.0-44.6x) and seed damage (1.8-8.6x). Cost-benefit analysis indicated that the red Roselle was more profitable than the green, the late crop was more profitable than the early, and protection at both vegetative and reproductive stages was more profitable than other spray regimes returning N440,291.25/ha, N755,291.5/ha, and N397,236.0/ha for leaf, calyx and seed valuation, respectively. Insecticidal protection of the crop has been shown to mitigate drop damage and return profit.

Keywords: Insect infestation, *Hibiscus sabdariffa*, Growth, Yield, Cypermethrin, Dimethoate

Introduction

Roselle (*Hibiscus sabdariffa* L) from the family Malvaceae is an important vegetable crop in tropical and sub-tropical regions of the world (5, 20). In Africa, the major producing countries of *H. sabdariffa* var. *sabdariffa* include Republic of Benin, Sudan, Cote D' Ivoire, Ghana, Niger, Burkina Faso, and Nigeria (17). The crop has many domestic and industrial uses (5). Locally, the dried red calyx is processed into a colorful cold beverage and the green calyces and bracts are also used to prepare soup/sauce. In many parts of the world, leaves and stalks are consumed as green vegetable/salad (12,17). In Nigeria, Roselle is intercropped with staple food (e.g. yam, maize, sorghum) or oilseed (e.g. Beniseed) crops, or planted along field margins preponderantly by women; they add value to the crop by developing products for the market (14). Different phenological stages of Roselle (seedling, flowering and fruiting stages) are attacked by various insect pests some of them causing economic losses (2,1). In the peasantry, polycultural system of crop production in Nigeria, the inclusion of two or more malvaceous crop is prevalent, thus encouraging cross infestation and damage by insects (3). In spite of the crop's economic prospect, research on the impact of insect infestation on plant damage and yield is limited thus, the aim of this to study.

Materials and Methods

Experimental Site: The study was conducted during 2016 cropping season at the Teaching and Research Farm, Federal University of Agriculture, Makurdi, Benue State. Located in the southern guinea savanna zone of Nigeria. Makurdi lies between 7°44N and 8°35E with an altitude of 228m above sea level.

Seed Source: Two types of Roselle, the red calyx *H. sabdariffa* var. *sabdariffa* and the green calyx *H. sabdariffa* var. *altissima* were purchased locally

Layout: Randomized Complete Block Design (RCBD) with split-split-plot arrangement of treatments replicated three times. Treatments were early- (mid-June) and late-sown (mid-August) crop as the main plot treatments, the varieties *H. sabdariffa* var. *altissima* (Green calyx) and *H. sabdariffa* var. *sabdariffa* (Red calyx) served as the sub-plots treatment and application of 1000ml a.i/ha formulation of cypermethrin + dimethoate at: vegetative growth stage, reproductive growth stage, vegetative and reproductive growth stages, and an untreated control as the sub-sub plot treatments. Each experimental plot measured 5 m x 5m (the Roselle types were planted on a ridge at a spacing of 1m X 1m between and within rows, respectively.); adjacent plots were separated by 1 m alley while space between adjacent replications was 2 m.

Application of Treatment: At vegetative stage, spraying commenced at 3 weeks after planting (WAP) and was repeated four times at weekly intervals while at the reproductive phase spraying commenced at 50 % flowering and was similarly repeated four times.

Data Collection

Plant damage data: At 8 WAP and at harvest, plants in 1 m x 1 m section of rows 2 and 4 of each plot were visually examined, number of insect perforated leaves, flower, pod and calyces per plant per plant quantified. Leaf damage was assessed using leaf damage score where: 0=no leaf damage, 1=25% leaf damaged, 2=26-50% damaged leaf, 3=70% damaged leaves, 4=100% damaged leaves, 5=Total defoliation

66 After pod harvest, drying and shelling was done in order to determine seed yield. One hundred seeds were then selected at
 67 random/plot, soaked in water and the floated seeds (indicative of damage) were counted. Percentage seed damage was
 68 then computed using this formula:

$$\text{Percentage seed damage} = \frac{\text{Number of floated seeds}}{\text{Total number of seeds}} \times 100$$

69 Yield Parameters

70 At harvest, 1 m x 1 m section of rows 2 and 4 were randomly selected and all the leaves of the plants within harvested.
 71 Yellow leaves as well as entangled weeds were removed and the fresh edible/marketable and unmarketable leaves were
 72 then weighed per plot. The calyx and pods from the three inner rows of each plot was harvested and weighed. The calyx
 73 was then sundried and weighed. The pods on plants in the three inner were picked, counted and weighed. The number of
 74 pods/plant then was then computed. Twenty pods were selected at random, shelled, and the seeds gathered and weighed.
 75 The number of seeds/pod was computed. All pods from the three inner rows of each plot were shelled and the seeds
 76 weighed to determine the seed weight/plot. A random of 100 seeds was taken and weighed.

77 **Cost: benefit analysis:** Cost benefit analysis was calculated based on the method of (19). Total crop protection expenses
 78 were calculated by multiplying per spray cost with the total number of sprays throughout the crop growing period, benefit
 79 per hectare was determined by subtracting plant protection expenses from the total income generated per hectare which
 80 was determined based the present market price of the leaves, seeds and calyces of Roselle. Due to fluctuation in prices
 81 throughout the year, price per kilogram of Roselle leaves, calyces and seeds were fixed at 500, 1100 and ₱1150 per kg

respectively for the analysis. Cost benefit ratio of each spray regimes for the different planting dates was worked out by subtracting income of control from net income of spray regimes and the product was divided by total cost of crop protection for each treatment.

Data Analysis -All data were subjected to analysis of variance (ANOVA) using Genstat Software Package and significant means ($P < 0.05$) were separated using Fisher's least significant difference (FLSD) at 5% level of probability.

RESULTS

Effect of cropping season, Roselle variety and spray regimes on plant damage parameters

Numbers of insect- perforated leaves per plant were 1.4 – 1.6 higher ($P < 0.05$) in the early- than in the late-sown crop at both 8WAP and at harvest. There were no significant differences ($P > 0.05$) in leaf damage score index, numbers of insect-perforated calyces (8WAP and AH) and flowers/plant (8WAP), number of insect-perforated pods/plant and percentage seed damage at harvest. However, the number of insect- perforated pods/plant at harvest were significantly 1.9x higher ($P < 0.05$) in the late than the early crop. At 8WAP, insect damage in terms of the numbers of insect-perforated leaves, calyces, flowers per plant, as well as in terms of leaf damage score, was significantly ($P < 0.05$) more on the green- than on the red-type Roselle. At harvest both Roselle types differed significantly ($P < 0.05$) only in the number of insect-perforated pods/plant and percentage seed damage with higher damage occurring on the green-type, but varietal differences in numbers of insect- perforated leaves, calyces and pods/plant as well as in the leaf score index were not significant

($P>0.05$). The untreated plots had significantly greater insect damage except in the plots sprayed at the vegetative stage only where leaf perforations at 8 WAP and at harvest were significantly higher and where fruit boring at 8 WAP was statistically comparable. At 8WAP, calyx perforation was reduced by 61.0 % in the plots sprayed at vegetative stage and by 99.1 % in the plots sprayed at both vegetative and reproductive stages, at harvest, the values were 62.5 % and 96.5 %, respectively. Number of insect perforated pods/plant significantly decreased at both 8WAP and at harvest by 24 and 50.8% in plots sprayed at the vegetative stage, 98 and 99 % reduction were recorded in vegetative and reproductive sprayed plots. Significant reduction over the untreated plot were also observed in the number of insect perforated flowers in the vegetative and vegetative and reproductive sprayed plots amount to 63.87 and 98.95 % respectively. (Table 1)

Effect of cropping season, Roselle variety and spray regime some yield parameters

Planting Roselle early resulted in significantly ($P<0.05$) 1.7x marketable leaf yield than planting the crop late but fresh calyx yield was 1.10 x higher in the late than in the early crop. However, no significant differences ($P>0.05$) in weight of unmarketable leaves, pod, seed, 100-seed weight, percentage dry matter and number of pods/plant (8WAP and AH). In seven out of the 10 yield parameters quantified, the red-type Roselle gave significantly better performance than the green-type (Table 2). Increase in productivity ranged from 5.8 % in 100-seed weight to 34.8 % in fresh pod yield. Differences in weight of unmarketable leaves and the numbers of pods/plant at both 8WAP and at harvest were not significant. All spray regimes resulted in significant increases in yield and component of yield above the level in unsprayed plots (Table 2). The higher the frequency of spraying, the higher the increase in yield. In all yield parameters, the plots sprayed at both vegetative and reproductive stages differed significantly from the plots sprayed at vegetative or reproductive stage alone.

123 **Benefit-cost analysis of production of marketable fresh leaves, dry calyx, and seeds of Roselle**

124 **Across cropping season and variety, profit from the sale of fresh leaves, dry calyx and seeds was 4.0-10.8 fold greater in**
 125 **the plots sprayed at the vegetative and reproductive stages than in other treatment plots. Of the three plant produce, dry**
 126 **calyx production in the early crop was the most profitable (=N=862,030:00), followed by seed production in the late crop**
 127 **(=N=776,854:00), and leaf production of the early crop (=N=674,450:00).The red-calyx Roselle gave >2-fold the profit margin**
 128 **of the green-calyx Roselle (=N=2,094,535:00)and overall profit of the late crop exceeded that of the early crop**
 129 **(=N=4,541,047) by =N1,368,590:00. For seed production, spraying the early crop at the vegetative stage was not profitable.**

130

131

132

133

134

135 **Table 1: Main effects of cropping season, Roselle variety and spray regimes on plant damage parameters**

Variable	Number of insect perforated leaves/plant*		Leaf damage score		Number of insect perforated calyces/plant		Number of insect perforated flowers/ plant	Number of insect perforated pods/plant		% Seed damage
	8WAP	AH	8WAP	AH	8WAP	AH	8WAP	8WAP	AH	AH
Cropping season										
D1	50.73	55.50	2.79	3.08	0.64	1.56	6.05	0.98	1.91	40.21
D2	35.04	35.30	2.17	2.54	0.91	1.64	11.42	1.86	3.16	43.58

FLSD_{0.05}	10.91*	5.66*	ns	ns	ns	ns	ns	0.80	ns	ns
Roselle variety										
V1	52.67	46.80	2.63	2.83	1.03	2.15	10.90	2.07	3.63	43.29
V2	33.10	44.00	2.30	2.79	0.52	1.05	6.56	0.77	1.44	40.50
FSLD_{0.05}	4.14*	ns	0.16*	ns	1.08*	ns	1.70*	0.71*	0.82*	0.70*
Spray regimes										
S1	64.15	59.20	1.92	3.58	0.83	1.62	8.99	2.43	3.95	47.00
S2	48.55	34.50	3.50	1.92	0.14	0.32	0.81	0.18	0.48	27.50
S3	3.97	26.50	1.17	1.25	0.02	0.15	0.26	0.05	0.18	9.67
S4	54.88	41.40	4.30	4.50	2.13	4.32	24.88	3.03	8.03	83.43
FLSD_{0.05}	8.95*	11.66*	0.33*	0.30*	0.33*	0.54*	3.66*	0.63*	0.85*	3.10*
Interaction										
DXV	ns	*	ns	ns	*	ns	*	ns	ns	*
DXS	*	*	*	*	ns	ns	*	*	*	ns
VXS	*	ns	*	ns	*	*	*	*	*	*
DXVXS	*	*	*	ns	ns	ns	ns	*	ns	*

D1=Early sown Roselle, D2= Late sown Roselle, V1=Green type, V2=Red type,S1=Vegetative stage only, S2=Reproductive stage only, S3=Vegetative and Reproductive stage, S4= Untreated Control, WAP=Weeks after planting, AH=At harvest

144 **Table 2: Main effects of cropping season, Roselle variety and spray regimes on plant on some yield parameters**

145

Variable	Leaf yield (Kg/ha)		Calyx yield (Kg/ha)		Number of pods/plant		Pod yield (Kg/ha)		Seed yield (kg/ha)	100-Seed Weight (g)
	Marketable	Unmarketable	Fresh	Dry	8WAP	AH	Fresh	Dry		
Cropping Season										
D1	1827.32	1014.14	2340.89	250.75	9.58	83.20	2103.10	253.96	221.77	2.32
D2	1105.25	645.15	2587.54	288.35	13.49	73.40	2286.09	309.18	302.17	2.33
FLSD _{0.05}	139.83*	ns	89.47*	ns	ns	ns	ns	ns	ns	ns
Roselle Variety										
V1	1256.02	764.36	1973.81	220.51	9.58	64.10	1732.78	230.88	211.44	2.26
V2	1676.55	894.93	2954.62	318.58	13.50	92.50	2656.41	332.25	304.15	2.40
FSLD _{0.05}	376.83*	ns	415.91*	46.88*	ns	ns	563.93*	70.95*	56.49*	0.04*
Spray Regimes										
S1	425.69	735.77	1727.22	202.83	7.12	51.50	1569.22	222.98	150.74	1.93
S2	1532.36	500.43	2067.66	212.24	12.72	73.40	1877.49	198.26	202.10	2.58
S3	3867.73	152.02	5387.78	589.26	22.73	165.80	4731.27	634.16	633.65	3.16
S4	37.35	540.37	674.20	73.85	3.60	22.50	600.40	70.87	57.36	1.64
FLSD _{0.05}	350.84*	308.13*	618.62*	74.51*	1.10*	17.8*	581.93*	90.10*	67.97*	0.09*
Interaction										
DXV	*	ns	ns	ns	ns	ns	ns	ns	ns	ns
DXS	*	*	*	*	*	*	*	*	*	ns
VXS	*	ns	Ns	ns	*	ns	*	*	ns	ns
DXVXS	*	ns	*	*	*	ns	*	ns	ns	ns

146 D1=Early sown Roselle, D2= Late sown Roselle, V1=Green type, V2=Red type,S1=Vegetative stage only, S2=Reproductive
147 stage only, S3=Vegetative and Reproductive stage, S4= Untreated Control, WAP=Weeks after planting, AH=At harvest

148

149

150

151

152 Table 3: Benefit-cost analysis of production of marketable Roselle fresh leaves at Makurdi in the 2016 cropping season

Croppin g Season	Crop Variety	Crop Stage Protected	Cost of Protection (=N=/ha) ³	Benefit from Protection (=N=/ha) ⁴	Benefit : Cost ratio
Early	HSS ¹	Vegetative	15200.00	50650.00	1.86
		Reproductive	15200.00	323850.00	19.83
		Vegetative + Reproductive	30400.00	674450.00	21.95
		Unsprayed control	0.00	37560.00	
	HSA ²	Vegetative	15200.00	40033.33	1.22
		Reproductive	15200.00	293435.00	15.97
		Vegetative + Reproductive	30400.00	578550.00	18.82
		Unsprayed control	0.00	36700.00	
Late	HSS ¹	Vegetative	15200.00	29675.00	2.95
		Reproductive	15200.00	36035.00	3.37
		Vegetative + Reproductive	30400.00	383280.00	4.98
		Unsprayed control	0.00	0.00	
	HSA ²	Vegetative	15200.00	12955.00	1.85
		Reproductive	15200.00	61115.00	2.07
		Vegetative + Reproductive	30400.00	124885.00	3.42
		Unsprayed control	0.00	0.00	

153 1HSS= Hibiscus sabdariffa var. sabdariffa

154 **2HSA= Hibiscus sabdariffa var. altissimma**
 155 **3Summed over the cost of insecticide, equipment and labour for application.**
 156 **4Difference between income from sale of plant produce and cost of protection. Income was based on the market price of**
 157 **=N=500:00 /kg of fresh leaves**

158

159

160

161

162

163

164

165

166

167 **Table 4: Benefit-cost analysis of dry Roselle calyx at Makurdi in the 2016 cropping season**

Cropping Season	Crop Variety	Crop Stage Protected	Cost of Protection (=N=/ha)³	Benefit from Protection (=N=/ha)⁴	Benefit : Cost ratio
Early	HSS¹	Vegetative	15200.00	81017.00	2.34
		Reproductive	15200.00	331080.00	18.80
		Vegetative + Reproductive	30400.00	862030.00	27.36
	HSA²	Unsprayed control	0.00	60577.00	
		Vegetative	15200.00	67388.00	3.25
		Reproductive	15200.00	250956.00	15.33

Late	HSS ¹	Vegetative +	30400.00	728787.00	23.88
		Reproductive			
		Unsprayed control	0.00	33154.00	
		Vegetative	15200.00	359097.00	14.35
		Reproductive	15200.00	105976.00	7.97
	HSA ²	Vegetative +	30400.00	847224.00	18.90
		Reproductive			
		Unsprayed control	0.00	156200.00	
		Vegetative	15200.00	247139.00	13.55
		Reproductive	15200.00	229066.00	-8.55
		Vegetative +	30400.00	583125.00	16.20
		Reproductive			
		Unsprayed control	0.00	56320.00	

1HSS= Hibiscus sabdariffa var. sabdariffa

2HSA= Hibiscus sabdariffa var. altissimma

3Summed over the cost of insecticide, equipment and labour for application.

**4Difference between income from sale of plant produce and cost of protection. Income was based on the market price
=N=1,100:00/kg of dry calyx.**

181

182

183 **Table 5: Benefit-cost analysis of production of seeds of Roselle at Makurdi in the 2016 cropping season**

Cropping Season	Crop Variety	Crop Stage Protected	Cost of Protection (=N=/ha) ³	Benefit from Protection (=N=/ha) ⁴	Benefit : Cost ratio
Early	HSS ¹	Vegetative	15200.00	-9565.00	0.01
		Reproductive	15200.00	14263.00	1.94
		Vegetative + Reproductive	30400.00	55861.50	2.84
		Unsprayed control	0.00	5462.50	
	HSA ²	Vegetative	15200.00	-11370.50	0.09
		Reproductive	15200.00	7627.50	1.34
		Vegetative + Reproductive	30400.00	16842.00	1.47
		Unsprayed control	0.00	2495.50	
Late	HSS ¹	Vegetative	15200.00	330605.00	14.65
		Reproductive	15200.00	229140.50	16.08
		Vegetative + Reproductive	30400.00	776854.00	26.55
		Unsprayed control	0.00	123050.00	
	HSA ²	Vegetative	15200.00	287894.00	17.00
		Reproductive	15200.00	145995.50	-12.15
		Vegetative + Reproductive	30400.00	739387.00	17.28
		Unsprayed control	0.00	44620.00	

184 **1HSS= Hibiscus sabdariffa var. sabdariffa**

185 **2HSA= Hibiscus sabdariffa var. altissima**

186 **3Summed over the cost of insecticide, equipment and labour for application.**

187 **4Difference between income from sale of plant produce and cost of protection. Income was based on the market price of**

188 **N=1,150:00/kg of seed**

DISCUSSION

The number of insect perforated leaves was more in the early crop than the late crop. This can be attributed to higher number of leaf beetles being more abundant in the early than the late sown crop. However, greater number of pods were found to be perforated in the late sown crop compared to early sown Roselle could be due to the presence of more pod burrowing insects in the late than in the early sown crop. (14) reported okra having more perforated leaves in the early sown crop than the late sown crop this he stated might be due to lower populations of flea beetles in the late sown crop. In a study conducted separately by (11) and (8) they pointed out that planting date can an important tool when planning farm operations so that crops can avoid possible injury by emerging during period of low insect activities. (10) pointed out that biotic or abiotic stress can affect the growth, performance and yield of plants in both agricultural and natural system.

The green and red Roselle types showed differential reactions to infestation and damage by the insect pests. Green Roselle types recorded more damage leaves, calyx, pod, flower and seed than the red type, that is to say the green Roselle type is preferred by insect species more than the red type. It has been reported by (14) that some okra varieties differ in their response to insect infestation and damage. (16) reported more infestation by spiny bollworm *E. insulana* on White than the Sudani and Masri varieties. Phytophagous insects have been reported to discriminate among various host plants, this might be as a result of changes in leaf hardness or as a result of chemical changes by phago-stimulants or the presence of secondary metabolites.(6). Morphological features may produce physical stimuli or bar insect activities (15). (9) also pointed out that insect feeding activity is diminished in many crops because of morphological characteristics which may include pubescence, tissue characteristics and gummy exudates. The red type posses a number of insect and was able to

yield more than the green type this might suggests tolerance of red variety insect infestation, as it was able to produce substantially calyx and leaves. The red Roselle type was noticed to possess spike like structures on pods, this might have served as a deterrent to the insect pests. (9) attributed tolerance in some crops to changes in photosynthetic partitioning which led to high yield because of slight damage of photosynthetic tissues.

Spray plots recorded highest growth parameters, yield parameters and lowest damage parameters. This is in line with the findings of (14) who reported that sprayed okra varieties recorded more number of leaves, total dry matter, lesser number of damage leaves than the unsprayed plots.(18) also reported that spraying cypermethrin + Dimeothate on 2 okra varieties led to significantly taller plants, more branches and leaves. (7) reported that Deltamethrin treated Roselle plants recorded higher calyx yield than the untreated control. The findings of this study is also in line with that which was reported by (4), they pointed out that Okra treated with Oxymatrine-based insecticides recorded lower damage parameters tan the untreated control in both early and late sown okra in Kumasi Ghana. Insecticidal protection of the crop during the vegetative +reproductive stages has been shown to mitigate drop damage and return profit for the two Roselle types.

REFERENCE

1. Abdel-Moniem, A.S.H and Abd El-Wahab, T.E. (2006). Insect pests and predators inhabiting Roselle plants, *Hibiscus sabdariffa* L., a medicinal plant in Egypt. *Archives of Phytopathology and Plant Protection*. 39(1): 25-32
2. Abdel-Moniem, A.S.H, Abd El-Wahab, T.E. and Farag, N.A. (2011): Prevailing insects in Roselle plants, *Hibiscus sabdariffa* L., and their efficiency on pollination. *Archives of Phytopathology and Plant Protection*. 44(3):242–252

- 232 3. Adewole, M. M. (2010). Host Preference in Family Malvaceae by *Podagrica* spp (Coleoptera: Chrysomelidae): an
 233 Oligophagus Herbivore. B. Agric project report. Department of Crop Protection, University of Agriculture, Abeokuta.
 234 42pp
- 235 4. Aetiba, N., P. and Osekre , E., A. (2016). Management of insect pests of okra (*Abelmoschus esculentus* L. Moench)
 236 using oxyamtrine-based insecticide. *Advances in Research*. 6(1): 1-7
- 237 5. Ahmed, M. E., Salheldeen E. A., Abdelrhim A. J., Moayad M. B., Khlid A. I (2012). Effect of weeds on Calcies yield of
 238 *Hibiscus sabdariffa* L in traditional agricultural sector of Sudan. *International Journal of Plant Research*. 2(2): 1-5
- 239 6. Akoroda, O.O (1985). Morphotype diversity in Nigeria land-races of *Corchorus olitorus* L. *Journal of Horticultural*
 240 *Science*. 60(4): 557-562.
- 241 7. Alao F. O., Adebayo T. A., Olaniran O.A. and Akanbi, W.B. (2011).Preliminary evaluation of the insecticidal potential
 242 of organic compost extracts against insect pests of okra (*Abemoschus esculentus* (L.) Moench). *Asian Journal of*
 243 *Plant Science and Research*. 1 (3):123-130
- 244 8. George,K.(2003).FleaBeetle: Organic control options. national centre for appropriate technology (NCAT). Online at:
 245 <http://attra.ncat.org/attra pub/ PDF/fleabeetle.pdf> (accessed July 2016).
- 246 9. Hanelt, P. (1990). Taxonomy, evolution and history. In: Rabinowitch, H. and Brewster, J. (Eds), Onions and allied
 247 crops. 1–26. CRC Press, Boca Raton, Florida, U.S.A.
- 248 10. Higley, L.G., Browde, J.A. and Higley, P.M. (1993). Moving towards new understandings of biotic stress and stress
 249 interactions. In: Buxton, D.R, Shibles, R., Forsberg, R. A., Blad, B. L., Asay, K. H., Paulson, G. M. and
 250 Wilson,R.F.(Eds) *International Crop Science*,749–754. Crop Science Society of America, Madison, WI,U.S.A.
- 251 11. Kemble, A. (1995). Integrated pest management practices for the production of vegetables. ministry of agriculture,
 252 plant protection and regulatory services directorate, Ghana and German development cooperation (GTZ).
- 253 12. Mahadevan, N. and Shivali, K.P. (2009). *Hibiscus sabdariffa* Linn. An Overview. *Natural Product Radiance*. 8: 77- 83.
- 254 13. McClintock, N. (2004). Roselle in Senegal and Mali. LEISA, Magazine on low external input and sustainable
 255 ariculture.20(1): 8-10
- 256 14. Mohammed, I.G., Osipitan, A.A., Pitan, O.R. and Atayese, M. (2013).Evaluation of 15 varieties of okra (*Abelmoschus*
 257 *esculentus* (L.) Moench to field infestation by flea beetles (*Podagrica* spp). *African Entomology*. 21(1):70-78.
- 258 15. Norris, D.M. and Kogan, M. (1980). Biochemical and morphological bases of resistance. 23–61. In: Maxwell, F.G. and
 259 Jennings, P.R. (Eds) *Breeding Plants resistant to insects*. John Wiley and Sons, New York, U.S.A.
- 260 16. Ottai, M.E.S, Abdel-Moniem, A.S.H. and El-Mergawi, R.A. (2004). Effect of variety and location on growth and yield
 261 components of Roselle, *Hibiscus sabdariffa* L. and its infestation with the spiny bollworm *Earias insulana* (Boisd).
 262 *Archives of Phytopathology and Plant Protection*. 37:215–231

- 263 17. Oyewole, C. I and Mera. M. (2010). Response of Roselle (*Hibiscus sabdariffa* L.) to rates of inorganic and farmyard
264 fertilizers in the Sudan savanna ecological zone of Nigeria. *African Journal of Agricultural Research*. 5(17):2305-
265 2309.
- 266 18. Samaila, A. E. and Oaya, C. S. (2014). Efficacy Of insecticidal sprays control of insect pests on the growth and yield
267 of okra at Vinde – Fufore, Adamawa State, Nigeria. *Journal of Agriculture and Veterinary Science*. 7(4):1-4
- 268 19. Shabozoi, N.U., Abro, G.H., Syed, T.S. and Awan, M.S. (2011). Economic appraisal of pest management options in
269 okra. *Pakistan Journal of Zoology*. 43(5):869-878.
- 270 20. Sualey, A. (2013). The influence of spacing and plant age at first harvest on the growth, development, and yield of
271 three Roselle (*Hibiscus sabdariffa* L.) Cultivars. PhD Thesis University of Ghana, Legon.126pp