

Original Research Article

Effect of Nitrogen and Phosphorus on the Yield of Sesame

ABSTRACT

The experiment was conducted at the Research farm of Sher-e-Bangla Agricultural University, Sher-e-Bangla Nagar, Dhaka, during February 2014 to June 2014. BARI Til-3 variety was developed by Bangladesh Agricultural Research Institute (BARI), Gazipur. This experiment consisted of two factors; Factor-A: four levels of nitrogen viz. (i) N_0 (Control), (ii) N_1 (80 kg ha⁻¹), (iii) N_2 (100 kg ha⁻¹) and (iv) N_3 (120 kg ha⁻¹) and Factor B: four levels of phosphorus viz. (i) P_0 (Control), (ii) P_1 (20 kg ha⁻¹), (iii) P_2 (30 kg ha⁻¹) and (iv) P_3 (40 kg ha⁻¹). The experiment was laid out in a Randomized Complete Block Design (RCBD) with three replications. Data were recorded on different parameters such as the number of capsule plant⁻¹, number of seeds capsule⁻¹, the weight of 1000 seeds, seed yield ha⁻¹, stover yield ha⁻¹. The study reveal that combination different level of nitrogen and phosphorous have significant influence on yield of sesame. Highest seed yield ha⁻¹ of sesame (1652 kg) and highest harvest index of sesame (37.33%) were recorded from N_3P_2 (120 kg N ha⁻¹ with 30 kg P ha⁻¹). Therefore, the results suggest that the combined use of 120 kg N ha⁻¹ and 30 kg P ha⁻¹ gave the highest yield of sesame.

Keywords: Sesame, BARI Til-3, Nitrogen, Phosphorous, Sesamum indicum

1. INTRODUCTION

The oldest cultivated plants in the world and indigenous oil plant is known as sesame (*Sesamum indicum* L.) is one of longest history in Indian sub-continent. It is under cultivation in Asia for over 5000 years [1]. Sesame is an important source of edible oil has diverse nutritive values. It is one of the popular oilseeds in Bangladesh which occupies the second position after mustard among the edible oils [2]. Total area coverage of sesame is 87,000 hectares with an annual production of 97,000 metric tons [3]. Its average yield in Bangladesh is 889 kg ha⁻¹ [4]. It has multiple uses for mixing with various food items. Sesame oil is used as hair tonic from very old age in the country. Therefore, it is traditionally cultivated in the different parts of Bangladesh. Among the oil crops, sesame (*Sesamum indicum* L.) has the highest oil content of 46 - 64% [5]. Despite being such an important crop, the productivity of sesame in Bangladesh is very low (889 kg ha⁻¹) in comparison to the global level [6]. Nutrient management is very important for yield improvement of crops [7]. Nitrogen and Phosphorus are important plant nutrients which help in growth and development of plant and ultimately improved crop yield. They involve in many biochemical functions in the physiological system of the plant. Application of nitrogen fertilizer significantly enhanced the growth, nitrogen uptake and yield attributes over control. Nitrogen is the most dynamic nutrient element and becomes the first limiting nutrient as land use intensifies [8-9]. It is taken up in the highest amount by crops and its role in plants cannot be easily substituted [7]. Its supply in the soil is the most important factor limiting growth and yield [10]. Increases in N supply within limits are associated with increase in leaf area and weight, carboxylases and chlorophyll content, all of which determine the photosynthetic activities of leaf and ultimately dry matter production and allocation to the various organs of a plant [11]. Phosphorus is essential parts of skeleton of plasma membrane, nucleic acid, many coenzymes, organic molecules and phosphorylated compounds in plant system [12]. It plays an important role in energy transfer reactions and oxidation-reduction process. Lack of phosphorus, therefore, hampers metabolic

process such as the conversion of sugar into starch and cellulose. Phosphorus is mostly concentrated in the reproductive organ of plant contributing to seed development. A seed needs enough phosphorus and its deficiency, therefore, causes shriveled seed. Thus phosphorus is an important nutrient for seed development and seed filling contributing to better yield formation [13]. Consequently, it increases seed yield of sesame especially under irrigation condition [14]. Therefore, the study was undertaken to observe the crop performance under capricious nitrogen and phosphorus levels and to estimate the optimum and economic levels of nitrogen and phosphorus for yield of sesame. Among the agronomic manipulation, proper nutrient management plays a vital role in getting higher yield. Present investigation was carried out to find the response and sort out the optimum dose of nitrogen and phosphorus fertilizers on yield of sesame varieties.

2. MATERIAL AND METHODS

The experiment was conducted at the Research farm of Sher-e-Bangla Agricultural University, Sher-e-Bangla Nagar, Dhaka, from February to June 2014. The experimental site was located at 23°77' N latitude and 90°3' E longitude with an elevation of 8.5 meters from sea level. The soil of the experimental site belongs to Tejgaon series under the Agro-ecological zone, Madhupur Tract (AEZ - 28), which falls into Deep Red Brown Terrace Soil. Initial soil samples were collected from the experimental plots to a depth of 0-15 cm from the surface before the initiation of the experiment and analyzed in the laboratory (Table 1).

Table 1. Physical and Chemical Properties of the Experimental Soil

Soil properties	Value
A. Physical Properties	
1. Particle Size	
% Sand	29.04
% Silt	41.8
% Clay	29.16
2. Soil Texture	Clay Loam
B. Chemical Properties	
1. Soil pH	5.80
2. Organic Carbon (%)	0.78
3. Organic Matter (%)	1.35
4. Total N (%)	0.08
5. C : N ratio	9.75 : 1
6. Available P (ppm)	22
7. Exchangeable K (me/100g soil)	0.18
8. Available S (ppm)	18

BARI Til-3, a popular variety of sesame developed by Bangladesh Agricultural Research Institute (BARI), was used as planting material for the experiment. The experiment was laid out in a Randomized Complete Block Design (RCBD) with three replications. Individual plot size was 1.2 × 2 m². The row-to-row and seed to seed distance were 30 and 5 cm respectively. Two factors were considered for the study. Factor A, conducted with four different level of Nitrogen (viz. N₀=Control, N₁=80 Kg ha⁻¹, N₂=100 Kg ha⁻¹, N₃=120 Kg ha⁻¹), and Factor B, consisted of four different level of Phosphorous (viz. P₀= Control, P₁= 20 Kg ha⁻¹, P₂=30 Kg ha⁻¹, P₃=40 Kg ha⁻¹). The land was four successive ploughing and cross ploughing and followed by laddering to have a desirable tilth. Experimental plots were fertilized with 5 t ha⁻¹, 45 kg ha⁻¹, 5 kg ha⁻¹, 10 kg ha⁻¹ Cowdung, MoP, ZnSO₄ and Boron respectively except Urea and TSP that were applied as per treatment [2]. All necessary intercultural operations were done whenever required. Five sample plants plot⁻¹ were selected randomly before harvesting of the crop for recording the data of Number of capsules plant⁻¹, Number of seeds capsule⁻¹ and 1000-seed weight. Then plants were harvested, bundled, tagged and recorded the Seed yield and Stover yield per plot and converted it to t ha⁻¹. Biological yield (t ha⁻¹) and Harvest index (%) was calculated by following formula:

$$\text{Biological yield} = \text{Seed yield} + \text{Stover yield}$$

$$\text{Harvest index (HI)} = (\text{Grain yield} \times 100) / \text{Biological yield}$$

2.1 Statistical Analysis

All the collected data were analyzed following the Analysis of Variance (ANOVA) technique and the mean differences were adjudged by Duncan's Multiple Range Test (DMRT) [15] using a computer operated program named MSTAT-C.

3. RESULTS AND DISCUSSION

3.1 Number of Capsule Plant⁻¹

Combined effect of different levels of nitrogen and phosphorus showed significant differences on number of capsule plant⁻¹ of sesame (Table 2). Results designated that the highest number of capsule plant⁻¹ (29.58) was recorded from N₃P₂. On the other time, the lowest number of capsule plant⁻¹ of sesame (14.97) was recorded from N₀P₀. Similar results were found by Maiti and Jana [16]. Mondol *et al.* [17] observed that number of seeds capsule⁻¹ was increased significantly with increasing nitrogen rates.

3.2 Number of Seeds Capsule⁻¹

Combined effect of different levels of nitrogen and phosphorus showed significant differences on number of seed capsule⁻¹ of sesame (Table 2). Results revealed that the highest number of seed capsule⁻¹ of sesame (79.85) was recorded from N₃P₂. The lowest number of seed capsule⁻¹ of sesame (40.59) was recorded from N₀P₀ which was statistically identical with N₀P₁ and N₀P₃. Mondol *et al.* [17] observed that number of seeds capsule⁻¹ was increased significantly with increasing nitrogen rates. Maiti and Jana [16] stated that application of 30 kg P₂O₅ ha⁻¹ produced significantly the highest capsules and capsules plant⁻¹ as compared to other levels of phosphorus.

3.3 1000 Seed Weight (g)

Combined effect of different levels of nitrogen and phosphorus showed significant differences for 1000 seed weight (g) of sesame (Table 2). Results were expressed that the highest 1000 seed weight (3.97 g) was recorded from N₃P₂ which was significantly different from all other treatment combinations. Again, the lowest 1000 seed weight of sesame (2.69 g) was recorded from N₀P₀ which was statistically similar with N₀P₁ followed by N₀P₂ and N₀P₃.

Table 2. Combined Effect of Nitrogen and Phosphorus on Yield Contributing Parameters of Sesame

Treatments	Yield contributing parameters of sesame		
	Number of Capsule Plant ⁻¹	Number of Seeds Capsule ⁻¹	1000 Seed Weight (g)
N ₀ P ₀	14.97	40.59	2.69
N ₀ P ₁	17.63	43.25	2.77
N ₀ P ₂	18.97	54.92	2.85
N ₀ P ₃	17.97	44.59	2.81
N ₁ P ₀	20.30	49.25	2.90
N ₁ P ₁	21.50	54.69	3.19
N ₁ P ₂	27.63	74.79	3.94
N ₁ P ₃	25.22	63.12	3.56
N ₂ P ₀	20.97	50.59	2.99
N ₂ P ₁	28.38	75.84	3.95
N ₂ P ₂	23.26	58.26	3.26
N ₂ P ₃	25.58	69.12	3.81
N ₃ P ₀	21.33	51.59	3.10
N ₃ P ₁	24.29	60.65	3.67
N ₃ P ₂	29.58	79.85	3.97
N ₃ P ₃	26.92	73.20	3.88
LSD _{0.05}	0.7153	3.923	0.09133
CV (%)	6.55	10.18	5.36

N₀: 0 kg N ha⁻¹, N₁: 80 kg N ha⁻¹, N₂: 100 kg N ha⁻¹, N₃: 120 kg N ha⁻¹

P₀: 0 kg P ha⁻¹, P₁: 20 kg P ha⁻¹, P₂: 30 kg P ha⁻¹, P₃: 40 kg P ha⁻¹

3.4 Seed Yield (kg ha⁻¹)

Statistically influence was examined for seed yield ha⁻¹ of sesame was observed by combined effect of different levels of nitrogen and phosphorus (Table 3). Results exposed that the highest seed yield ha⁻¹ of sesame (1652 kg) was recorded from N₃P₂. The treatment arrangement of N₂P₁ (1637 kg) and N₁P₂ (1558 kg) also showed significantly higher seed yield ha⁻¹ and that was expressed as second and third highest seed yield ha⁻¹ respectively but significantly different from all other treatment combinations. The lowest seed yield of sesame (978.80 kg) was recorded from N₀P₀ followed by N₀P₁ (1056 kg) and N₀P₃ (1252 kg) which was the second lowest and third seed yield ha⁻¹ respectively. The results obtained from the treatment combination of N₃P₃ and N₂P₃ also gave promising seed yield but significantly lower than the treatment arrangement of N₃P₂. Kanade *et al.* [18] observed that expressively higher grain yield was obtained with 50 kg N ha⁻¹ and 25 kg P₂O₅ ha⁻¹ compared to 25 kg N ha⁻¹ and 12.5 kg P₂O₅ ha⁻¹. Itnal *et al.* [19] opined that application of 50 kg ha⁻¹ N + 25 kg P₂O₅ ha⁻¹ produced the highest yield, which was 69 percent greater than control. Thorve *et al.* [20] opined that yield attributes and yield of *Sesamum* was increased with every successive increased level of N and P fertilizer and were maximum with 37.5 kg N ha⁻¹ + 18.5 P₂O₅ ha⁻¹.

3.5 Stover Yield (t ha⁻¹)

Statistically influence was examined for stover yield ha⁻¹ of sesame was observed by combined effect of different levels of nitrogen and phosphorus (Table 3). Results exposed that the highest stover yield of sesame (3.10 t ha⁻¹) was recorded from N₁P₁ which was closely followed by N₂P₀, N₀P₂, N₀P₃ and N₃P₀. The lowest stover yield of sesame (2.65 t ha⁻¹) was recorded from N₀P₀ (Control) treatment which was statistically similar with N₀P₁ (2.68 t ha⁻¹) followed by N₁P₂ (2.71 t ha⁻¹) treatment. Jagvir *et al.* [21] observed that stover yield *mustard* is increased significantly with the application of recommended dose of mixed fertilizer (NPKS).

3.6 Biological Yield (t ha⁻¹)

Combined effect of nitrogen and phosphorus had significant influence biological yield of sesame (Table 3). Results exposed that the highest biological yield of sesame (4.56 t ha⁻¹) was recorded from N₁P₁ which was statistically similar with N₂P₀ followed by N₂P₂, N₃P₀ and N₃P₂. The lowest biological yield of sesame (3.63 t ha⁻¹) was recorded from N₀P₀ treatment.

3.7 Harvest Index

Combined effect of nitrogen and phosphorus had significant influence harvest index of sesame (Table 3). Results exposed that the highest harvest index of sesame (37.33%) was recorded from N₃P₂ which was statistically similar with N₂P₁ (36.77%) and N₁P₂ (36.49%) and that was expressed as second and third highest harvest index respectively. The lowest harvest index of sesame (27.06%) was recorded from N₀P₀ which was statistically similar to N₀P₁ (28.33%) followed by N₀P₂ and N₀P₃.

Table 3. Combined Effect of Nitrogen and Phosphorus on Yield Parameters of Sesame

Treatments	Yield Parameters of Sesame			
	Seed yield (kg ha ⁻¹)	Stover yield (t ha ⁻¹)	Biological Yield (t ha ⁻¹)	Harvest Index (%)
N ₀ P ₀	978.8	2.65	3.63	27.06
N ₀ P ₁	1056	2.68	3.73	28.33
N ₀ P ₂	1281	3.06	4.17	29.56
N ₀ P ₃	1252	3.05	4.3	29.15
N ₁ P ₀	1320	2.91	4.23	31.24
N ₁ P ₁	1464	3.1	4.56	32.12
N ₁ P ₂	1558	2.71	4.27	36.49

N ₁ P ₃	1481	2.84	4.33	34.28
N ₂ P ₀	1431	3.08	4.51	31.68
N ₂ P ₁	1637	2.82	4.33	36.77
N ₂ P ₂	1477	2.97	4.46	33.17
N ₂ P ₃	1514	2.8	4.32	35.13
N ₃ P ₀	1429	3.03	4.46	32.07
N ₃ P ₁	1481	2.87	4.35	34.09
N ₃ P ₂	1652	2.78	4.43	37.33
N ₃ P ₃	1520	2.75	4.27	35.66
LSD _{0.05}	4.329	0.05273	0.07457	1.306
CV (%)	7.39	11.56	10.28	9.43

N₀: 0 kg N ha⁻¹, N₁: 80 kg N ha⁻¹, N₂: 100 kg N ha⁻¹, N₃: 120 kg N ha⁻¹
P₀: 0 kg P ha⁻¹, P₁: 20 kg P ha⁻¹, P₂: 30 kg P ha⁻¹, P₃: 40 kg P ha⁻¹



4. CONCLUSION

Form the above findings, it can be concluded that combined effect of nitrogen and phosphorus, N₃P₂ (120 kg ha⁻¹ nitrogen with 30 kg ha⁻¹ phosphorus) had the best performance in respect of yield and yield contributing characters compared to the combinations of control treatment of nitrogen and phosphorus. Therefore, the present experimental results suggest that the combined use of 120 kg N ha⁻¹ and 30 kg P ha⁻¹ along with recommended doses of other fertilizer would be beneficial to increase the yield of sesame variety BARI Til-3.

REFERENCES

- Pham TD, Nguyen TDT, Anders CC, Bui TM. Morphological evaluation of sesame (*Sesamum indicum* L.) varieties from different origins. Australian J. Crop Sci. 2010;4(7): 498-504.
- BARI (Bangladesh Agricultural Research Institute). *Krishi Projukti Hatboi* (Handbook of Agro-technology). Bangladesh Agril. Res. Inst. Joydebpur, Gazipur 1701. 2004;129-132 (In Bangla).
- AIS (Agricultural Information Service). *Krishi Diary* (Agricultural Diary). Agril. Info. Service. Khamarbari, Fanngate, Dhaka-1215. 2014;14 (In Bangla).
- BBS (Bangladesh Bureau of Statistics). Statistical Yearbook of Bangladesh. Stat. Div. Minis. Planning, Govt. People's Repub. Bangladesh. 2011.
- Raja A, Hattab LO, Gurusamy G, Vembu G, Suganya S. Sulphur application on growth and yield of sesame varieties. Int. J. Agril. Res. 2007;2(7): 599-606.
- FAO. FAOSTAT. 2014. <http://www.fao.org/faostat/en/>
- Olaniji, JO, Akanbi WB. Effects of cultural practices on mineral compositions cassava peel compost and its effects on the performance of cabbage (*Brassica oleraceae* L.). Journal of Applied Biosciences. 2008;8(1): 272-279.
- Tiessen KD, Flaten, CA, Grant R, Karamanos E, Burton DL, Entz MH. Efficiency of fallbanded N: Effects of application date, landscape position and inhibitors. In: proc. of 2004 Manitoba Agronomist's conference, Winnipeg. 2003 pp.118-132.
- Akanbi WB, Akande MO, Adediran JA. Suitability of composted maize straw and mineral nitrogen fertilizer for tomato production. Journal of Vegetable Science. 2005;11(1): 57-65.
- Akanbi WB, Akande MO, Baiyewu RA, Akinfasoye JO. Effect of Maize Stover Compost and Nitrogen Fertilizer on Growth, Yield and Nutrient Uptake of Amaranth. Moor Journal of Agricultural Sciences. 2000;1: 6-15.
- Akanbi WB, Oyediran GO, Olaniran OA, Adeyeye SA, Akande MO, Adediran JA. Effects of organic and inorganic fertilizers and their combination on Growth, Nutrient uptake and shoot yield of Celosia (*Celosia argentea* L.). 2006; Science Focus 11-(1) 84—90.
- Panday SN, Sinha BK. Plant Physiology. Vikas Publishing House Pvt. Ltd. 5 Ansari Road, New Delhi, 11002.1986;112-130.
- Sahrawat KL, Islam MS. Phosphorus requirements and management of oil-seeds. In Phosphorus Requirements for Sustainable Agriculture in Asia and Oceania. IRRI (Int. Rice Res. Inst.). Los Banos. Laguna, Philippines. 1990;pp: 371-381.

- 211 14. Kaul AK, Das ML. Oilseeds in Bangladesh. Bangladesh Canada Agriculture Sector Team,
212 Ministry of Agriculture, Govt. of the People's Republic of Bangladesh. Dhaka. 1986;pp: 36-98.
- 213 15. Gomez KA, Gomez AA. Statistical procedures for Agriculture Research. Second Edition.
214 Published by John Wiley and Sons, New York. 1984;pp. 680.
- 215 16. Maiti D, Jana PK. Effect of different levels of nitrogen and phosphorus on yield and yield attributes
216 of sesame. J. Oilseeds Res. 1985;3(1): 252-259.
- 217 17. Mondol DK, Sounda G, Panda PK, Ghosh P, Maitra S, Roy DK. Effects of different irrigation
218 levels and nitrogen doses on growth and yield of sesame (*Sesamum indicum* L.). Indian
219 Agriculturist. 1997;41(1):15-21.
- 220 18. Kanade VM, Chavan SA, Khanvilkar SA. Effects of sowing dates and fertilizers levels on yield of
221 sesamum. J. Maharashtra Agril. Univ. 1992;17: 12-14.
- 222 19. Itnal CJ, Halemani HL, Radder GD, Surkod VS, Sajjan GC. Response of sesamum genotypes to
223 application of fertilizers in drylands. J. Maharashtra Agric. Univ. 1993;18(3): 374-375.
- 224 20. Thorve SB, Katwate MT, Jadhav JD. Response of sesame (*Sesamum indicum* L.) cultivars under
225 varying levels of fertilizers under rainfed conditions. Asian J. Soil Sci., 2011;6(1): 1-10.
- 226 21. Jagvir, S., Monga, D. and Deshmuch, M.-S. (2004). Direct and residual effect of sulphur on
227 growth, yield and quality of cotton (*Gossypium hirsutum*)-mustard (*B. juncea*) cropping system.
228 Journal of Cotton Research and Development, 18(2), 172-174.