

# Chemistry and Mineralogy of Soils Derived from Different Parent Materials in Southeastern Nigeria

E. O. Azu, Donatus<sup>1\*</sup> and V. E. Osodeke<sup>2</sup>

<sup>1</sup>*Department of Horticulture and Landscape Technology, Akanu Ibiam Federal Polytechnic, Unwana, Afikpo Ebonyi, Nigeria.*

<sup>2</sup>*Department of Soil Science and Meteorology, Michael Okpara University of Agriculture, Umudike, Abia State, Nigeria.*

## Original Research Article

### ABSTRACT

Knowledge of the soil minerals is an important index in understanding the soil fertility properties since mineral surfaces serve as both the source and sink of plant nutrients. Soils developed from contrasting parent materials viz: Coastal Plain Sand, Sandstone, Basalt and Shale at different soil horizons (0-20 cm, 20-40 cm and 40-60 cm) in South Eastern Nigeria were studied for mineralogical properties. Soil samples were collected from the different parent material in three replicates and a total of thirty six composite samples were collected. These samples were subjected to X-ray diffraction analysis with Siemens D5000 diffractometer using Cuk $\alpha$  radiation with Iron(Fe) filter ( $\lambda = 1.5409\text{\AA}$ ) at 40kv and 30mA, at a scan rate of  $2^\circ$  per minute. The experimental design was a 4 x 3 factorial in Randomized Complete Block Design (RCBD). Statistical analysis revealed significant ( $P < 0.05$ ) variation in soil physical and chemical properties with parent material and soil depth except organic matter "OM", total nitrogen "TN", magnesium " $\text{Mg}^{+2}$ " total exchangeable acidity "TEA" and base saturation "BS" where variations were not significant. However, interaction between parent material and soil depth was only significant in influencing the pH. Top soil layers had a significant greater amount of OM, TN, AP, cations, Sand and Silt. Generally, soil of shale formation was superior, followed by basaltic soil in these soil fertility indicators, while soil of Coastal Plain Sand formation had the least amounts. Parent material and depth played dominant roles in the type and distribution of clay minerals in the study area. The concentration of these minerals varied with parent material and soil depth. Significant variations with parent material were observed in the minerals identified except in Halloysite and Chlorite where non-significant variations were observed. While variations with depth were statistically significant ( $P < 0.05$ ) in Chlorite, Goethite, Hematite, Montmorillonite and Quartz, the influence of the interaction between parent material and soil depth were only significant in Gibbsite, Kaolinite and Montmorillonite. Soil derived from Coastal Plain Sand had, Kaolinite (9.94 to 28.83 %) and Quartz (56.4 to 87.3 %) as the dominant clay minerals and these decreased with depth. Soil formed in shale had mixed clay mineralogy of montmorillonite (14.37 to 21.88 %), goethite (23.58 to 25.09 %), hematite (20.39 to 24.19 %), gibbsite (8.48 to 12.10 %), kaolinite (6.89 to 3.56 %) and others with depth. Soil formed in sandstone has Kaolinite accounted for 26.42 to 31.26 %, goethite, 7.77 to 14.87 %, and quartz accounted for 35.4 to 51.8 %. Other minerals identified were: hematite (4.58 to 14.62 %), gibbsite (1.70 to 5.23 %), while soils derived from basalt had kaolinite (14.64 to 27.44 %), quartz (23.7 to 42.5 %), hematite (11.0 to 23.14 %), goethite (13.94 to 26.48 %) as the dominant crystalline minerals present in the soil. The study concludes that the mineralogy of Southeastern Nigeria consists of quartz, kaolinite, hematite, goethite and gibbsite with traces of smectite in soil derived from shale as the dominant minerals and their percent mass occurrences

\*Corresponding author: E-mail: donblessed01@gmail.com;

varied with soil parent material and depth. Since the mineralogy of the soils studied is mainly kaolinitic, the soils can be said to be low in terms of fertility and therefore require adequate soil fertility management strategies to enhance their fertility and productivity.

*Keywords: Soil chemistry; clay mineral; parent material and soil depth.*

## 1. INTRODUCTION

The challenge for agriculture over the coming decades will be to meet world increasing demand for food in a sustainable way (Idris and Ahmed, 2012). Declining soil fertility and management of plant nutrients have made this task more difficult (Gruhn *et al.*, 2000). With increasing demand of agricultural production due to the drastic increase in global population, there is urgent need to outline strategies that will ensure sustainable production of food through a holistic approach to soil nutrient management (Jones *et al.*, 2013). A decisive and holistic approach to soil fertility study will in addition to investigating the nutrient content in a soil include a study of the mineralogical properties.

Soil minerals serve as both sources and sinks of essential plant nutrients. As primary minerals that originally formed at high temperatures and pressures in igneous and metamorphic rocks are weathered in soils, they release plant nutrients into the soil solution. New minerals form in the aqueous phase of soil environments. These secondary minerals which dominate the clay fraction of the soil (Clay minerals) serve as sources of nutrients themselves, or they precipitate or adsorb essential elements, keeping them from being taken up readily by plants. In many cases, secondary minerals serve as important reservoirs where nutrients are held strongly enough to prevent leaching, yet weakly enough to allow plants to draw on them to meet their nutritional needs. The availability of different nutrient elements such as P, Ca, K, NH<sub>4</sub>, Mg, etc is largely influenced by the clay minerals present in such soils.

Soils show tremendous variability with regard to their mineralogical properties (Nsalambi and Christopher, 2010). The types and amount of minerals present in soils from different regions vary greatly, hence the mineralogical constitution of soil is rather complex.

The mineralogical composition of the clay fraction of a soil is one of the critical factors in determining soil chemical and physical properties. In natural soils, Borggaard and Elberling (2004), concluded that mineralogy can be decisive for soil sensitivity and resilience,

including the capacity of the soils to resist stress and recover after exploitation. Mineral types and amounts are influenced by several factors such as climate, topography, vegetation and bedrock type (Libo *et al.*, 2015). Clay minerals are excellent tracers of weathering processes of bedrocks, especially some typical clay minerals such as kaolinite, montmorillonite and illite (Tang, *et al.*, 2002). Quantitative analysis of clay minerals and their alteration provides invaluable information on the pedogenic history of soils (Wilson, 1999). Clay mineralogy in soils helps in better understanding soil genesis and development (Moore and Reynolds 1997). Moreover, the types and amount of clay minerals are considered as important constraints on the physical and chemical properties of soil.

Kaolinite, Quartz and different forms of iron (Fe) and aluminium (Al) oxides have been reported as the dominant minerals in the clay fraction of the highly weathered soils of most tropical and subtropical regions (Nitzsche *et al.*, 2008; Schaefer *et al.*, 2008). The concentration of these minerals in the soil and their characteristics such as crystalline and specific surface area vary with the parent materials, intensity of weathering, composition of soil solution and drainage conditions (Schaefer *et al.*, 2008) and may also be affected by soil management system (Silvaneto *et al.*, 2008).

The mineralogy and geochemistry of different geological formations such as shale, sandstone, coastal plain sand etc, influence to a large extent the nature of the overlying soils (Szilas *et al.*, 2005). Norra *et al.* (2006) observed that the mineralogical and geochemical components of surface soils reflect the soil parent material and pedogenic minerals. Parent material has been recognised as an important factor in soil formation since the earliest scientific consideration of soils (Jenny, 1980). According to Joffe (1949) the formation of soil in a region can occur within a certain period of time depending on the parent material, climate, topography and vegetation of the region (Buol *et al.*, 1980; Dinc *et al.*, 1987). Different parent materials affect the morphology and chemistry of soils under the same conditions, such as topography and vegetation (Irmak *et al.*, 2007). Differences in

physical, chemical and mineralogical properties of soils are related primarily to parent material (Washer and Collins, 1988).

The clay mineral characteristics of a given soil type influence largely the capacity of the soil to supply nutrient to plants and the kinetics of its ability to replenish the soil solution after the depletion of the rhizosphere (Hinsinger *et al.*, 2009). Different types of clay mineral hold and retain different kind and amount of nutrients. Therefore characterising the clay minerals present in a particular soil can be an important index in understanding and possibly predicting the degree to which the soil can retain and supply nutrients to plants. Also, the soil clay mineralogy is closely related to soil fertility and the differences in soil clay mineralogy cause great differences in soil fertility. Thus, characterisation of soils especially as it relates to mineralogical distribution provides useful information on their behaviour, fertility status and their agricultural productivity potentials since mineral surfaces serve as potential sites for nutrient retention and subsequent release to plants (Hinsinger *et al.*, 2009). Much work on the fertility status of the highly weathered soils of Southeastern Nigeria has been done, but little or none of these works provided information on the mineralogy. The unavailability of detailed information on the mineralogical properties of soils of Southeastern informed the choice of this study.

## 2. MATERIALS AND METHODS

### 2.1 Location

The study area, southeastern Nigeria lies between latitudes 4°20' and 7°25'N and longitudes 5°25' and 9°51'E (Nwagbara, 1997). It is made up of Abia, Akwa Ibom, Anambra, Cross River, Ebonyi, Enugu, Imo, and River States and occupies a land mass of about 12 % of the total land area of Nigeria (Odurukwe *et al.*, 1995). It covers an area of about 37,845 km<sup>2</sup> land mass with 60 percent of this area covered by the tropical rainforest (Njoku *et al.*, 2006). It is bounded on the east by the Republic of Cameroon, on the North by Benue State and on the south by the Bight of Benin (Odurukwe *et al.*, 1995).

The climate is essentially humid Tropical with annual rainfall total not exceeding 3500mm (Njoku, *et al.*, 2006). There are two marked seasons in the study area: the wet and the dry seasons. The wet season begins in late April and

last, till October with heavy rainfalls recorded in the months of June, July and September. There is usually a decline of rainfall in August. This is usually referred to as "August break" (Obasi *et al.*, 2015). Current mean maximum temperature of the area is 32°C while the mean minimum is 21°C, and a relatively high mean annual relative humidity, exceeding 75 % (Njoku *et al.*, 2006).

Vegetation stretches from mangrove swamp in the coast through rainforest to derived savanna in the interior. According to Okorie and Okpala, (2000), the vegetation is essentially secondary forests tending towards derived savannah because of population pressure and repeated annual bush burning.

### 2.2 Geology

The extensive soil systems and landscape of South-eastern Nigeria are underlain by sedimentary materials of different ages and different formations (Igwe *et al.*, 2015). The geological formations were described by Ofoegbu (1985) as materials within the "Benue trough". They are mainly weathered sandstones of Palaeocene/Pleistocene age and shales of the basement complex of Precambrian age (Oriajiaka, 1975). The underlying geology consist of heterogeneous materials, namely Basement Complex, Beach Sands, Coastal Plain Sand, Mangrove Swamp Deposits, Sandstones, Somberirio Warri Deltaic Deposits, Recent and Sub-recent Alluvium (FDALR, 1985). Crystalline rocks mainly gneisses and related rocks (metamorphic) are found around Oban hills and Obudu highlands, while basalt (igneous rock) is present around Ikom in Cross River State (Jungarius and Lavett 1964). Most of the reference are however underlain by sedimentary rocks from the earlier ones of the lower cretaceous era, through those of tertiary period (e.g. Imo shale group, the coastal plain sands, the Bende-Amaeke formation etc) to those of the quaternary period (alluvium) (Oriajiaka, 1975).

### 2.3 Soil of the Area

Most soils of this area belong to the order "Ultisol" and are classified as typic Hapludult (Federal Department of Agriculture and Land Resources, 1985; Nwaogu and Ebeniro, 2009). Soils formed on coastal plain sand and sandstone occupy most of Imo, Abia and Akwa Ibom States, a reasonable part of Rivers, Bayelsa States and a little of Cross River State. These soils are usually acidic, have low CEC, low base saturation and low fertility status (Okunami, 1981). Soils formed on shale are

reddish-brown and gravelly. They have high clay content, poorly drained, acidic with high exchangeable cations but are generally fertile. Soils formed from basalt are highly weathered, strongly acidic and a good number of them are low in available P, having high fixation capacity for P (Osodeke and Uba, 2005).

## 2.4 Soil Sampling

Thirty six soil samples representative of twelve (12) locations of contrasting parent materials comprising of Abakiliki (Shale), Afikpo (shale), Ikom (Basalt), Bende (Shale) Umudike (Coastal plain sand), Nssuka (Basalt), Etung (Basalt), Ikot-Ekpene (Coastal plain sand), Aba (Coastal plain sand), Amaeke (Sandstone), Ohafia (Sandstone), and Okigwe (Sandstone), at three depths (0-20 cm, 20-40 cm and 40-60 cm) were collected using soil auger (Table 1 and Fig. 1).

## 2.5 Sample Preparation

The soil samples were air-dried, crushed and sieved through a 2mm sieve (Pansu and Gautheyrou, 2006). They were bagged and labeled accordingly for laboratory analyses. The samples chemical and mineralogical content were analysed according to standard Laboratory procedures.

## 2.6 Laboratory Analyses

Particle size distribution: This was determined using the Bouyocous hydrometer method as described by Benton (2001). Soil pH was determined in soil to water and soil to  $\text{CaCl}_2$  at the ratio of 1:2.5, using glass electrode pH meter (Udo *et al.*, 2009). Organic carbon was determined by the wet oxidation method of Walkley and Black as described by Pansu and Gautheyrou (2006). Total nitrogen determination was done by the macro Kjeldahl digestion method (Simmone *et al.*, 1994). Exchangeable acidity was determined by the 1M KCl extraction procedure as described by Udo *et al.*, (2009). Exchangeable basic cations ( $\text{K}^+$ ,  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{Na}^+$ ) were determined by the ammonium acetate method (Carter and Gregoich, 2008). Ca and Mg in the extract were determined using the atomic absorption spectrophotometer, while K and Na were read using the flame photometer. Effective Cation Exchange Capacity (ECEC) was obtained by summation of all the exchangeable cations and exchangeable acidity as described by Udo, *et al.*, (2009). Base Saturation was obtained mathematically with:

$$\text{BS (\%)} = \frac{\text{TEB}}{\text{ECEC}} \times \frac{100}{1}$$

BS = Base saturation, TEB = Total Exchangeable Bases, ECEC = Effective Cation Exchange Capacity.

## 2.7 Mineralogical Analysis

The qualitative determination of the mineralogical properties of the soil samples using X-ray diffraction were carried out at Sci-Ba Laboratory and Scientific Consultant, Cape Town South Africa. About 2g of each soil sample was dried, pulverised and milled using McCrone Mill with agate grinding element in a Jar. The unique grinding action of the mill rapidly reduces the soil particles to sub-micrometer sizes and mixes the sample for homogenisation required for qualitative analytical methods (Approximately 10  $\mu\text{m}$ ). Pre-analysis sample preparation procedures included the removal of soluble salts and gypsum by washing out according to the method of Konse and Rich, (1959). Carbonates were removed by neutralisation with sodium acetate (pH =5), according to Grossman and Millet, (1961), while organic matter was removed by oxidation with  $\text{H}_2\text{O}_2$  (Konse And Rich, 1959). These samples were saturated with Magnesium ( $\text{Mg}^{2+}$ ) and Potassium ( $\text{K}^+$ ) using magnesium chloride ( $\text{MgCl}_2$ ) and potassium chloride (KCl) solutions. About 0.5g of dried and prepared soil sample each was used for the X-ray analysis. The X-ray studies were performed with Siemens D5000 diffractometer using  $\text{CuK}\alpha$  radiation with Iron (Fe) filter ( $\lambda = 1.5409\text{\AA}$ ) at 40kv and 30mA, at a scan rate of  $2^\circ$  per minute (Osabor *et al.*, 2009). The diffraction patterns were obtained with the aid of computer, while the  $2\theta$ , d-values and peak intensities yielded by the powder patterns were used to identify the minerals (Akhirevbulu *et al.*, 2010). The interpretation of the diffractograms obtained for each sample was done by comparing the peaks obtained with those of standard minerals established by Brown (2008) and Joint Committee of Powder Diffraction Standards (1980).

## 2.8 Statistical Analysis

This study was considered a 4 x 3 factorial in Randomized Complete Block Design (RCBD) in which the parent material and soil sampling depths were the two factors under consideration.

The data obtained were subjected to simple descriptive statistics and analysis of variance (ANOVA) as outlined by Steel and Torrie (1980)

to test the differences between means. Least Significant Difference ( $FLSD_{0.05}$ ) was used to separate the means that were significantly difference.

### 3. RESULTS AND DISCUSSION

#### 3.1 Physical and Chemical Characteristics of the Soils Used for the Study

The physical and chemical characteristic of the soils used in the study are presented in Table 2. The results of the soil analysis showed that there were significant variability in both the physical and chemical properties among parent materials and sampling depth. Particle size analysis showed that the soils varied from sandy-loam, sandy-clay-loam to clay-loam. Soils of shale and basalt formations were generally clayey-loam irrespective of depth, while soils of coastal plain sand formation were more of sandy-clay-loam especially at the surface horizon. The variability observed in the textural properties of the soils studied could be a reflection of the differences in the parent materials (Enwezor *et al.*, 1990). Parent material and soil depth significantly ( $P < 0.05$ ) influenced the textural properties of the soils studied. The sand component ranged from 77.1 % in soil of coastal plain sand formation to 23.8 % in soil formed from shale. The results showed that sand is the predominant particle size in soils formed on coastal plain sand. The sand component decreased with soil depth which is in accord with the findings of Aki and Antigha, (2015). Generally, it was observed that soils formed from shale and basalt had the least amount of sand, which is in line with the reports of Oti, (2015) and Osodeke and Ubah, (2005) for soils of southeastern Nigeria.

Silt fraction ranged from 26.53 % in soils of basalt formation to 10.07 % in soil of shale formation and this did not show a consistent distribution trend down the soil profile. Similarly, soils formed from Shale and Basalt had the highest percentage of silt, while soils formed from Coastal Plain Sand had the least proportion of silt. It was also observed that only the effect of parent material was significant ( $P < 0.05$ ) in influencing the silt distribution in these soils.

Percentage clay fraction distribution among the soils ranged from 62.2 % to 12.6 %. The clay content generally increased with soil depth and the highest clay contents were recorded in soils formed from Shale and Basalt. Other researchers such as Osodeke and Ubah, (2005), Wissen *et*

*al.*, (2015) have also reported similar trend in clay distribution of tropical soils. The relative low clay content at the surface horizon (0 -20 cm), may be due to the sorting of soil materials by biological and agricultural activities, clay migration or combination of these activities as earlier reported by Malgwi *et al.*, (2000) and Adegbenro *et al.*, (2011).

Soil pH varied widely with soil depth and parent material. Apart from soils formed from coastal plain sand which exhibited slightly reverse trend, soil reaction in all soil types was slightly acidic at the surface horizon, grading towards strongly acidic down the soil profile. This observation can be due to high leaching potentials that characterised the acid sands of southeastern Nigeria as conditioned by the heavy tropical rainfall leading to loss of basic cations from the surface horizons as noted by Osodeke (1999) and Udo *et al.*, (2009). Upper horizons receive maximum leaching by rainfall, and by dissolved carbonic acids and organic acids which remove metal cations (e.g.  $Ca^{++}$ ,  $K^{+}$ ) and replace them with  $H^{+}$  ions, but the lower horizons are not so strongly leached, in fact, the dryer areas may accumulate calcium and other materials removed from the upper soil. The strongly acidic conditions observed at the lower depths in soils formed from shale and basalt could be attributed to the increasing clay content with its associated high Fe and Al concentrations and 1:1 clay minerals as observed in this study. The highest pH ( $H_2O$ ) of 5.83 was observed in soil of shale formation at 0 -20cm depth while the least (3.10) was noted in basaltic soils (Table 2). Other researchers (Onwuka *et al.*, 2007; Eneje and Azu 2009) have reported acidity in most soils of Southeastern Nigeria. Low pH is basically an indication of the presence of  $H^{+}$  or acidic cations which displaces the basic cations and result to low nutrient and low crop yield (Brady and Weil, 2008). The parent material, soil depth and their interaction generally showed significant influence on soil pH.

The organic carbon and organic matter content of the soils varied from 2.50 % in soil of shale formation to 0.90 % in basaltic soil and 4.33 % in soil formed from shale to 1.53 % in soil of basalt formation respectively. Variability on both organic carbon and organic matter were observed, but at 5% probability level, only the soil depth showed significant difference. Organic carbon and organic matter content of these soils, irrespective of parent material declined exponentially with depths, probably due to a decrease in rooting density, litter fall, agricultural activities and soil

management practices, which agrees with the report of Bowman and Savory (1992).

Result of soil organic carbon and organic matter exhibited typical surface build-up from litter and root activities. There was no consistent variability in soil organic carbon and organic matter as related to the parent materials, which may probably due to the fact that organic carbon is build up from plant activities and human manipulations on natural soils. Osodeke, (1992) and Orhue and John (2015), have reported a decreasing trend in organic carbon content down the soil profile in most tropical regions of Nigeria. The present values fall within the low and very low category in broad rating of organic carbon (Udo *et al.*, 2009) and since organic matter content is related to soil fertility status (Enwezor *et al.*, 1990), the present values are indications that these soils are low in plant nutrients.

The total nitrogen ranged from 0.27 % in soil of Shale formation to 0.08 % in soil of Sanstone and the nitrogen content followed similar trend of decreasing in quantity with soil depth as organic carbon with the exception of some soils formed on coastal plain sand where direct relationship between total nitrogen and soil depth were observed. This observation could be attributed to the large pore spaces and thus, high porosity that is associated with sand which encourages the eluviation from upper horizons and illuviation of nitrogen at higher depths especially in high rainfall zones (Brady and Weil, 2008). Similarly, this observation could be viewed from the angle of leaching away of nitrates from soil surface horizons due to the relatively high mobility of nitrate in soils, especially in highly porous soils. Most of the soils were low in total nitrogen content except the soil formed from shale which had moderate levels of nitrogen in relation to other soil types and these values were within the critical level set by Adeoye and Agboola (1984) for soils of the humid tropics.

The order of abundance of exchangeable bases in the soils is  $\text{Ca} > \text{Mg} > \text{K} > \text{Na}$ . The exchangeable  $\text{Ca}^{2+}$  ranged from 4.37 Cmol/kg in soil of Shale formation to 1.72 Cmol/kg in soil Coastal Plain Sand formation. Generally,  $\text{Ca}^{2+}$  content decreased with soil depth and soils formed on shale had greater  $\text{Ca}^{2+}$  content relative to other soil types. With the exception of soil of Coastal Plain Sand formed soil at sub-soil levels, all the soils had  $\text{Ca}^{2+}$  level above the critical level of 2.0 cmol/kg reported by Agboola and Corey, (1982). It was also noted that soils from Coastal Plain Sand had the least values of

$\text{Ca}^{2+}$  and these values increased with increased soil depth, indicating evidence of high leaching.

The exchangeable potassium ( $\text{K}^+$ ) varied from 0.37 Cmol/kg in soil of Shale formation to 0.14 Cmol/kg in soils formed from Coastal Plain Sand. Magnesium ( $\text{Mg}^{2+}$ ) and Sodium ( $\text{Na}^+$ ) varied from 1.24 cmol/kg in Basaltic soil to 1.33 cmol/kg in soil of Sandstone formation. The total exchangeable bases were low in line with the reports of other researchers (Adeoye and Agboola 1984; Enwezor *et al.*, 1989; Osodeke and Ubah 2005). In relation to parent materials, it was observed that the exchangeable bases varied and the order of abundance is shale > basalt > sandstone > coastal plain sand. Sodium content did not show a uniform variation among the soils of contrasting parent material. The trend in the distribution of  $\text{Na}^+$  among different soil types was contrary to that of other basic cations.

The highest amounts of  $\text{Na}^+$  were found in soil formed from Coastal Plain Sand while soil formed from Shale had the least values and the order of the relative abundance of  $\text{Na}^+$  is coastal plain sand > sandstone > basalt > shale. Sodium is a dispersing agent and its high proportion in sands may be responsible for their coarse nature and the strong aggregation and plasticity in clay soils may be an indication of low  $\text{Na}^+$  content (Brady and Weil 2008).

While the total exchangeable bases decreased with soil depth for most soil types, the total exchangeable acidity generally increased with soil depth irrespective of parent material. A similar observation was made by Oti, (2015) for some soils of southern Nigeria. Both ECEC and base saturation were low which is an indication of low soil fertility as reported by Udo and Ogunwale (1977).

### 3.2 Mineralogical Composition of the Soils of Studied Area

The XRD results of the mineralogical analysis showed the mineralogical assemblages of the different soil samples. The relative abundance of clay minerals distinguished in the selected soil samples is presented in Table 3. The results indicated the predominating presence of kaolinite quartz, hermatite, goethite and gibbsite within the soils irrespective of the parent material. Identification of other secondary minerals was difficult, because their peaks were obscured by the greater peaks of the major minerals, "kaolinite and quartz". All formations contained small amount of hollysite, but only shale



contained measurable amount of this mineral. Other researchers Odoma *et al.*, (2013), Onyeogu *et al.*, (2016), Akhirevbulu *et al.*, (2010) Osabor *et al.*, (2009) and Nweke *et al.*, (2015) made similar observations for soils of southeastern Nigeria. However, montmorillonite, vermiculite, illite were present in trace amount in some of the soils, especially soils formed on shale at lower soils horizons.

Parent material and depth played dominant roles in the type and distribution of clay minerals in the study area. The concentration of these minerals varied with parent material and soil depth. Significant variations with parent material were observed in the minerals identified except in Halloysite and Chlorite where non-significant variations were observed. While variations with depth were statistically significant ( $P < 0.05$ ) in Chlorite, Goethite, Hematite, Montmorillonite and Quartz, the influence of the interaction between parent material and soil depth were only significant in Gibbsite, Kaolinite and Montmorillonite. The vertical distribution pattern of goethite and hematite, in profile varied with parent material, but contrary to the report of Mehmood *et al.* (2015), the interactive effect of depth and parent material was not significant. This result emphasizes the influence of parent material and soil depth on clay mineral formation and distribution as reported by other Researchers ((Szilas *et al.*, 2005; Norra *et al.*, 2006).

### **3.3 Mineralogical Composition of Soils Derived from Coastal Plain Sand at Different Soil Depths**

Soils formed on Coastal Plain Sands showed that the dominant minerals present were kaolinite and quartz and this is similar in all the pedogenic horizons and this agreed with the report of Chikezie *et al.*, (2010) for soils derived from coastal plain sands of Southeastern Nigeria. Kaolinite accounted to 9.94 to 28.83 %, while Quartz accounted to 56.4 to 87.3 % and these decreased with depth. Other minerals which were present in trace amounts included Goethite (1.41 to 8.81 %), Haematite (0.75 to 3.46 %), Gibbsite (0.36 to 1.33 %) and Mica (0.103 %). The high dominance of quartz in the clay fraction of these soils clearly explains their grittiness and also suggests clay to be of residual origin (Akhirevbulu *et al.*, 2010). The presence of high amount of quartz, which acts as abrasive according to Apuo-Nwosu *et al.*, (2011), could lead to poor physico-chemical performance of the soil. Also, the presence of high amount of

kaolinite in soil of Coastal Plain Sand formation indicates that they must have been subjected to strong chemical weathering with very good leaching, favoured by tropical and subtropical humid climates (Onyeogu *et al.*, 2016).

Similarly, Chikezie *et al.*, (2010) noted that the dominance of kaolinite suggests that the soils are at an advanced stage of weathering. High leaching, together with low pH and removal of basic cations have been reported as the ideal conditions for Kaolinite formation in soils (Branhisel and Bertesch 1992). Warm and humid climate with good drainage have been reported for kaolinite formation (Nadimi and Farpoor 2011; Khormali and Abtahi 2003). Since such an environment is found in the study area at present, kaolinite formation is highly favoured.

Kaolinite is a low activity clay and the effective surface of kaolinite ( $10\text{-}20\text{ m}^2$ ) is restricted to its outer faces, thus, kaolinite has low cation exchange capacity (CEC) with values ranging from 3-15 Cmol/kg (Brady and Weil, (2008). Kaolinite containing soils however, make good bases for road beds and building foundations because of the absence of interlayer spacing between them, but usually poor for agricultural production due to the low CEC except with good fertiliser management policies and practices. When related to the soil nutrient properties, it can be observed that soils formed on coastal plain sands were generally low in fertility status, which can be attributed to the high dominance of kaolinite and quartz in the clay fraction of the soils.

### **3.4 Mineralogical Composition of Soils Derived from Shale at Different Soil Depths**

Their mineralogical compositions of soil of shale formation differed significantly from those derived from other parent materials Table 3. X-Ray Diffraction analysis data showed that soil derived from shale had mixed clay mineralogy of montmorillonite, goethite, hematite, gibbsite, kaolinite and others with depth and smectite was the dominant clay mineral in this soil especially at lower depth.. The presence of smectite was in agreement with the CEC and swelling properties of the soils. The level to very gently undulating landscape, high pH and saturation of the soils with water for a period of time in the study area may have ensured the accumulation of bases and therefore the formation of smectite and other 2:1 clay mineral (Buol *et al.*, 1980; Dinc *et al.*, 1987; Yesilsoy, 1994).

The mineralogy is kaolinitic in the upper 0-20cm (6.89 to 3.56 %) of the profile and mixed mineralogy at the lower depths. At lower depths, the dominant minerals were montmorillonite (14.37 to 21.88 %), goethite (23.58 to 25.09 %), hermatite (20.39 to 24.19 %), gibbsite (8.48 to 12.10 %) and Illite (0.68 to 1.26 %). Chlorite and vermiculite were also present in small amounts (0.05 to 0.483% and 0.523 to 1.283% respectively) at lower depths. The mineralogy of soil of shale formation was much more complex than mineralogy of other soil formations studied. Apart of quartz and kaolinite which decreased in occurrence with depth, other minerals generally increased in proportion with depth. Nweke *et al.*, (2015) reported that illite and montmorillonite are the predominant clay minerals in soil of Abakaliki which is formed from shale. Other researchers have shown that montmorillonite is a significant component of several soils derived from shale in Southeastern Nigeria (Jungerius and Levelt, 1964; Chikezie *et al.*, 2010).

The maximum amount of montmorillonite noticed at lower horizons of soil derived from shale could be as a result of the highly base accumulating environment, such as poor drainage, high water **table** and deep horizons which encourages montmorillonite formation. Barshad, (2006) stated that conditions that impose on the soil a base accumulating environment induces montmorillonite formation. Some researchers have claimed that excess  $Ca^{++}$  content in the soil would increase the formation of montmorillonite but decrease the formation of kaolinite (Yesilsoy, 1994). Many researchers have also reported that montmorillonite was the most abundant clay mineral in most of the soils formed on shale parent material (Singer, 1989; Yilmaz, 1990). Most soils of shale formation under the study areas have hydromorphic properties, poor internal drainage conditions which retard the alteration of these minerals and may account for the dominance of montmorillonite on these soils (Chikezie *et al.*, 2010). Similarly simple transformation of illite to montmorillonite may play a role in increasing the montmorillonite concentration, thereby decreasing the illite concentration on the surface soils, since illite is the main precursor mineral for the formation of smectite in soil, particularly at the surface

horizon. Climatic condition of the study area provides a leaching environment that promotes detachment and release of  $K^+$  from illite. Moreover, the high amount of  $Mg^{2+}$  notice in soil of shale formation, probably created the favourable condition for montmorillonite through transformations at the soil surface ( Owliaie *et al.*, 2006).

Trace to very low amount of vermiculite was observed mostly at the middle horizon (20-40 cm) of the soil. Even though, the alteration of mica is the main pathway for the formation of pedogenic vermiculite in soils (Churchman and Lowe 2012), but Barshad (2006) noted that conditions that impose on the soil an intermediate to moderate environment between highly base accumulating and highly base depleting, favour vermiculite formation.

The soils also contain, in low amounts, kaolinite and illite minerals. For kaolinite formation, the ratio of Si/Al must be under 2.0, the basic cation content must be less,  $pH < 7.0$  (Buol *et al.*, 1980; Dinc *et al.*, 1987; Dixon and Weed, 1989; Sumner, 2000). According to this theorem, the rainfall must be in a sufficient amount to translocate silicon and basic cations to a certain limit in the profile. Furthermore, a low amount of kaolinite mineral might have formed gradually over a very long period of time, due to high weathering and leaching of basic cations as conditioned by heavy tropical rainfall. Also, simple transformation of illite to montmorillonite may play a role in decreasing the illite concentration on the surface soils.

Soil derived from shale had significant higher concentration of Hermatite and Goethite followed by the groups of soils derived from basalt, sandstone and coastal plain sand in that order. Mehmood *et al.*, (2015) also reported similar observations in soils studied. The higher goethite and hermite content in shale derived soil may be due to higher clay content and lothogenic hermatite (Memo *et al.*, 2011; Mehmood *et al.*, 2015) The high hermatite and goethite content at the 40-60 cm horizon may be due to weathering at surface and the resultant release of these minerals that are subsequently leached down and accumulate in the 40-60 cm horizon of soil, relative at higher stage of soil development as reported by Memoon (2008).

**Table 1. Soil parent materials and sample locations**

**Please arrange all data in standard MS Word table format as shown below**




--	--

Parent Material	Sample Location (Town and Cities)
Coastal plain sand	Umudike, Ikot-Ekpene, Aba
Shale	Afikpo, Abakiliki, Bende
Sandstone	Amaeke, Ohafia, Okigwe
Basalt	Ikom, Nsukka, Etung





Fig. 1. Map of Southeastern Nigeria showing sample locations



**Table 2. Some physical and chemical properties of the soils used for the study**  
Please arrange all data in standard MS Word table format as shown below

[illegible]

Parent ECEC	BS Material	Depth (cm)	Particle size distrib. (%)			Texture	pH		Org.C (%)	OM	TN (%)	Na (%)	K (%)	Ca (%)	Mg (%)	TEB (%)	TEA Cmol/kg	
			Sand	Silt	Clay		CaCl <sub>2</sub>	H <sub>2</sub> O										
Basalt	0-20	40.7	26.5	32.8	CL	4.03	5.03	1.90	3.27	0.22	0.09	0.28	3.22	1.24	4.70	2.72	7.42	63
	20-40	26.3	20.3	53.4	CL	3.43	4.33	1.27	2.13	0.16	0.06	0.22	2.76	1.12	4.18	3.32	7.50	55
	40-60	24.6	16.3	59.3	CL	3.10	4.07	0.90	1.53	0.10	0.05	0.21	1.82	0.34	2.41	3.47	5.88	41
CPS	0-20	77.1	10.3	12.6	SCL	3.93	4.90	1.83	3.30	0.13	0.15	0.17	1.91	0.72	3.55	2.45	5.67	59
20-40	72.7	10.1	17.4	SCL	4.20	5.13	1.40	2.30	0.18	0.10	0.20	2.13	1.69	4.41	2.42	7.00	64	
40-60	69.6	11.1	19.3	SCL	4.00	5.00	1.10	1.43	0.16	0.10	0.14	1.72	0.57	3.19	3.11	6.29	51	
Shale	0-20	44.4	19.4	36.2	CL	4.67	5.83	2.50	4.33	0.27	0.07	0.37	4.37	2.43	6.83	2.81	9.64	68
20-40	30.1	16.6	53.3	CL	4.40	5.30	1.53	2.73	0.23	0.02	0.32	3.89	1.68	5.30	3.43	8.73	59	
40-60	23.8	14.6	62.2	CL	3.43	4.90	1.13	2.00	0.12	0.04	0.22	2.93	1.23	3.74	3.95	8.03	48	
SST	0-20	56.9	19.2	23.9	SCL	4.10	5.03	1.97	3.40	0.18	0.11	0.26	3.01	1.70	5.08	3.03	8.44	60
20-40	51.0	16.6	32.7	CL	3.63	4.70	1.70	2.97	0.14	0.09	0.24	2.97	1.62	4.92	3.31	8.56	57	
40-60	45.4	12.4	42.2	CL	3.50	4.57	1.13	1.97	0.08	0.06	0.17	2.65	1.33	4.20	3.43	7.63	54	



0.560	20-40	0.000	0.000	24.54	3.82	14.88	0.00	28.08	0.00	0.00	0.290	32.80
0.000	40-60	0.000	0.383	26.48	10.31	23.14	0.08	14.64	0.00	0.67	0.620	23.70
CPS	0-20	0.103	0.000	1.41	0.36	0.75	0.00	9.94	0.02	0.00	0.00	0.220
87.30		0.000										
72.60	20-40	0.000	0.000	2.84	1.33	2.56	0.00	19.83	0.00	0.00	0.00	0.513
		0.000										
56.40	40-60	0.000	0.000	8.81	1.23	3.46	0.00	28.83	0.400	0.00	0.00	0.907
Shale		0.000										
22.50	0-20	0.017	0.050	23.58	9.62	20.39	0.68	6.89	0.377	14.37	1.137	
		0.523										
13.40	20-40	0.027	0.047	24.43	12.10	21.65	0.92	5.09	0.290	20.09	0.537	
		1.283										
0.627	40-60	0.337	0.483	25.09	8.48	24.19	1.26	3.56	0.000	21.88	0.053	8.20
SST	0-20	0.210	0.000	7.77	1.70	4.58	0.31	31.26	0.557	0.00	1.267	51.80
0.600												
38.70	20-40	0.000	0.307	12.12	3.66	10.57	0.16	30.63	0.683	0.00	0.00	0.357
		1.793										
1.017	40-60	0.000	0.867	14.87	5.23	14.62	0.13	26.42	0.643	0.16	0.227	35.40
GRAND MEAN	0.071		0.178	15.49	5.42	12.65	0.29	18.97	0.248	4.76	0.579	40.50
0.576												
LSD (0.05) DE	NS		0.328	3.129	NS	2.592	NS	NS	NS	NS	1.537	NS
5.20	NS											
LSD (0.05) PM	NS		NS	3.614	2.303	2.993	0.734	3.828	0.431	1.775	NS	NS
6.00	0.529											
LSD (0.05) PM X DE	NS		NS	NS	3.900	NS	NS	NS	6.630	NS	3.074	NS
NS	NS											

CPS= Coastal plain sand, Sandstone, Herm = hermatite, Mot = Montmorillonite, Verm = Vermiculite, PM=Parent material, DE=Depth.

**Please arrange all data in standard MS Word table format as shown below**

[illegible]

	Sand (%)	silt (%)	clay (%)	pH	Org.C (%)	TEB	TEA (Cmol/kg)	ECEC	BS (%)
Chlorite	-0.354	-0.24	0.468	-0.429	-0.479	-0.253	0.617*	0.26	-0.508
Geothite	-0.944**	0.429	0.937**	-0.155	-0.199	0.088	0.715**	0.371	-0.376
Gibbsite	-0.809**	0.427	0.785**	0.097	-0.20	0.210	0.507	0.394	-0.178
Hermatite	-0.914**	0.330	0.933**	-0.092	-0.182	0.109	0.724**	0.402	-0.364
Illite	-0.497	0.059	0.545	0.422	0.100	0.346	0.556	0.595*	-0.037
Kaolinite	0.327	0.171		-0.414	-0.313	-0.064	-0.097	-0.165	-0.113
Mica	0.197	-0.075	-0.201	0.168	0.216	0.391	0.166	0.483	0.149
Mont	-0.494	0.051	0.543	0.482	0.107	0.358	0.467	0.544	0.020
Quartz	0.910**	-0.424	-0.901**	0.300	0.154	-0.211	-0.805**	-0.560	0.316
Verm	-0.339	0.266	0.305	0.048	0.160	0.474	0.445	0.683*	0.136

Mont = Montmorillinite, Verm = Vermiculite.

\*Correlation is significant at the 0.05 level (2-tailed). \*\*Correlation is significant at the 0.01 level (2-tailed)

The high CEC and relatively higher fertility status observed in soils from shale could be a reflection of the high content of high activity clays such as montmorillonite in these soils. The large CEC which is consequent of the large surface area of these minerals increase their capacity to retain and supply large quantities of nutrients such as Ca, Mg, K and  $\text{NH}_4$ , thus, these soils tend to be of high fertility, especially when proper irrigation management is practiced (Brady and Weil, 2008).

### 3.5 Mineralogical Composition of Soils Derived from Sandstone at Different Soil Depths

Soil formed in sandstone has a dominant clay mineralogy that consists of kaolinite, goethite and quartz. Okusami *et al.*, (1985) had also reported the dominance of these minerals in soils of sandstone formation. Kaolinite accounted for 26.42 to 31.26 %, goethite, 7.77 to 14.87 %, and quartz accounted for 35.4 to 51.8 %. Other minerals identified were: hermatite (4.58 to 14.62 %), gibbsite (1.70 to 5.23 %), chlorite (0.307 to 0.867 %), illite (0.13 to 0.31 %), mica (0.557 to 0.683 %) and vermiculite (0.60 to 1.79 %).

### 3.6 Mineralogical Composition of Soils Derived from Basalt at Different Soil Depths

X-rays diffraction analysis of soils formed from basalt revealed that kaolinite (14.64 to 27.44 %), quartz (23.7 to 42.5 %), hematite (11.0 to 23.14 %), goethite (13.94 to 26.48 %) were the dominant crystalline minerals present in the soil. While the quartz and kaolinite content decreased with depth, goethite and hematite increased down the profile, but gibbsite was fairly constant. Esu *et al.*, (2015) have also reported the dominance of kaolinite, quartz and iron oxides in a basaltic soil of Ikom. They stated that basaltic soils are highly weathered to the extent that they no longer possess any further weathering potential. They have indeed reached the oxic state of weathering.

### 3.7 Correlation between the Major Soil Minerals and Some Soil Properties

Correlation analysis between the major soil minerals identified and some soil properties is presented in Table 4. Sand showed negative correlation with most of the major minerals identified except quartz and mica where the



correlation was positive. The highly significant negative correlation ( $P < 0.01$ ) between sand and goethite, gibbsite and hermatite is an indication that soils with sand as dominant texture have a low or insignificant amount of these iron oxides. Similarly, the highly positive correlation between sand and quartz shows that the higher the sand contents of a soil, the more the occurrence of quartz. The reverse trend was observed when sand was correlated with goethite, gibbsite, hermatite and quartz. Therefore, soils rich in clay, have more goethite, gibbsite, montmorillonite, illite and chlorite content, but less in quartz and clay.  $P^H$  correlated negatively with goethite, hermatite and kaolinite, but positively with montmorillonite, illite, and quartz.

Significant negative correlation was observed between total exchangeable acidity and quartz, the correlation between total exchangeable acidity and goethite, hermatite, illite were positively significant. Base saturation showed negative correlation with chlorite, goethite, gibbsite, and hermatite, but positively with montmorillonite, mica, vermiculite and illite.

#### 4. CONCLUSION

The result of the present investigation on the chemistry and mineralogy of soils of contrasting parent materials at different depth suggests that the soils were generally poor in terms of fertility except the soil of shale formation where fertility indicators was relatively higher than in other soil types. Parent material and depth played dominant roles in the type and distribution of clay minerals in the study area. Mineralogy revealed the dominance of quartz, kaolinite in most of the soil studied. However a mixed mineralogy of quartz, kaolinite, goethite, hermatite and smectite was observed in soil derived from shale. Knowledge of different types of clay minerals in soil together with their distribution has greater relevance in assessing the long term nutrient supplying power of soil to crop. Also, such study will provide an important index in formulating a sound fertiliser programme for a given set of soil and crop. If suitable and efficient drainage programmes are employed, it is certain that soil of shale formation will be a potential source for plant cultivation in Southeastern Nigeria. However, for optimum and sustainable crop production in other soil types, external fertiliser input and other fertility enhancement strategies must be adopted. A more detailed study of soil minerals as they relate with the chemical, physical and biological attributes of the soils of Southeastern Nigeria will not only provide a

broad knowledge of the fertility status of these highly weathered soils, but will help farmers to formulate an effective nutrient management programmes in general for a soil and particularly for a soil type for sustainable food production.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

- Adegbenro, R.O.; Ojetade, J.O. and Amusan, A.A.(2011). Effect of topography on phosphorus forms and distribution in soils formed in schistbin lfe area. *Journal of Agriculture and Veterinary Sciences* 5(1):86-105.
- Adeoye, G. O. and Agbola, A. A. (1984).Critical level of soil pH, available P, K, Zm, and Mn and maize content of P, Cu, and Mn in the sedimentary soils of Southeastern Nigeria.*Fertilizer Research*. 6: 65-71.
- Agboola, A.A. and Corey, R.B.(1982). The relationship between soil pH, organic matter, available phosphorus, exchangeable calcium, magnesium and nine elements in the maize tissue, *Soil Science*. 115(5), 367-375.
- Akhirevbulu, O.E.; Amadasun, C.V.O.; Ogunbajo, M.I. and Ujuanbi, O.(2010). The geology and mineralogy of clay occurrences around Kutigi Cental Bida Basin, Nigeria. *Ethiopian J. Environmental Studies and Management*, 3(3):49-57.
- Aki, E.E and Antigha, N.R.B.(2015). Some physical properties of soils overlying limestone parent material in southeastern Nigeria. *Nigerian Journal of Soil Science*, 25:1-7.
- Apugo-Nwosu, T.U; Mohammed-Dabo, I.A.; Ahmed, A.S.; Abubakar, G.; Alkali, A.S. and Ayilara, S.I.(2011). Studies on the suitability of Ubakala bentonic clay for oil well drilling mud formation.*British Journal of Applied Science. Technol*.1(4):152-171.
- Barshad, I.(2006). Chemistry of soil development. In:Beer, F.E. (Editor). *Chemistry of the soil*. Raynold Pub. Corp. New York.
- Benton, Jones J. Jr. (2001). Laboratory Guide for Conducting soil test and plant analysis. CRC Press Boca Raton, Washington DC.
- Borggaard, O.K. and Elberling, B.(2004).Estimation of phosphate sorption capacity by means of a pedotransfer function. *Geoderma* 118:55-61.

- Bowman, R.A. and Savory, D.J.(1992). Phosphorus distribution in calcareous soil profiles of the Central Plains.*Soil Science Society of America Journal*, 56:423-426.
- Buol, S.W., F.D. Hole and R.J. McCracken, (1980). *Soil Genesis and Class*. Iowa State Univ. Press, USA.
- Brady, N. C. and Weil, R. R. (2008). *The Nature and Properties of Soils*. 13<sup>th</sup> Ed. Prentice Hall Inc. New Jersey, pg 201-226.
- Branhisel, R.L. and Berstesch, P.M.(1992). Chlorites and hydroxyl interlayer vermiculite and smectite . In: Dixon, J.E. and Weeds S.B.(eds). *Minerals in soil environments* , SSSA Book Ser. 1, SSA, Madison, pp 729-788.
- Brennan, F. P., Moynihan, E., Griffiths, B. S., Hillier, S., Owen, J., Pendrowski, H. and Avery L. M. (2014). Clay mineral type effect on bacterial enteropathogen survival in soil. *Sci. Total Environ.* 468,: 302-305.
- Brown, G.; Brindley, G. W. (1980) X-ray diffraction procedures for clay mineral identification. In: Brindley, G. W.; Brown, G. (Ed.). *Crystal structures of clays minerals and their X-ray identification*. London: *Mineralogical Society*,. p. 305-360.
- Carter, M. R. and Gregorich, E. G. (2008). *Soil Sampling and Methods of Analysis 2<sup>nd</sup>* (ed). Can Soc. Soil Sci. 1224p.
- Chikezie, I.A.; Eswaran, H.; Asawalam, D.O. and Ano, A.O. (2010). Characterization of two benchmark soils of contrasting parent materials in Abis State, southeastern Nigeria. *Global Journal of Applied Sciences* 16(1):23-29.
- Churchman, G. and Lowe, D.J.(2012). Alteration, formation and occurrence of minerals in soils. In: *Handbook of Soil Science: Properties and processes*. CRC Press. Pp1-72
- Dinc, U., H. Ozbek, A.S. Kapur and S. Senol, (1987). *Toprak Genesi ve Snflandrlmas* (In Turkish). Cukurova. Univ. Press, Turkey, pp: 379.
- Eneje, R. C. and Azu E. O. (2009). Algae compost effect on soil nutrient status and aggregate stability. *Nigerian Journal of Soil Science*. 19(2):118-126
- Enwezor, W.O. Ohiri, A.C, Opuwaribo, E.E and Udo, E.J. (1990). Literature Review on soil fertility investigations in Nigeria. Federal ministry of Agricultural and Natural Resources, Lagos. 460pp.
- Enwezor, W.O.; Udo, E.J.; Usoroh, N.J.; Ayotade, K.A.; Adepetu, J.A.; Chude, V.O. and Udegbe, C.I. (1989). Fertilizer use and management practices for crops in Nigeria. Series No.2. Bobma Publishers, Ibadan p. 18.
- Esu, I.E.; Uko, E. and Aki, E.E.(2015). Morphological, physicochemical and mineralogical properties of soils developed from basalt at Ikom, Cross River State, Nigeria. *Nig. J. Soil Sci.* 25:168 – 180.
- FDALR (1985). Soil Map of Nigeria project. Pp63-148.
- Grossman, R.B., Millet, J.L. (1961). Carbonate removal from soils by a modification of the acetate buffer method. *Soil Science Society of America Journal* 25, 325-326.
- Gruhn, P.; Goletti, F. and Yudelman, M.(2000). Integrated nutrient management, soil fertility and sustainable agriculture: Current issue and future challenges, food, Agriculture and the Environmental Discussion. Paper 32
- Hinsinger, P.; Bengough, A.G.; Vetterlein, D. and Young, I.M. (2009) Rhizosphere: biophysics, biogeochemistry, and ecological relevance. *Plant Soil* 321: 117–152
- Idris, O.A. Amel and Ahmed, H. Sirelkhatim (2012). Phosphorus sorption capacity as a guide for phosphorus availability of selected Sudanese soil series. *African Crop Science Journal*, 20(1):59-65.
- Igwe, C.; Zarei, M. and Stahr, K. (2015). Analysis of clay mineralogy of some southeastern soils using X-ray diffraction and fluorescence. *Geophysical Research*, 11: 622-634.
- Irmak, S., Sururu, A.K. and Aydogdu, I.H.(2007). Effects of Different Parent Material on the Mineral Characteristics of Soils in the Arid Region of Turkey. *Pakistan Journal of Biological Sciences*. 10(4):528-536
- Joint Committee on Powder Diffraction Standard (1980): *Mineral Powder Diffraction File: vols I and II*. Publ. intern. Center for Diffraction, Dam Parklane, U.S.A. pp 4-53.
- Jenny, H. (1980). *Soil Resource: Origin and Behaviour*. Springer Verlag, New York.
- Joffe, J.S., 1949. *Pedology*. Somerset Press Inc., Somerville, pp: 622.
- Jones, D.L., Pual Cross, P. and Wither, P.J.A. (2013). Nutrient stripping: the global disparity between food security and soil nutrient stocks. *Journal of Applied Ecology*. 10(11): 1365-2664.
- Jungerius, P.D. and Levelt, W.M.(1964). Clay mineralogy of soils over sedimentary rocks in Eastern Nigeria. *Soil Science*, 97:89 – 95.

- Khormali, F. and Abtahi, A.(2003). Origin and distribution of clay minerals in calcareous arid and semi-arid soils of Fars Province. *Clay Mineral.* 38:511-527.
- Konze, G. W. and Rich, C. I. (1959). Mineralogy Methods. In: C. I. rich, L. F. Satz and G. W. Kunze (eds). Certain properties of selected Southeastern United States Soils and Mineralogical Procedures for the study. Southern coop, Series Bull, Pp.135-146.
- Libo, Hao, Qiaoqiao, Wei, Yuyan, Zhao, Jilong, Lu and Xingum, Zhao (2015).Optimization method of quantitative calculation of clay minerals in soil.*J. Earth System Sci.* 124(3):675-680.
- Malgwi, W.B.A.; Ojanuga, G.; Chude, V.O.; Kparamwang, T. and Raji, B.A.(2000). Morphological and physical properties of soils of Ife and Ondo areas. *Nigerian J. Soil Sci. Research*, 1:58-64.
- Mehmood, A., Mohammad, S. A., Mohammad, I. and Shah, R. (2015). Iron oxides forms quantification in relation with soil genesis in soil parent materials. *Journal of Biodiversity and Environmental Sciences*, 6(5):383-390.
- Memon M. (2008). Role of Fe-oxides for predicting phosphorus sorption in calcareous soils. Ph.D thesis, University Karlsruhe, Germany.
- Memon M., Memon K.S., Akhtar M.S. and Stuben, D.(2009). Characterization and Quantification of Iron Oxides occurring in low concentration in soils. *Commun Soil Sci plant anal* **40**, 1625-178.
- Nadimi, M. and Farpoor, M.H.(2013). Genesis and clay mineralogy of soils on different geomorphic surfaces in Mahan-Joupar area, central Iran. *Arab. J. Geosci.* 6:825-833
- Nitzsche, R. P., Percival, J. B., Torrance, K. K., Striling, J. A. R. and Bowen, J. T. (2008). X-ray diffraction and infrared characterization of oxisols from Central and Sotheastern Brazil.*Clay Minerals*,43(4):549-560.
- Njoku, J. C., Okpara, D. A. and Asiegbo, J. A. (2006). Growth and yield response of sweet potato to inorganic nitrogen and potassium in a tropical ultisol. *Nig. Agric. J.* 32:30-40.
- Norra, S., Lanka-Panditha, M., Kramar, U., and Stuben, D.(2006). Mineralogical and geochemical patterns of Urban Surface Soils, the example of Pforzheim.*Applied Geochemistry*, 21:2064-2081.
- Nsalambi, V. Nkongola and Christopher, J. Plassmeyer (2010). Effect of vegetation type on soil physical properties of Lincoln University Living Laboratory *Research Journal of Forestry* 4:1-13.
- Nwagbara, M.O. (1997). Temperature characteristics and variability over cities in southeastern Nigeria (1965 – 1994).M.Sc thesis, Department of Geography, University of Port-Harcourt.
- Nwaogu, E. N. and Ebeuiro, C. N. (2009).Greenhouse evaluation of the performance of turmeric grown on soils of different parent materials in southeastern Nigeria. *ASN 43<sup>rd</sup> Annual Conf. Proc.* 864.
- Nweke, O.M.; Igwe, E.O. and Nnabo, P.N.(2015). Comparative evaluation of clays from Abakaliki formation with commercial bentonite clays for use as drilling mud. *Afr. J. Environmental Science and Technology*, 9(6):508-518.
- Obasi, A. I. Ejpe, I. I., Igwe, E. O. and Nnachi, Enwo, E. (2015).The physical properties of soils within major dumpsites in Abakaliki Urban, southeastern Nigeria and their importance to groundwater contamination.*International Journal of Agriculture and Forestry.* 5(1):17-22.
- Odoma, A.N.; Obaje, N.G.; Omada, J.I.; Idakwo, S.O. and Erbacher, J.(2013).Paleoclimate Reconstuction during Mamu Formation (Cretaceous) Based on Clay Mineral Distribution. *J.Appl. Geology and Geophysics* 1(5):40-46.
- Odurukwe, S. O. Aruebunwa, F. A., Iloka, A. W., Udeabor, A. and Ibedu, M. A. (1995).Physical Environment of Southeast Nigeria. In: Indigenous Fallow and Multipurpose Tree and Shrub Species in the farming system southeastern zone of Nigeria. A Report of Diagnostic Survey.NRCRI. Umudike Pubs.
- Okorie, P. E. and Okpala, I. E. (2000).Effect of animal manure and inorganic fertilization on ground flora development at two degraded sites in Umudike.*Nigerian Journal of Sustainable Agriculture and the Environment*, 2(1):84-88.
- Okunami, T. A. (1981). Land and pedogenic characterization of selected wetlands in West Africa with emphasis on rice production.Ph.D thesis, University of Minnesota, St. Paul, Minnesota, U.S.A.
- Onwuka, M.I., Osodeke, V.E. and Okolo, N.A.(2007). Amelioration of soil acidity using cocoa husk ash for maize production in Umudike area of southeastern Nigeria .*Tropical and Subtropical Agroecosystem* 7:41-45.

- Onyeogu, T., Uzoegbu, M.U. and Ideozu, R.U.(2016). Clay minerals Assessment from Maastichtian Syclinal, Afikpo, Nigeria. *International Journal of Science*.6(9):746-747.
- Orhue, Ehi Robert and John, Kingsley (2015).Phosphorus status in some soils of mid-western Agro-ecological zone of Nigeria.*Nigerian Journal of Soil Science*, 25:31-37.
- Oriajiaka, S.O.(1975). Geology In: Nigeria in maps: Eastern States. Ofofata, G.E.K.(ed). Ethiope publishing house, Benin city, Nigeria, pp.3-7.
- Osabor, V.N.; Okafor, P.C.; Ibe, K.A. and Ayi, A.A.(2009). Characterization of clays in Odukpiani, southeastern Nigeria.*African Journal of Pure and Applied Chemistry*. 3(5):79-85.
- Osodeke, V.E. and Kamalu, O.J. (1992).Phosphorus status of Hevea growing soils of Nigeria.*Indian Journal of Natural Rubber Research*, 5:107 – 112.
- Osodeke, V. E. (1996). Effects of rainfall and land-use on the management of the highly degraded soils of southeastern Nigeria.Proceedings of the 5<sup>th</sup> Annual conference of the Nigerian Society of Biological conservation, Umudike.
- Osodeke, V.E.(1999). Determination of P requirement of rubber seedlings using sorption isotherms.*Journal of sustainable Agriculture and Environment*. 2(1):139-143.
- Osodeke, V. E. and Uba, A. F. (2005).Determination of phosphorus fraction in selected soils of southeastern Nigeria.*International Journal of Natural and Applied Sciences (IJNAS)*, 1(1):10-14.
- Oti, Nnenna Nnannaya (2015). Yield decline of major crops induced by erosion on the ultisols of Owerri, southeastern Nigeria: Maize response to natural erosion. *Nigerian Journal of Soil Science*,25:8-19.
- Owliaie, H.R., Abtahi, A. and Heck, R.J.(2006). Pedogenesis and clay mineralogical investigation of soils formed on gypsiferous and calcareous materials, on a transect, Southwestern Iran. *Geoderm*. 134:62-81
- Pansu, M. and Gautheyrou, J. (2006). Handbook of Soil Analysis: Mineralogical Organic and Inorganic Methods. Springer, p995.
- Schaeffer, C. E. G. R., Fabris, J. D. and Ker, J. C. (2008). Minerals in the clay fraction of Brazilian Latosols (oxisols): A Review. *Clay Minerals*, 43(1):137-154.
- Silvaneto, L. F., Inda, A. V., Bayer, C., Dick, D. P. and Tonin, A. T. (2008). Iron oxides in tropical and subtropical Brazilian oxisols under long-term no tillage. *Soil Science Society of Brazil Journal*, 32(5):1873-1881.
- Simmone, E. N., Jones, J. B., Mills, A. H., Smittle, A. A. and Hussey, C. G. (1994). Comparison of analytical methods for nitrogen analysis in plant tissues.*Common Soil Science*, 24:1609-1616.
- Singer, A., (1989). Palygorskite and sepiolite group minerals. *Soil Sci. Soc. Am.*, 53: 829-871.
- Steel, R.G.D. and Torrie, J.H. (1980). Principles and procedures of statistics. A biometrical approach (2<sup>nd</sup> Edn). McGraw-Hill, New York, USA. pp. 20-90
- Szilas, C., Moberg, J.P.,Borgaard, O.K and Semoka, J.M.R.(2005). Mineralogy of characteristic well-drained Soils of Sub-humid to humid Tanzania. *Acta Agriculture Scandinavica B-Soil and Plant Science*, 55:241-251.
- Tang, Y. J., Jia, J. V. and Xie, X. D. (2002).Environmental significance of clay minerals. *Earth Sci. Front*, 9(2):537-344.
- Udo, E.J and Ogunwale, J.A(1977) Phosphours fraction absorption by Nigeria soils. *Soil science society of America proceeding* 36:879-883
- Udo,E. J., Ibia, T. O., Ogunwale, J. A., Ano, A. O. and Esu, I. E. (2009). Manual of Soil, Plant and Water Analysis. Sibon Books Ltd. Lagos, Nigeria. P183.
- Washer, N.E. and M.E. Collius, (1988). Genesis of adjacent morphologically distinct soils in Northwest Florida. *Soil Sci. Soc. Am. J.*, 52: 191-196.
- Wilson, M.J.(199). The origin and formation of clay minerals in soil: past, present and future perspectives. *Clay Mineral*. 34:7-25
- Wissem, H., Jean, A. M., David, P., Noura, Z. and Mongi, S. (2015). Phosphorus isotherm sorption in semi arid soil. *International Journal of Engineering and Technology*, 2(3):1995-2005.
- Yesilsoy, M. (1994). Toprak bitki ve su ilipkisi. C.U. Yay. No. 89, Adana, Turkey, pp: 144, (In Turkish).
- Yilmaz, K., (1990). Harran Ovasii Topraklarinin Mineralojik Karakteristikleri. C.U. Yay. Doktora Tezi, Adana, Turkey, pp: 186.