

Case Study

Anthrax outbreak investigation among humans and animals in northern

Ghana: Case report

Abstract

Background: Anthrax is a bacterial infection that affects both animals and humans. It is mainly caused by gram positive bacterium, *Bacillus anthracis*. It is an acute, specific infectious disease of mainly grass eating animals including cattle, sheep, goats and pigs and common in areas where livestock are raised. It affects humans when exposed to the fur, blood or carcass of an infected animal. A single case of anthrax in any given geographical area has the potential to result in an outbreak in both humans and animals if not handled effectively and in a timely manner. Anthrax disease occurrence is irregular in the northern regions of Ghana with yearly outbreaks.

Case Presentation: We report of an outbreak investigation of anthrax in both humans and animals in April, 2013. The outbreak was from Gbengo community in the Bawku West district of Upper East Region of northern Ghana. There were three human deaths due to this particular outbreak, representing a case fatality rate of 0.032 (per 1000 of a total district population 94,034). Laboratory confirmed anthrax (*Bacillus anthracis*) with a very high case fatality rate (100%) also affected domesticated herbivorous mammals (17 cattle 3 sheep, 2 goats) of all ages and sexes.

Conclusion: The case revealed that anthrax outbreaks remain a major problem in northern Ghana with high case fatality rate. It also highlights the importance of forecasting anthrax outbreak. In response, public health action to anthrax control with continued public health programming to avert future outbreaks is needed. In addition, both health systems and veterinary services need to collaborate in anthrax surveillance for early case detection and response to prevent deaths.

26 **Key Words:** Anthrax; outbreak; human, infectious disease; fatality rate, Ghana.

27 **Background**

28 Anthrax is a life threatening infectious disease that affects all warm blooded animals
29 including humans [1]. Anthrax is caused by *Bacillus anthracis* which is irregularly
30 distributed worldwide in places where repeated outbreaks occurs [2, 3]. Outbreaks of Anthrax
31 occur more frequently in developing countries like Ghana as compared to developed
32 countries [4, 5]. Rasko and colleagues observed that there are 89 known strains of anthrax
33 with the most widely recognized being the virulent Ames strain [6]. The virulent Ames strain
34 was detected in September, 2001 anthrax attacks in the United States, when some individuals
35 were sent anthrax through the mailing or postal system [6, 7].

36 *Bacillus anthracis* are gram positive bacterial spores which are soil-borne with a long life
37 span. Thus, the spores are usually present at animal burial sites of anthrax-killed animals for
38 many decades. For example, anthrax spores have been known to re-infect animals over 70
39 years after burial sites of anthrax-infected animals were disturbed [8, 9]. Anthrax spores can
40 be found on all continents including the Antarctica [6]. It mostly infects wild and
41 domesticated herbivorous mammals which ingest or inhale the spores while grazing.
42 Ingestion has been reported to be the most common route by which herbivores contract
43 anthrax [10, 11]. Carnivorous predators living in the same environment may become infected
44 themselves. Humans usually become infected with anthrax via the oral, cutaneous or
45 pulmonary routes [12, 13]. Infection in humans could occur during direct contact with
46 domesticated livestock, hunting, butchering or during food preparation and handling.
47 Exposure to blood and other tissues from anthrax infected animals via inhalation or direct
48 inoculation through broken skin has been noted to cause anthrax [13]. Direct contact with
49 anthrax spores from an animal's fur, hide or wool has also been found to cause anthrax [12,
50 13].

51 The clinical presentation or post-mortem analysis is critical stages that aid in identifying a
 52 suspected outbreak especially in low resource settings like Ghana. A suspected case of
 53 anthrax in animals occurs when the animal suffers a sudden death accompanied with one of
 54 the following signs; absence of rigor mortis (legs not stiff), heavy bleeding from the nose,
 55 mouth and other natural orifices, subcutaneous swellings, rapid bloating and dark non-
 56 clotting blood [14]. In humans, anthrax presents with an acute onset which can manifest in
 57 various forms such as cutaneous, gastro-intestinal, inhalation and meningeal [15, 16, 17]. The
 58 cutaneous manifestation of anthrax is the commonest which occurs after contact with anthrax
 59 spores [15]. This results in a development of a skin lesion which evolves over 2 to 6 days
 60 after contact with the spores from a papular through a vesicular stage. This often leads to the
 61 formation of a depressed black eschar invariably accompanied by oedema that may be mild
 62 or extensive [15, 18]. However, systemic forms are sporadic and manifest through the
 63 gastrointestinal, pulmonary or meningeal systems. Gastrointestinal manifestations occur after
 64 2 to 5 days after ingesting infected meat which is usually characterized by nausea, vomiting,
 65 anorexia and fever [19, 20]. Pulmonary (inhalation) anthrax occurs after inhaling *bacillus*
 66 *anthracis* spores which often present as a brief prodrome resembling acute viral respiratory
 67 illness, followed by rapid onset of hypoxia, dyspnoea and hyperthermia with X-ray evidence
 68 of mediastinal widening [16, 21, 22]. Additionally, it has been reported that acute onset of a
 69 high fever possibly with convulsions, loss of consciousness and other meningeal signs are
 70 consistent with meningeal inflammation [23, 24]. Anthrax disease when detected at the early
 71 stages of the infection, some strains respond well to antibiotic treatment [25, 26]. There are
 72 also effective vaccines against anthrax infection [27, 28, 29].

73

74 Over the past 12 years anthrax outbreaks have been irregular in the three northern regions of
 75 Ghana with 65% of the districts in the three regions reportedly having recorded at least one

76 outbreak of anthrax disease in livestock resulting in human deaths from 2000 to 2012 [30].
 77 Within that period, 272 cases of human anthrax had been suspected, 18 confirmed with 15
 78 deaths. In the Upper East Region (UER) of Ghana, there were anthrax outbreaks in cattle at
 79 Bongo-Soe and Bongo-Ayopia between February and March of 2013 [31, 32].
 80 The Upper East Regional Health Office was notified of the outbreak in Gbengo community
 81 through news websites. Other radio, television and news media houses in the country
 82 similarly reported of the outbreak including deaths of cattle and humans in Gbengo
 83 community. According to the report, the village has experienced sudden deaths in animals
 84 due to a strange disease unknown to them. There was no prior official report to the regional
 85 health office to these news reports; neither from the district health officials nor the veterinary
 86 services. The news item created public anxiety and debate with all attention drawn to the
 87 health authorities in the region. A team was immediately mobilized and tasked by the
 88 Regional Director of Health Services to investigate into the causes of the unusual deaths and
 89 confirm the alleged anthrax outbreak in the Bawku West district. Even though anthrax
 90 outbreak was suspected, this was not officially confirmed by any official investigation. The
 91 outbreak investigation team set out to investigate the suspected outbreak with the following
 92 specific objectives: to clinically diagnose the suspected anthrax cases in both animals and
 93 humans, to determine the extent of the possible outbreak, to identify the source of the
 94 infection, identify the populations at risk, to determine the anthrax vaccination coverage in
 95 animals and to implement control and preventive measures. It also sought to determine
 96 rainfall figures in the area for the first quarter of the year.

97 **Methods**

98 **Study Area**

99 Gbengo is a sub-district in the Bawku West district in UER (Fig.1a and 1b) of northern
 100 Ghana. It shares a common border with Burkina Faso to the north, East Mamprusi district to

the south, Bawku Municipal to the east and Nabdam district to the west. The district has a total land area of 1,070sq km with a human population of 94,034. The livestock in the district include: 26,679 cattle, 28,542 sheep, 34,167 goats and 3,567 donkeys [33]. The people of Bawku west district depend largely on livestock for both socio-cultural and economic purposes.

Study Design

The investigation adopted the One Health Approach [34]. That is, involvement of the regional health system and veterinary service in UER. The regional director of health services, regional director of agriculture and the regional veterinary officer took part in the investigation. The One Health Approach recognizes that the health of humans, animals and ecosystems are interconnected. The approach involves applying a coordinated, collaborative, multidisciplinary and cross-sectoral approach to address potential or existing risks that originate at the animal-human-ecosystems interface [34].

The investigation involved a review of past reports of outbreak investigations in UER as well as an in-depth analysis of data collected between February and April 2013 as a result of the outbreak. Two Focused Group Discussions (six in each group) were held with livestock farmers and local community members. A total of seven interviews with a cross-section of the community members from grazing sites and livestock farmers were conducted to provide an account of what happened to their animals and humans in Gbengo community for the past two months. Six interviews were also conducted with various stakeholders (i.e. district director of health, district and regional veterinary officers, and district and regional disease control officers).

Three case definition categories were used in the investigation to determine suspected, probable and confirmed cases. A case is said to be suspected when it is compatible with the clinical description and has an epidemiological link to confirmed or suspected animal cases (or contaminated animal products) from February 16, 2013 to April 18, 2013 in and around Gbengo community in Bawku West district. A probable case is a suspected case that has a positive reaction to allergic skin test (in non-vaccinated individuals) and a confirmed case was defined as a suspected case that is laboratory-confirmed. This was done by using a laboratory process called venipuncture: collection of blood from a vein located in the cubital fossa or the dorsum of the hand. The blood sample is then sent to a laboratory where serology tests are done to isolate *Bacillus anthracis*. Active case searches were also conducted in the communities based on the case definitions.

Veterinary records on anthrax vaccination coverage in the area and the history of anthrax outbreaks in the area as well as rainfall figures for the period were reviewed. Disease line list in the district hospital from February to April 2013 was also reviewed.

Laboratory investigations

The team collected sample from a dead young bull in Gbengo on the 6th April to the regional laboratory and confirmed positive for anthrax through the isolation of *Bacillus anthracis*.

Data Analysis

The results of the outbreak investigation were categorized into four thematic areas which include: Animal anthrax, Human anthrax, Veterinary records and Rainfall pattern. We performed descriptive analysis of the outbreak data by person, place and time. The findings

are presented as frequency distributions, percentages, mean $\pm$ SD, range, and rates (attack rates, case-fatality rates etc.).

Results

Animal anthrax

The total population of animals in the outbreak community - Gbengo was 214 cattle, 247 sheep 302 goats, 84 donkeys, 27 dogs and no record of pigs. Gbengo community is a Muslim community. All age groups of cattle were affected during the outbreak period with the 6-8 years age group been the most affected, 41.2% (7/17) and 3-5years age group least affected 5.9% (1/17). This is shown in Figure 3. Also, all sex groups were affected with males mostly affected 64.7% (11/17). However, there was no significant difference between mortality in males and females. This is illustrated in Figure 4. All livestock (cattle, sheep and goats) were affected in the outbreak except donkeys that were spared. Mortality in cattle was 7.9% (17/214) and that of sheep and goats were 1.2% (3/247) and 0.7% (2/302) respectively as shown in Figure 5. We could not characterize deaths in other livestock by age and sex nor develop an Epi curve because livestock farmers could only give accurate data on casualties in cattle other than sheep and goats.

The outbreak started in February with a cattle death on the 16th and 21st of February 2013. It increased to two deaths on March 2, 2013 and dropped again to one death each on the 3rd, 7th, 8th, 10th and 11th March 2013. The deaths increased to two on March 17 and dropped again to one each on the 20th, 22nd, 23rd and 25th March 2013. On April 4th and 6th 2013, there was also a death each reported. There has since been no death reported after April 6, 2013 (figure 3).

Interviews with livestock farmers revealed that their animals grazed on their farms around the community and drank at the nearby White Volta River. It was revealed that since the predominant religion in the area is Islam, dead carcasses were immediately buried since it is a taboo to eat them. Most dead animals are buried at the river bank as it was easier to excavate. They could not recall how many goats and sheep died as more value is placed on cattle than the other animals.

Human Anthrax

The investigation revealed that Gbengo community in the Bawku West district had three confirmed cases of human anthrax out of six suspected cases. There were three (3) deaths giving a case fatality rate of 50%. The anthrax mortality rate was 0.032 (per 1000 of a total district population 94,034). One (1) out of the six cases was a female. All the cases were from the Gbengo community (figure 2)

Veterinary Records

A review of veterinary records revealed that UER has had yearly outbreaks of anthrax. There was an anthrax outbreak in March, 2012 in Googo, a community which is about 12km away from Gbengo the current outbreak village. The most recently confirmed anthrax outbreak occurred in Bongo district the previous month, March 2013 in the same region. It was also observed from the records that animals in Gbengo have not been vaccinated for over 10yrs (Table 1). The region has not had any programmed vaccination for several years except ad hoc vaccinations during outbreaks targeted at the outbreak area and its surrounding.

Rainfall Pattern in the Outbreak Area, April 2013

Our investigations also revealed that there were two rainfall figures recorded during the first quarter of 2013 in the area. These were 42mm and 6mm on the 1st and 18th March, 2013 respectively. This is shown in Table 2.

Focus Group Discussion and other Stakeholder Interviews

During our FGD and stakeholder interviews, we observed that both health system and community-related challenges, mainly poor surveillance activities (notifications and reporting) by staff in the region which was largely attributed to low staff strength and limited capacity and lack of anthrax vaccination in animals due mainly to logistical problems. We also noticed that livestock farmers have over the years benefited from free vaccination services and thus shy away from fee payment for anthrax vaccination. Additionally, group and family ownership of animals poses a major problem for livestock when they are to be either vaccinated or treated for fee payment.

Discussions

Anthrax outbreak in humans in the Upper East region has always been associated with livestock infection. In recognition of the complex interplay of social, behavioural, cultural, economic, human, animal and environmental health factors in the causation and transmission of anthrax in northern Ghana, a One Health Approach methodology was used in this investigation. This is one of the most effective approaches to curb anthrax and other zoonotic diseases [34]. The number of deaths reported both in livestock and humans were an accumulation from February to April 2013. We noticed that communal and family ownership of kraals has been a challenge to the treatment and vaccination of livestock especially cattle. While most cattle owners are not able to respond to routine veterinary activities as there is inadequate veterinary staff available, the available staff lacks necessary logistics making

service delivery a challenge. From the findings, the lack of vaccination of livestock against anthrax for many years is a major risk factor for periodic outbreaks. This is particularly worrying considering that the vaccine is locally produced and should be available. This makes education of livestock farmers on vaccination an important component of any control and prevention measure in northern Ghana. Anthrax disease is more common in developing countries where there is lack of widespread veterinary or human public health programmes [35]. Unfortunately, in Ghana, there has not been a sustainable anthrax vaccination programme in animals, a situation which has left Gbengo community over ten years without vaccination coverage. This lack of vaccinations leaves a highly susceptible animal and human population in the outbreak area to anthrax infection.

The importance of educating farmers to take livestock as a business and learn to invest in it through routine treatment and vaccination especially against anthrax disease must be stressed. The poor attitude of livestock farmers to avail their animals for vaccination was evident in the Focus Group Discussions (FGDs) and is one of the factors that militated against the control of the disease. In addition, the unwillingness of cattle farmers to pay for vaccination of their livestock was also noted. Since anthrax disease affects animals and humans, it should be seen as a health development priority that requires a multi-sectorial approach, a strategy that requires the collaboration of various ministries and other international agencies. This is because elimination of anthrax in northern Ghana using the One Health Approach is only effective if there is better sectorial balance among existing groups and networks, especially between veterinarians and physicians, and increased participation of environmental and wildlife health practitioners, as well as social scientists and development actors [32, 36]. It is imperative for the strengthening of veterinary and medical services by including relevant

stakeholders in policy formulation and priorities in the decision making process towards control and prevention measures.

Consistent with previous studies by Hampson and his colleagues [37], we observed that the outbreak in Gbengo has affected domesticated herbivorous mammals (cattle sheep, goats) of all ages and sexes and has caused 100% deaths in all livestock that were known to be infected. Ingestion is thought to be the most common route by which herbivores contract anthrax [38]. Livestock in this community are free range and thus are highly exposed to anthrax spores. Though there were donkeys and dogs in the community, there were no records of their infection. There is evidence that anthrax is rare in dogs and cats too [39, 40]. However, though donkeys are susceptible to anthrax, they remain uninfected in the community because they might not have come into contact with the spores.

We observed that even though some forms of anthrax respond well to antibiotic treatment, the cases were not reported to the veterinary services or the Ghana Health Services for attention which could have prevented some deaths. Three human deaths out of six reported cases in humans were confirmed to be due to anthrax even though potentially additional deaths may have occurred outside healthcare facilities.

Outbreaks originating from soil borne infections primarily occur in the warm season. Epidemics tend to occur in association with periods of marked climatic or ecological changes such as heavy rainfall, flooding or drought [42, 43]. We observed that Gbengo recorded two rainfall figures in March. This could have caused a change in the climate causing grasses to sprout out and with animals grazing so close to the soil thereby ingesting or inhaling anthrax spores and getting the infection.

It is therefore important that community members are educated on the dangers of anthrax and its prevention. We also observed that even though there seem to be collaboration between the health services and the veterinary services, this is weak and needs to be addressed. During the outbreak, there was a ban on slaughtering and movement of livestock in the region to prevent further spread.

Study Limitations

The outbreak investigation had some limitations as a period of less than 3 months was quite short. Secondly, the refusal of some of the community members to divulge information was a setback and limitation.

Conclusion

The investigation revealed an outbreak of laboratory confirmed anthrax (*Bacillus anthracis*) with a very high case fatality rate (100% in animals and 50% in humans) in Gbengo community. There are still major challenges with anthrax outbreaks such as lack of awareness among community members, high poverty levels and irregular vaccination of livestock. Livestock vaccination rates can be improved by taking advantage of the local production of the vaccine to increase coverage. Annual outbreak of the disease and poor vaccination coverage in northern Ghana is a major problem that needs urgent attention by all stakeholders (e.g. Ghana Health Service, Ministry of Agriculture and Veterinary Service). A One Health Approach calls for collaboration and communication among stakeholders in order to utilize the synergies in anthrax control. In addition, improve anthrax surveillance and response preparedness is urgently needed to increase early case detection and reduce fatality. Increasing public information and education on the diseases etiology is important.

Consent

Verbal consent was sought from community members and livestock farmers before interviews were conducted. Permission was also granted by the Bawku west district hospital, veterinary and health departments at both district and regional levels before records were reviewed.

REFERENCES

1. Chen H, Bao W, Wang Y, Zhang K, Wang F. Clinical and epidemiological investigation of a fatal anthrax case in China. *J Infect Dev Ctries*. 2015 Feb 19;9(2):214-7.
2. Cherkasskiy, B. L. (1999). "A national register of historic and contemporary anthrax foci". *Journal of Applied Microbiology* 87 (2): 192–195. doi:10.1046/j.1365-2672.1999.00868.x. PMID 1047594
3. Booth M, Donaldson L, Cui X, Sun J et al. Confirmed *Bacillus anthracis* infection among persons who inject drugs, Scotland, 2009-2010. *Emerg Infect Dis*. 2014 Sep;20(9):1452-63. doi: 10.3201/eid2009.131481.
4. Nirmal KT, Karma W, Tshering D et al. Investigation and Control of Anthrax Outbreak at the Human–Animal Interface, Bhutan, 2010. *Emerg Infect Dis*. 2014; 20(9): 1524–1526.
5. Ohnishi N, Maruyama F, Ogawa H, Kachi H et al. Genome Sequence of a *Bacillus anthracis* Outbreak Strain from Zambia, 2011. *Genome Announc*. 2014;2(2). pii: e00116-14. doi: 10.1128/genomeA.00116-14.
6. Rasko DA, Worsham PL, Abshire TG, Stanley ST, Bannan JD, Wilson MR, Langham RJ, Decker RS, Jiang L (2011). "*Bacillus anthracis* comparative genome analysis in support of the Amerithrax investigation". *Proc Natl Acad Sci U S A* 108 (12): 5027–32. doi:10.1073/pnas.1016657108. PMC 3064363. PMID 21383169.
7. Goel AK. Anthrax: A disease of biowarfare and public health importance. *World J Clin Cases*. 2015 Jan 16;3(1):20-33.
8. "Crossrail work stopped after human bones found on site". *London Evening Standard*. 2009. [Online]. Available from: <http://www.standard.co.uk/news/crossrail-work-stopped-after-human-bones-found-on-site-6759649.html> [Accessed March 9, 2015]
9. Guillemin J. Anthrax: The Investigation of a Deadly Outbreak. *N Engl J Med* 2000; 342:1373 May 4, 2000 DOI: 10.1056/NEJM200005043421815


10. Dragon, DC, Baxer DE, Mitchell J, Wollen N, (2005) Natural Dissemination of *Bacillus Anthracis* Spores in Northern Canada. *Applied and Environmental Microbiology* 71: 1610-1015
11. Hudson JA, Daniel RM, Morgan HW (2006). "Acidophilic and thermophilic *Bacillus* strains from geothermally heated antarctic soil". *FEMS Microbiol Lett* 60 (3): 279–282.
12. Bayindir Y, Bayraktar M, Karadag N, Ozcan H, Kayabas U, Otlu B, Durmaz R, Doganay M. Investigation and analysis of a human orf outbreak among people living on the same farm. *New Microbiol.* 2011 Jan;34(1):37-43.
13. Frankel AE, Kuo SR, Dostal D, Watson L, Duesbery NS, Cheng CP, Cheng HJ, Leppla SH. Pathophysiology of anthrax. *Front Biosci (Landmark Ed)*. 2009 Jan 1;14:4516-24.
14. Saskatchewan Ministry of Health. Anthrax in Animals - Human Precautions. 2010. [Online]. Available from: <http://www.health.gov.sk.ca/adx/asp/adxGetMedia.aspx?DocID=e7c86ba2-1a15-44a0-ac3f-3397c2e8c0c1&MediaID=4474&Filename=Anthrax+in+Animals+-+Human+Precautions.pdf&l=English> [Accessed March 9, 2015].
15. Recep T, Bilal S, Ozcan D, Alicem T et al. Cutaneous anthrax in Southeast Anatolia of Turkey 2014. (doi:10.3109/15569527.2014.880844)
16. Coggeshall KM, Lupu F, Ballard J, Metcalf JP et al. The sepsis model: an emerging hypothesis for the lethality of inhalation anthrax. *J Cell Mol Med*. 2013 Jul;17(7):914-20. doi: 10.1111/jcmm.12075.
17. Hatami H, Ramazankhani A, Mansoori F. Two cases of gastrointestinal anthrax with an unusual presentation from Kermanshah (western Iran). *Arch Iran Med*. **2010**;13(2):156-9.
18. Nandi AK, Kamal MM, Alam MA, Rahman F, Uddin MJ, Baidya NR, Mostafa SM. Cutaneous anthrax in a school teacher. *Mymensingh Med J*. 2014;23(2):372-4.
19. Owen JL, Yang T, Mohamadzadeh M. New insights into gastrointestinal anthrax infection. *Trends Mol Med*. 2015;21(3):154-163.
20. Ghodratollah M, Abbas A, Mehrdad K. Gastrointestinal anthrax: clinical experience in 5 cases. *Caspian J Intern Med*. 2013 Spring; 4(2): 672–676.
21. Sykes A, Brooks T, Dusmet M, Nicholson AG, Hansell DM, Wilson R. Inhalational anthrax in a vaccinated soldier. *Eur Respir J*. 2013;42(1):285-7.
22. Twenhafel NA. Pathology of inhalational anthrax animal models. *Vet Pathol*. 2010;47(5):819-30.

23. Khoddami M, Shirvani F, Esmaeili J, Beladimogaddam N. Two rare presentations of fatal anthrax: meningeal and intestinal. *Arch Iran Med.* **2010**;13(5):432-5. doi: 010135/AIM.0013.
24. Albayrak F, Memikoglu O, Kurt O, Cokca F, Birengel S, Tekeli E. A case of anthrax meningitis. *Scand J Infect Dis.* 2002;34(8):627-8.
25. Mikesell P, Ivins BE, Ristoph JD, Vodkin, MH, Dreier TM, Leppla SH (1983-1123456). "Plasmids, Pasteur, and Anthrax" (PDF). *ASM News* 49: 320–2.
26. Migone TS, Bolmer S, Zhong J, Corey A, Vasconcelos D, Buccellato M, Meister G. Added benefit of raxibacumab to antibiotic treatment of inhalational anthrax. *Antimicrob Agents Chemother.* 2015;59(2):1145-51.
27. Ding G, Chen X, Zhu J, Duesbery NS, Cheng X, Cao B. A human/murine chimeric fab antibody neutralizes anthrax lethal toxin in vitro. . *Clin Dev Immunol.* 2013;2013:475809.
28. Hassett KJ, Vance DJ, Jain NK, Sahni N, Rabia LA et al. Glassy-state stabilization of a dominant negative inhibitor anthrax vaccine containing aluminum hydroxide and glycopyranoside lipid a adjuvants. *J Pharm Sci.* 2015 Feb;104(2):627-39.
29. Wang HC, An HJ, Yu YZ, Xu Q. Potentiation of anthrax vaccines using protective antigen-expressing viral replicon vectors. *Immunol Lett.* 2015;163(2):206-13.
30. Veterinary Services Directorate Annual Report, 2012.
31. Veterinary Services Directorate First Quarter Report, 2013.
32. Opare C, Nsiire A, Awumbilla B , Akanmori BD. Human behavioural factors implicated in outbreaks of human anthrax in the Tamale municipality of northern Ghana
33. Ghana Statistical Service. 2010 Population and housing census. 2012 [Online]. Available from: http://www.statsghana.gov.gh/docfiles/2010phc/Census2010_Summary_report_of_final_results.pdf [Accessed March 3, 2015].
34. Mbabu M, Njeru I, File S, Osoro E, Kiambi S et al. Establishing a One Health office in Kenya. . *Pan Afr Med J.* 2014;19:106. doi: 10.11604/pamj.2014.19.106.4588.
35. Murthy S, Keystone J, Kissoon N. Infections of the developing world. *Crit Care Clin.* 2013;29(3):485-507.
36. Patocka J, Splino M. Anthrax toxin characterization. *Acta Medica (Hradec Kralove).* 2002;45(1):3-5.

- 393 37. Hampson K, Lamb T, Auty H, et al (2011) Predictability of Anthrax Infection in the
394 Serengeti, Tanzania. J Appl Ecol 48(6): 1333-1344
- 395 38. Dragon DC, Elkin BT, (2011) An Overview of Early Anthrax Outbreaks in Northern
396 Canada: Field Reports of the Health of Animals Branch, Agriculture, Canada. Arctic Vol. 54.
397 No. 1 pp 32-40
- 398 39. Langston C. Postexposure management and treatment of anthrax in dogs--executive
399 councils of the American Academy of Veterinary Pharmacology and Therapeutics and the
400 American College of Veterinary Clinical Pharmacology. AAPS J. 2005;7(2):E272-3.
- 401 40. John GB, Thomas VI, Luciana B. Management of Anthrax. *Clinical Infectious Diseases*.
402 2001;35(7): 851-858
403
- 404 41. Blackburn JK, Asher V, Stokke S, Hunter DL, Alexander KA. Dances with anthrax:
405 wolves (*Canis lupus*) kill anthrax bacteremic plains bison (*Bison bison bison*) in southwestern
406 Montana. J Wildl Dis. 2014;50(2):393-6.
- 407 42. World Health Organisation (2008) Anthrax in Humans and Animals. Fourth Edition,
408 WHO Library Cataloguing Publication Data.
- 409 43. Hugh-Jones M, Blackburn J. The Ecology of *Bacillus Anthracis*. Molecular Aspects of
410 Medicine. 2009; 3(356-367)

411

412 **Figure 1a & 1b:**



413




414

415

416 **Figure 2: Epi curve**

417




418

419

420

421 **Figure 3:**

422




423

424

425 **Figure 4:**

426

Number of Cattle Deaths by Sex in Gbengo, Bawku West District
(9th February - 18th April 2013)



427


428

429

430

431 **Figure 5:**

Number of Deaths by species in Gbengo between
9th February and 18th April 2013



432

433 **Table 1: Vaccination coverage of Animals in Bawku West District 2000- April 2013**

Anthrax Vaccination 2000 – 2013: Bawku West District, April 2013						
Year	Cattle	Sheep	Goats	Pigs	Dogs	Donkeys
2000	x	x	x	X	X	X
2001	x	x	x	X	X	X
2002	x	x	x	X	X	X
2003	x	x	x	X	X	X
2004	x	x	x	X	X	X
2005	x	x	x	X	X	X
2006	x	x	x	X	X	X
2007	x	x	x	X	X	X
2008	185	x	x	X	X	X
2009	x	x	x	X	X	X
2010	x	x	x	X	X	X
2011	x	x	x	X	X	X
2012	3,752	513	86	47	96	16
2013	1,240	307	161	0	65	28

434 X= No vaccinations carried out in the animal species in the corresponding year

435 Source: Veterinary Service Department, Upper East Region

436 **Table 2: Rainfall Recorded during the 1st Quarter of 2013**

Month	1st Week	2nd Week	3rd Week	4th Week
January	0	0	0	0
February	0	0	0	0

March	42mm	0	6mm	0
April	0	0	0	0

437 Source: Meteorological Service Department

438

439