

Original Research Article**Awareness and Status of Tetanus Toxoid
Immunization among Female Undergraduate
Students in a Nigerian University.****ABSTRACT**

Aims: The study aimed to determine the level of awareness and status of tetanus toxoid (TT) immunization among female undergraduates in Port Harcourt.

Study design: This was a cross sectional school based study carried out among female undergraduate students.

Place and Duration of Study: The study was carried out in University of Port Harcourt in May 2014.

Methodology: A pre-tested questionnaire was administered to 512 female students selected through a Multi-Staged sampling technique. Information sought included socio-demographics, awareness of TT vaccine, knowledge of the immunization campaign for women of childbearing age and TT immunization status. Data was analyzed using epi info version 7.1.3.3.

Results: A total of 512 students participated in the study; mean age was 22 ± 2.79 years. Four hundred and eighty five (94.7%) and 409 (79.9%) students had heard of tetanus and TT vaccine respectively. Commonest sources of information were health workers. 33.8% knew the target population for the TT immunization campaign and 18.2% knew that the correct number of doses of TT is 5 doses. Two hundred and ninety eight (58.2%) and 10 (2%) students had received at least 1 dose and 5 doses of TT respectively. Students who had knowledge of the target population for TT immunization campaign (OR=2.1362, $p=0.003$) and those who were confident in the effectiveness of TT (OR=2.4074, $p=0.000$) were twice more likely to receive at least a dose of TT vaccine.

Conclusion: There was poor knowledge of TT immunization campaign for women of childbearing age and very poor coverage of TT5 among the female undergraduate students in Port Harcourt. Female undergraduate students should be reached during TT immunization campaigns.

Keywords: [Tetanus toxoid, Female undergraduates, Students, Port Harcourt]

18 **1. INTRODUCTION**

19 Maternal and neonatal tetanus (MNT) is a public health problem which has continued to
20 plague most countries in the developing world, despite the availability of an effective
21 preventive vaccine [1]. Tetanus is an acute, infectious, non-communicable disease caused
22 by *Clostridium tetani* [1]. Every year 59,000 newborns die of this disease: this translates into
23 164 deaths daily or one death every 9 minutes of children under 1 month of age [1,2]. It
24 accounts for 7% of neonatal deaths globally [3]. Thirty thousand mothers also die from
25 tetanus after child birth annually, translating to 5% of maternal deaths [4]. Most of these
26 deaths occur in Africa and in East and Southern Asia among populations with limited or no
27 access to health care services and information related to safe birthing practices, with poor
28 socio-economic status and low literacy rate.

29 Many countries have been certified as maternal and neonatal tetanus free, due to global
30 efforts which have been directed at eliminating the disease [5]. Nigeria is one of the
31 remaining 16 countries in Africa that are yet to eliminate maternal and neonatal tetanus [5].
32 In 2000, Nigeria was among the 27 countries said to be responsible for 90% of the global
33 neonatal tetanus cases [4]. In the same year, the World Health Organization (WHO)
34 classified Nigeria among the 17 countries with more than 50% of their districts at risk for
35 maternal and neonatal tetanus because of their limited infrastructure (indicated by DPT3
36 immunization rate of 50% or less), serious manpower and logistic constraints [4]. In Nigeria,
37 Neonatal tetanus accounts for 20% of the neonatal deaths, with the burden of the disease
38 worst in the North East and North West parts of the country, where it accounts for up to
39 68.1% and 74.2% of the neonatal deaths in these regions respectively [1]. The major
40 predisposition to this disease is lack of maternal immunization with Tetanus Toxoid vaccine,
41 coupled with deliveries conducted by untrained caregivers in an unhygienic environment and
42 with unhygienic birth practices [1,4]. These conditions allow the tetanus spores to
43 contaminate maternal wounds during child birth and the umbilical cord at the time it is cut or
44 dressed after delivery [1,4].

The major means of preventing maternal and neonatal tetanus recommended by WHO is the immunization of women of childbearing age (15 to 49 years) with 5 doses of tetanus toxoid vaccine as early as possible [1,4]. The vaccine schedule is as follows: First dose (TT1) is to be administered at first contact, 2nd dose (TT2) at 4 weeks after the first dose, TT3 6 months after TT2, TT4 1 year after TT3 and TT5 is given 1 year after TT4 [6]. The complete 5 doses is expected to provide lifelong protection for the would be mother and also provide adequate partial immunity for her unborn baby against neonatal tetanus [6]. Data on the up take of the vaccine by women of childbearing age in Nigeria showed poor up take of the vaccine [7-10]. In Ibadan, Bamidele and Umoh [11] reported that 36.3% of female undergraduates were aware of the tetanus toxoid immunization schedule, though 35.8% had commenced the schedule, only 10.4% of their respondents had received TT2. Lack of information on the need for the vaccination schedule was the commonest reason for non-vaccination. The authors concluded that there was a wide gap between policy formation and practice [10]. For maternal and neonatal tetanus to be eliminated in Nigeria, this gap has to be bridged. Data on tetanus toxoid immunization status of female undergraduates in Port Harcourt are scarce, this present study therefore aimed to determine the level of awareness and status of tetanus toxoid immunization among female undergraduates in Port Harcourt.

2. MATERIAL AND METHODS

This was a school based cross-sectional descriptive study carried out among female undergraduate students of the University of Port Harcourt in May 2014. The University of Port Harcourt is the largest Federal tertiary institution in Rivers State, Southern Nigeria. Records show that as at the 2013 / 2014 session, there were 11,526 male and 9,674 female regular undergraduate students.

The minimum sample size for this study was calculated using the formula [12]

$$n = \frac{Z^2(pq)}{d^2}$$

75 E^2
76 Where n = minimum sample size

77 p= Prevalence of 35.8% reported by Bamidele and Umoh [10]

78 q= 100-p

79 E= Margin of error tolerated at 5%

80 Z= 1.96 at 95% confidence level

81 $n = \frac{1.96^2 \times 35.8 \times (100-35.8)}{5^2} = 353$
82

83 Thus minimum sample size of 353 was calculated.

84 Allowance for Design Effect (DE) = Sample size $\times 1\frac{1}{2}$ = 529.5

85 Allowance for attrition (5%) =26.5

86 Minimum sample size =529.5 + 26.5 = 556

87 However  60 students were recruited for the study. A multi – staged sampling technique
88 was used to recruit the students. The students were first stratified into the eight major
89 Faculties and Colleges of the University. One department was randomly chosen from each
90 of the eight faculties / colleges. From the chosen departments, two academic grade levels
91 were chosen by simple random sampling technique. The first 35 females in the register in
92 the chosen levels were recruited for the study, making a total of 70 females per department.
93 However 12 (2.1%) students did not give consent, another 36 (6.4%) did not return their
94 questionnaires and were excluded from the study. Eventually only 512 students participated
95 in the study.

96 The questionnaire used for the study was structured; self-administered and pretested by
97 administering it to 40 students in a department not included in the study and the pretest
98 results were also not included in the study. Information sought included socio-
99 demographics, awareness of tetanus and tetanus toxoid (TT) and immunization status. Data
100 was analyzed using the Epi Info version 7.1.3.3 statistical software. Proportions were
101 compared using the chi square test and a p value of less than 0.05 was regarded as
102 significant.

103

104 3. RESULTS

105

106 A total of 512 female students participated in the study. The age of the respondents ranged
107 from 15 to 31 years with a mean age of 22 ± 2.79 years. A total of 405 (79.1%) of the
108 respondents were less than 25 years of age, while 107 (20.9%) were 25 years and above.
109 Of the 512 respondents, 485 (94.7%) had heard about tetanus while 27 (5.3%) had not. Up
110 to 172 (33.6%) respondents heard about tetanus from a health worker, 150 (30.9%) from
111 school lessons, 67 (13.8%) heard about tetanus from the media, 40 (8.2%) from the media
112 and family members and 30 (6.2%) heard about tetanus from family members alone. A total
113 of 409 (79.9%) of the respondents had heard about tetanus toxoid while 103 (20.1%) had
114 not. Concerning the correct response as to the target population for tetanus toxoid
115 immunization campaigns, 173 (33.8%) of the respondents correctly answered "women of
116 childbearing age". Sixty four (12.5%) answered "pregnant women only", 15 answered
117 "married women only" and 260 (50.8%) had no idea.
118 Table 1 shows the relationship between the knowledge of the target population for TT and
119 some variables. Correct knowledge of the target population for TT was significantly higher
120 among the older age group ($P = 0.013$) and those in Medical faculties ($P = 0.001$). A
121 significantly increasing trend in the proportion of respondents with correct knowledge of the
122 TT target population was also noticed with advancing year of study ($P = 0.001$).
123 .

124 **Table 1: Relationship between knowledge of tetanus toxoid target population and**
125 **some variables**

Knowledge of Tetanus Toxoid target population				
Variable	Correct N (%)	Wrong N (%)	Total N	P value

Age group	<25 years	126 (31.1)	279 (68.9)	405	0.013*
	≥25 years	47 (43.9)	60 (56.1)	107	
Year of study	One	39 (26.5)	108 (73.5)	147	<0.001*
	Two	45 (25.3)	133 (74.7)	178	
	Three	14 (29.8)	33 (70.2)	47	
	Four	31 (33.3)	62 (66.7)	93	
	Five	3 (100.0)	0 (0.0)	3	
	Six	41 (93.2)	3 (6.8)	44	
College / Faculty	Medical	46 (71.9)	18 (28.1)	64	<0.001*
	Non - medical	127 (28.4)	321 (71.6)	448	

126 * = significant

127 Only 93 (18.2%) of the respondents knew that the complete number of shots for TT was 5
 128 shots. Other respondents gave various answers from 1 – 4 and 319 (62.3%) had no idea
 129 Table 2 shows the relationship between the knowledge of the complete dose of TT and
 130 some variables. Correct knowledge of the complete dose of TT was significantly higher
 131 among the older age group ($P = 0.001$) and those in Medical faculties ($P=0.001$).

132 **Table 2: Relationship between knowledge of tetanus toxoid complete dose and some**
 133 **variables**

Knowledge of Tetanus Toxoid complete dose					
Variable		Correct N (%)	Wrong N (%)	Total N	P value
Age group	<25 years	53 (13.1)	352 (86.9)	405	<0.001*
	≥25 years	40 (37.4)	67 (62.6)	107	
Year of study	One	14 (9.5)	133 (90.5)	147	<0.001*
	Two	24 (13.5)	154 (86.5)	178	
	Three	5 (10.6)	42 (89.4)	47	
	Four	9 (9.7)	84 (90.3)	93	
	Five	1 (33.3)	2 (66.7)	3	
	Six	40 (90.0)	4 (9.1)	44	
College / Faculty	Medical	39 (60.9)	25 (39.1)	64	<0.001*
	Non - medical	54 (12.0)	394 (88.0)	448	

134 * = significant

135 Concerning the effectiveness of TT, 309 (60.4%) thought it was effective, 32 (6.2%) thought
136 it was not and 171 (33.4%) had no idea.

137 A total of 298 (58.2%) of the respondents had received at least one dose of TT and 214
138 (41.8%) had not received any at all. The most common occasion on which TT was received
139 was after an injury (57.7%), routine immunization (32.6%) and during antenatal care
140 (10.1%).

141 Figure 1 shows the reasons for not receiving any dose of TT. Most of the respondents who
142 did not receive any dose of TT had no reason for not doing so. The next popular reason
143 seen among 59 of the respondents was not being aware.




Figure 1: Reasons for not receiving tetanus toxoid immunization.

Table 3 shows the relationship between having received at least a dose of TT immunization and some variables. Being in a medical department ($p < 0.001$), being confident in the

effectiveness of TT ($P= 0.001$), a higher year of study ($P= 0.001$), correct knowledge of TT target population ($P=0.001$) and correct knowledge of the complete dose of TT ($P=0.001$) were significantly associated with receiving at least a dose of TT.

Table 3: Relationship between receiving tetanus toxoid immunization and some variables

		Received Tetanus Toxoid immunization			P value
Variable		Yes N (%)	No N (%)	Total N	
Age group	<25 years	231 (57.0)	174 (43.0)	405	0.298
	≥25 years	67 (62.6)	40 (37.4)	107	
Year of study	One	72 (49.0)	75 (51.0)	147	<0.001*
	Two	100 (56.2)	78 (43.8)	178	
	Three	28 (59.6)	19 (40.4)	47	
	Four	55 (59.1)	38 (40.9)	93	
	Five	3 (100.0)	0 (0.0)	3	
	Six	40 (90.9)	4 (9.1)	44	
College / Faculty	Medical	52 (81.2)	12 (18.8)	64	<0.001*
	Non - medical	246 (54.9)	202 (45.1)	448	
Confidence in effectiveness of TT	Yes	215 (69.6)	94 (30.4)	309	< 0.001*
	No	83 (40.9)	120 (59.1)	203	
Knowledge of tetanus toxoid target population	Correct	133 (76.9)	40 (23.1)	173	<0.001*
	Wrong	165 (48.7)	174 (51.3)	339	

Knowledge of tetanus toxoid complete dose	Correct	72 (77.4)	21 (22.6)	93	<0.001*
	Wrong	226 (53.9)	193 (46.1)	419	

* = significant

Out of the 298 respondents that received at least a dose of TT, 10 (3.4%) had received 5 complete doses (2.0% of the total number of respondents), 15 (5%) had received 4 doses, 41 (13.8%) had received 3 doses, 85 (28.5%) had received 2 doses and 84 (28.2%) had received only TT1.

Table 4 shows the relationship between receiving the complete dose of TT and some variables. The older age group ($p = 0.022$), being confident in the effectiveness of TT ($P = 0.047$), correct knowledge of TT target population ($P=0.001$) and correct knowledge of the complete dose of TT ($P=0.001$) were significantly associated with receiving the complete dose of TT.

Table 4: Relationship between receiving complete dose of tetanus toxoid immunization and some variables

Received complete dose of Tetanus Toxoid immunization					
Variable		Yes N (%)	No N (%)	Total N	P value
Age group	<25 years	5 (1.2)	400 (98.8)	405	0.022*
	≥25 years	5 (4.7)	102 (95.3)	107	
Year of study	One	3 (2.0)	144 (98.0)	147	0.371
	Two	3 (1.7)	175 (98.3)	178	
	Three	1 (2.1)	46 (97.9)	47	
	Four	0 (0.0)	93 (100.0)	93	

	Five	0 (0.0)	3 (100.0)	3	
	Six	3 (6.8)	41 (93.2)	44	
College / Faculty	Medical	3 (4.7)	61 (95.3)	64	0.118
	Non - medical	7 (1.6)	441 (98.4)	448	
Confidence in effectiveness of TT	Yes	9 (2.9)	300 (97.1)	309	0.047*
	No	1 (0.5)	202 (99.5)	203	
Knowledge of tetanus toxoid target population	Correct	9 (5.2)	164 (94.8)	173	<0.001*
	Wrong	1 (0.3)	338 (99.7)	339	
Knowledge of tetanus toxoid complete dose	Correct	7 (7.5)	86 (92.5)	93	<0.001*
	Wrong	3 (0.7)	416 (99.3)	419	

167 * = significant

168 Table 5 shows the logistic regression analysis of some variables relating to receiving tetanus
 169 toxoid immunization by the respondents. After adjusting for various factors, only confidence
 170 in effectiveness of tetanus toxoid vaccine (OR=2.4074,P=0.000, Coefficient=0.8785) and
 171 the correct knowledge of the target population for tetanus toxoid (OR=2.1362,P = 0.003,
 172 Coefficient=0.759) significantly influenced receiving at least a dose of tetanus toxoid among
 173 the respondents. No factor was found to be significantly associated with the completion of
 174 the 5 doses of TT vaccine.

175 **Table 5: Logistic regression analysis of factors relating to receiving Tetanus Toxoid**
 176 **immunization**

Variable	Received at least a dose of TT immunization			Received complete dose of TT immunization		
	Odds Ratio	Coefficient	P-Value	Odds Ratio	Coefficient	P-Value
Age group (<25 / ≥25 years)	0.8543	-0.1575	0.549	2.4585	0.8996	0.2272
Faculty (Non medical / Medical)	0.9538	-0.0473	0.9348	1.7767	0.5748	0.4827
Confidence in effectiveness of tetanus toxoid (Yes / No)	2.4074	0.8785	0.000*	1.6147	0.4792	0.6757
Knowledge of tetanus toxoid target population (Correct / Wrong)	2.1362	0.759	0.0033*	8.4928	2.1392	0.0764
Knowledge of tetanus toxoid complete dose	1.2547	0.2269	0.5243	0.3082	-1.1771	0.1636

177 * = significant

178

179 **4.DISCUSSION**

180 Our data revealed that all the respondents in this study were within the childbearing age, the
 181 population of women recommended by WHO to receive 5 doses of tetanus toxoid
 182 immunization for the prevention of maternal and neonatal tetanus [4]. It also showed that

183 more than three quarter of them were aware of tetanus and tetanus toxoid vaccine. This high
184 level of awareness is comparable to what was reported by Qadir et al among female college
185 and University students in Karachi in Pakistan [13] and by Bamidele and Umoh among
186 female workers and undergraduates in a university and teaching hospital in Nigeria [11]. The
187 high educational qualification of the respondents in these studies may account for the high
188 level of awareness reported in these studies. The fact that the Commonest source of
189 information was health workers, followed by classroom lessons, demonstrates the important
190 role health facilities and schools can play in dissemination of health information to the
191 general public.

192 The results however indicate that this high level of awareness is not commensurate with the
193 level of Knowledge exhibited by the respondents. Only one third of the respondents knew
194 that the tetanus toxoid immunization campaign is targeted at women of childbearing age.
195 More than half of the respondents did not know the target population, while 15.4% of the
196 respondents were under the misconception that it is targeted at married women only or
197 pregnant women only. Similarly, a very small proportion (18.2%) of the respondents knew
198 that the complete dose for tetanus toxoid is 5 doses. This poor knowledge of the tetanus
199 immunization programme and the high level of misconception about the target population
200 may have contributed to the poor up-take of the vaccine by women of childbearing age in
201 Nigeria, as has been previously reported [7-10].

202 In this study, correct knowledge of the target population and the correct number of doses of
203 tetanus toxoid to be administered are positively associated with being older in age, higher
204 academic level and being in the Faculty of Medicine. This is not surprising because it shows
205 the positive relationship between higher educational attainment and general knowledge
206 about disease prevention. Students in Faculty of Medicine may have received detailed
207 information about the tetanus toxoid immunization programme in course of their academic
208 pursuits and were therefore operating at a higher knowledge level compared to the students
209 from other faculties. Bamidele and Umoh [11] and Basher [14] also reported similar findings

210 among medical students in Nigeria and India respectively. The study further showed that
211 majority (60.4%) of the students thought that TT is effective in preventing tetanus. This
212 confirms the report of Naeem et al who made similar observations [15], though in the
213 previous study, a higher proportion (89.5%) of respondents thought TT was effective.

214

215 The United Nations International Children's Fund (UNICEF), World Health Organization
216 (WHO) and United Nations Population Fund (UNFPA) recommended that in high risk
217 countries such as Nigeria, 90% of women of childbearing age should receive TT to achieve
218 and maintain elimination of maternal and neonatal tetanus [4]. In this study, only 58.2% of
219 the respondents had received at least one dose of TT, showing that Nigeria still has a long
220 way to go in this fight to eliminate maternal and neonatal tetanus. This result however
221 compares favourably with the 3.1% and 35.8%, 39.8% and 55.6% and 52% reported by
222 Orimadegun [9], Bamidele and Umoh[10] in South Western Nigeria, Qadir et al [13] and
223 Naeem et al[15] in Pakistan and Perry et al[16] in Bangladesh respectively. It is however
224 lower than the 70.9% reported among medical students in Mymensingh, India by Basher
225 [14]. These differences in observation maybe due to differences in level of awareness and
226 knowledge of tetanus toxoid immunization programme for women of childbearing age.

227

228 The study further indicate that the most common occasion for receiving TT among the
229 respondents was following an injury. Routine immunization was the second most common
230 reason for receiving TT. Orimadegun et al [9] also had similar observation in Western
231 Nigeria, but Bamidele and Umoh [11] reported routine TT immunization during pregnancy as
232 the most common reason for receiving TT by majority of their respondents. This difference in
233 observation maybe a reflection of differences in the age groups of respondents used in the
234 three studies. The Bamidele and Umoh's study involved older women who may have been
235 exposed to TT immunization during their several ante natal visits in health facilities.

236 The results in this study also showed that majority of the respondents who did not receive TT
237 had no reason for not doing so; the next commonest reason was lack of awareness. This
238 shows that with adequate community awareness campaign aimed at increasing the
239 knowledge of women of childbearing age about the danger of maternal and neonatal tetanus
240 and the need to prevent the disease by receiving 5 doses of TT immunization, the rate of
241 vaccination may improve and Nigeria may meet the 90% target needed for the elimination of
242 maternal and neonatal tetanus [4].

243 The study further revealed the very poor status of TT5 immunization coverage among the
244 respondents. Only 3.4% of the respondents who received at least one dose of TT received
245 the complete 5 doses, i.e.2% of all the respondents. This is not unexpected because the
246 poor knowledge of the respondents on TT immunization programme for women of
247 childbearing age, also affected their practice. Studies from other developing countries also
248 reported a very poor TT5 immunization coverage among women of childbearing age
249 [13,14,16].

250 Bivariate analysis showed that being in the medical faculty, being older in age, higher year of
251 study, being confident in the effectiveness of TT, having correct knowledge of the target
252 population and the complete dose of TT vaccine were positively associated with receiving TT
253 immunization. However, after multivariate logistic regression analysis, only being confident in
254 the effectiveness of TT vaccine in preventing tetanus and the knowledge of the target
255 population of TT immunization campaign significantly influenced receiving at least one dose
256 of the vaccine. This highlights the important role health education in schools, health facilities
257 and the mass media can play in boosting TT immunization coverage by women of
258 childbearing age.

259 The first limitation of the study is that the information sought is based on self-report and is
260 subject to over or under reporting. Secondly immunization received during childhood and the
261 adolescent period may not be easily remembered in adulthood.

262

263 **5. Conclusion**

264 There was a high level of awareness about tetanus and tetanus toxoid vaccine but poor
 265 knowledge of the tetanus toxoid immunization programme for women of child bearing age
 266 among female undergraduate students in Port Harcourt. There was also very poor coverage
 267 of TT5 among the students. Students who had knowledge of the target population of the TT
 268 immunization campaign and those who had confidence in the effectiveness of TT vaccine in
 269 preventing tetanus were twice more likely to receive at least one dose of TT vaccine
 270 compared to their counterparts who lacked such knowledge and also lacked confidence in
 271 the effectiveness of the vaccine. .

272 Extensive awareness campaign should be carried out in the university campuses on the
 273 danger of maternal and neonatal tetanus, the availability of a safe and effective preventive
 274 vaccine and the immunization schedule for women of childbearing age. Women of
 275 childbearing age in the University campuses should be reached during the sub immunization
 276 campaigns.

277

278

279

280 **ETHICAL APPROVAL**

281 Ethical approval was obtained from the Ethics Committee of the University of Port Harcourt
 282 Teaching Hospital.

283

284

285

286 **REFERENCES**

287

- 288 1. UNICEF. Maternal and Neonatal Tetanus (MNT) Elimination Initiative. Pampers
 289 UNICEF 2010 campaign launch. Why eliminate MNT? Make an invisible killer
 290 visible. Available at
 291 [http://www.unicef.org/corporate_partners/files/APPROVED_MNT_Report_05.06.10.](http://www.unicef.org/corporate_partners/files/APPROVED_MNT_Report_05.06.10.pdf)
 292 [pdf](http://www.unicef.org/corporate_partners/files/APPROVED_MNT_Report_05.06.10.pdf) . Accessed 11th August 2014.

- 293 2. Black RE, Cousens S, Johnson HL, Lawn JE, Rudan I et al. Global, regional and
294 national causes of child mortality in 2008: a systematic analysis. Lancet 2010; 375
295 (9730): 1969-1987.DOI: 10.1016/S0140-6736(10)60549-1.
- 296 3. Bryce J, Boschi-Pinto C, Shibuya K, Black RE. WHO estimates of the causes of
297 death in children. Lancet 2005; 365 (9465): 1147-1152. DOI: 10.1016/S0140-
298 6736(05)71877-8.
- 299 4. UNICEF/WHO/UNIFPA. Maternal and neonatal tetanus elimination by 2005.
300 Strategies for achieving and maintaining elimination. Available at
301 http://www.unicef.org/health/files/MNTE_strategy_paper.pdf . Accessed 8th August
302 2014.
- 303 5. Kiwanis/UNICEF. ELIMINATE maternal/neonatal tetanus. Maternal and neonatal
304 tetanus elimination announcements. Available at
305 sites.kiwanis.org/Kiwanis/en/theEliminateproject/Progress/Maternal_and_neonatal_t
306 etanus_elimination_updates.aspx. Accessed 5th August 2014.
- 307 6. WHO.Standards for maternal and neonatal care. Maternal immunization against
308 tetanus. Integrated Management of Pregnancy and ChildBirth (IMPAC). Geneva.
309 WHO 2006.
- 310 7. National Population Commission. Nigeria Demographic and Health Survey 2013.
311 Abuja. National Population Commission. October 2013.
- 312 8. UNICEF Health Statistics. The state of the world's children. New York. UNICEF
313 2006.
- 314 9. Orimadegun AE, Adepogu AA, Akinyinka OO. Adolescent girls' understanding of
315 tetanus and prevention: Implications for the disease control in Western Nigeria.
316 Front Public Health 2014; 2 (24) ; 1-7. DOI: 10.3389/fpubh.2014.00024.
- 317 10. Bamidele JO, Umoh SH. Awareness and compliance of women of childbearing age
318 in a Nigerian urban community with the TT1-TT5 immunization schedule. Sahel Med
319 J 2003; 6 (4) : 121-125. DOI: 10.4314/smj2.v6i4.12849.

- 320 11. Bamidele JO, Umoh SH. Awareness and compliance of antitetanus immunization
321 among adult females in a tertiary institution in Nigeria. Afri J Clin Exper Microbiol.
322 2004; 5(2): 208 – 216. DOI: 10.4314/ajcem.v5i2.7379.
- 323 12. Anderson DR, Sweeny DJ, Williams TA. Sampling and sampling distribution;
324 determining the size of sample. In: Introduction to statistics, concepts and
325 applications. Second edition. New York. West Publishing Company 1991; 215-64.
- 326 13. Qadir M, Murad R, Mumtaz S, Azmi AA, Rehman R et al. Frequency of tetanus
327 toxoid immunization among College and University female students of Karachi. J
328 Ayub Med Coll Abbottabad 2010; 22 (1) : 147-149. Available at
329 <http://www.ayubmed.edu.pk/JAMC/PAST/22-1/Murad.pdf>
- 330 14. Basher MS. Knowledge and practice about TT vaccination among undergraduate
331 female medical students. Mymensingh Med J 2010; 19 (4) : 520-523. PMID:
332 20956892.
- 333 15. Naeem M, Khan MZU, Abbas SH, Adil M, Khan A et al. Coverage and factors
334 associated with tetanus toxoid vaccination among married women of reproductive
335 age: A cross sectional study in Peshawar. J Ayub Med Coll Abbotabad 2010; 22 (3):
336 136-140. Available at <http://www.ayubmed.edu.pk/JAMC/PAST/22-3/Naeem.pdf>
- 337 16. Perry H, Weierbach R, Hossain I, Islam R. Tetanus toxoid immunization coverage
338 among women in zone 3 of Dhaka City: the challenge of reaching all women of
339 reproductive age in urban Bangladesh. Bull World Health Organ 1998; 76 (5) : 449-
340 457. PMC 2305782.

341