

Bioprospection for Anopheles mosquito's repellency effectiveness of natural volatiles from *Ocimum suave* growing in Dar es Salaam, Tanzania

Abstract

Essential oil was extracted from *Ocimum suave* leaves by water distillation and tested for repellency effectiveness against *Anopheles gambiae* adult mosquitoes. The percentage yield of the essential oil was 0.2%. Six concentrations of *O. suave* essential oil were applied on human skin of four volunteers, and the repellency effectiveness was analyzed by PoloPlus (LeOra software version 1.0, 2002-2014), revealed promising RC_{50} , RC_{75} , RC_{90} and RC_{99} with their confidence limits as 0.1161 mg/cm² (0.02067 - 0.1767 mg/cm²), 0.2823mg/cm² (0.22328 - 0.3654 mg/cm²), 0.4319 mg/cm² (0.35226 - 0.58862 mg/cm²) and 0.98934 mg/cm² (0.54731 - 0.99972 mg/cm²), respectively. *O. suave* essential oil exhibited higher *Anopheles* mosquitos' repellency effectiveness which merits further scientific attention for the development of natural repellents for the control of malaria and other mosquito borne diseases. These findings provides a scientific evidence and base for formulation for further mosquito repellency semi-field and field trials for the development of cheaper and affordable new mosquito repellent product(s) to meet human healthcare needs in the prevention and control of malaria and other mosquito transmitted infections.

Key words: *Ocimum suave*, essential oil, *Anopheles gambiae*, repellency

INTRODUCTION

Mosquitoes do not only cause nuisance to humans by biting them to get blood, but they also transmit diseases causing parasites and viruses. Female mosquitoes are responsible for the transmission of a number of human blood-borne diseases. Mosquitoes have been shown to transmit the mostly fatal and disabling diseases, the most prevalent being malaria, bancroftian filariasis, yellow fever, dengue fever and several arbovirus infections¹. The transmission of these infections to humans is a function of the availability of competent mosquito vectors; hence, disease control initiatives targeting mosquitoes remain the most successful mechanism for the prevention and control of malaria, bancroftian filariasis, yellow fever, dengue fever and several arbovirus infections². An estimated 3.4 billion people are at risk of malaria, of which 1.2 billion are at high risk of malaria morbidity and death³. There were an estimated 207 million cases of malaria in 2012 (uncertainty range: 135 – 287 million) and an estimated 627,000 deaths (uncertainty range: 473,000 – 789,000)³. 90% of all malaria deaths occur in sub-Saharan Africa³. In 2012, malaria killed an estimated 482,000 children under five years of age majority of

deaths occurred in sub-Saharan Africa. This translates to 1,300 children deaths every day, or one child dying almost every minute³.

Resistance to drugs and pesticides is taking malaria-control and eradication measures to even more demanding levels. The observed resistance shows strong positive correlation with heavy reliance on the pesticides and drugs currently on the market. In some parts of the world where malaria was once almost eradicated using synthetic pesticides such as DDT and antimalarial drugs, malaria has returned with new features not witnessed during the pre-eradication time. These included vector resistance to pesticides and malaria parasite resistance to drugs⁴. In Africa and some other parts of the world, the heavy reliance on pyrethroids has led to the emergence of pyrethroids-resistance in mosquitoes in malaria-endemic settings⁵. The vector resistance to pesticides challenge requires integrated approach that ensures that malaria parasites are not transmitted to humans.

The 2012 WHO Malaria Report estimates that, malaria related illness and mortality have been dramatically declined by the integrated efforts including the distribution of long lasting insecticide treated mosquito nets (LLINs), efficient case management with potent antimalarials, source reduction and selective household spraying⁶. This depicts the current context that, where mosquitoes' multiplication and resistance to pesticides is increasing in response to the prevailing climate change and the use of pesticides available on the market, an integrated approach in dealing with the malaria problem is now becoming an inevitable scientific endeavor. This entails, *by bioprospection, discovery of new compounds with more pesticidal and repellency effectiveness, oviposition deterrence and mosquito attractants* to be part of the integrated vector control initiative that will be applied based on local studies.

Moreover, environmental concerns that synthetic chemicals contaminate the environment due to their poor biodegradability and their direct toxicity to humans and other non-target organisms have made humans to dislike their use, thereby calling for search for safer pesticides and repellents of natural sources that are more biodegradable, more target specific and relatively of low toxicity as one of the constituents of the integrated approach in the fight against mosquito borne diseases, malaria in particular⁷.

The biting behavior of Anopheles mosquitoes is intriguing as at the moment they have been shown to bite people before retiring to bed or wait and bite when people get out of the net at night even when insecticide treated nets (ITNs) are used; hence this challenge calls **an** additional mosquito control tool to completely curtail human-mosquito contacts when not under ITNs. Measures to prevent humans from mosquito bites are currently the most important measure for the control of malaria². In comparison to vaccines and chemoprophylaxis as means of personal protection, repellents are an inexpensive and practical means of personal protection against nuisance and disease vector bites, in conjunction with other or when other control measures are not practical⁸⁻¹⁰. The mosquito repellent products commonly available on the market contain *N,N*-diethyl- 3-toluamide (DEET) as the active ingredient¹¹. However, DEET has operational disadvantages due to its allergic reactions and toxicity to man¹². DEET has additional disadvantages due to its ability to act as a good solvent for synthetic materials and plastics, which has led to the search for safer alternative synthetic and natural repellents². Previously, a list of 346 plant species from 276 genera and 99 families were shown to have various effects against mosquitoes such as toxicity, oviposition deterrent and repellency¹³.

In this bioprospection initiative for eco-friendly, safer and effective natural repellents we have put on test essential oil from *O. suave* as a possible candidate for incorporating into an integrated malaria vector control initiative, as additional means of malaria prevention and control, and to provide a useful complement to insecticide treated nets (ITNs).

Materials and methods

Study area

The laboratory study was carried out at Ubwari laboratory of the Amani Centre of the National Institute for Medical Research (NIMR) situated 1 km near Muheza town.

Plant materials

Leaves of *O. suave* were collected from Pugu forest near Dar es Salaam in Tanzania.

Hydro-distillation

The extraction process was carried out in the Phytochemistry Laboratory at the Department of Traditional Medicine Research in NIMR in Dar es Salaam, Tanzania. The fresh leaves were grinded using an electric blender and immediately transferred into a round bottomed flask and mounted on Clevenger apparatus for hydro-distillation to obtain essential oil. Temperature was first set at 100°C and later, after boiling was reduced to 50 °C. The condensed vapor and afforded oil was separated using a separating funnel by taking advantage of their mutual immiscibility.

The percentage yield was determined using the formula:

$$\text{Percentage yield} = \frac{\text{Weight of essential oil}}{\text{Weight of leaves}} \times 100$$

Laboratory based mosquito repellency assay

Adult *Anopheles gambiae* aged between 5 and 9 days were obtained from a laboratory colony maintained at 27 ± 2°C and 70-80% relative humidity. Larvae were fed a 3:1 mixture of dog biscuits, and yeast powder. 12:12 (light:dark) photoperiod was maintained. Adults were provided with a 10% sucrose solution. Observations of repellency were made using female mosquitoes starved for the preceding 7 hours. Serial dilutions of candidate repellent were made with ethanol to identify effective dose range. Four volunteers applied incremental doses of 0.028, 0.056, 0.121, 0.186, 0.251 and 0.316g/590cm² were applied on the test forearm. A single test comprised of continuous use of the same mosquitoes by the same volunteer and was completed in one day. Two replicates of this process were obtained using different batches of *An. gambiae* mosquitoes in two days (each replicate test for one extract was completed in one day). 1 ml of ethanol was applied evenly using a syringe to ≈590 cm² of the forearm skin between the wrist and elbow and allowed to dry (for 1 min) and the arm was inserted into a cage containing 50 mosquitoes. Before insertion of the arm into the cage containing 50 female mosquitoes, the hands were protected by medical gloves. The forearm applied with diluents (ethanol) was inserted into the cage first and the number of mosquitoes that landed in three minutes period was recorded. During testing, the volunteers were advised to avoid movement of the arm. The control forearm was carefully withdrawn and this arm was then treated with the lowest dose of repellent in 1 mL alcohol and

allowed to dry for one minute. The treated arm was placed in the cage for another three minute period and observed for mosquito landings. The number of landings was also recorded. This procedure was repeated for each additional incremental repellent dose. Successive tests were carried out one after the other without delay and the repellent dose at each test was calculated as the sum of the doses applied to arrive at the cumulative dose for each test.

In the control experiment, 1 mL ethanol was applied on the other forearm (left arm) and allowed to dry. This forearm was inserted in the cage for three minutes to verify that the number of landings is approximately ≥ 10 per one minute, as was observed at the beginning of the experiment. If the rate was < 10 females in one minute, the results of this experiment were discarded. Data analysis was carried out using **by** PoloPlus (LeOra software version 1.0, 2002-2014), employing probit analysis model, to obtain RC_{50} , RC_{75} and RC_{99} , and their confidence limits.

Percentage repellency was calculated as:

$$\text{Mosquito repellency (\%)} = \frac{100(1 - T)}{C}$$

Where:

T = Number of probing mosquitoes on treated arm;

C = Number of probing mosquitoes on control arm.

Results and discussion

The yield of *O. suave* oil was 0.2%. The six different concentrations of *O. suave* essential oil provided percentile repellence concentrations, $RC_{(\%)}$ (percentile effective doses) as summarized in table 1 below. It was observed that increasing concentration of the repellent oil led to increased percentage repellency. The results revealed promising RC_{50} , RC_{75} , RC_{90} and RC_{99} with their confidence limits as 0.1161 mg/cm² (0.02067 - 0.1767 mg/cm²), 0.2823mg/cm² (0.22328 - 0.3654 mg/cm²), 0.4319 mg/cm² (0.35226 - 0.58862 mg/cm²) and 0.98934 mg/cm² (0.54731 - 0.99972 mg/cm²), respectively.

163 Table 1: Percentile repellence concentrations of *O. suave* essential oil

RC₅₀	Lower limit (mg/cm ²)	RC₇₅	Lower limit (mg/cm ²)	RC₉₀	Lower limit (mg/cm ²)	RC₉₉	Lower limit (mg/cm ²)	SLOPE(SE)	χ²
	Upper limitL (mg/cm ²)		Upper limitL (mg/cm ²)		Upper limitL (mg/cm ²)		Upper limitL (mg/cm ²)		
0.1161	0.02067	0.2823	0.22328	0.4319	0.35226	0.68934	0.54731	0.041+- 0.008	2.697
	0.1767		0.3654		0.58862		0.99972		

164
165 This is clearly depicted by the PoloPlus regression line, figure 1, which appears to be sigmoid in
166 nature. The fact that the essential oil is obtained by using a simple technology that requires no
167 intensive training, and that raw materials are readily available in rural areas, this product
168 promises to be very cost effective and hence accessible to even the poverty stricken households
169 in the rural areas. Previous studies have shown that *Ocimum suave* essential oil can afford up to
170 74% repellency efficacy¹⁰. The study reported the concentration of the essential oil in
171 formulations used and not the amount of the oil on a/per unit skin surface area. This study goes
172 an extra mile to show the concentration of the essential oil on human skin (amount of essential
173 oil per skin surface area) that confers the herein claimed percentage protection at various
174 concentrations of the essential oil within the tested concentration (dose) range, figure 1. *O. suave*
175 essential oil exhibited higher *Anopheles* mosquitos' repellency effectiveness which merits
176 further scientific attention for the development of natural repellents for the control of malaria and
177 other mosquito borne diseases.




Figure 1. PoloPlus regression line showing percentage response (repellency) of *O. suave* essential oil against *An. gambiae* at 3min.

Conclusion

These findings provides a scientific evidence and base for formulation for further mosquito repellency semi-field and field trials for the development of cheaper and affordable new mosquito repellent product(s) to meet human healthcare needs in the prevention and control of malaria and other mosquito transmitted infections. Further research on the essential oil should include 1) toxicological and safety assessment of extracts 2) testing against other mosquito species including wild population 3) investigating complete protection time and the best formulation (concentration, solvent and other additives, etc.) to make the essential oil unfold its full potential effectiveness both in terms of efficacy and complete protection time 4) repellency

testing on other arthropods 5) **complete protection time testing** 6) Description of the mode of action.

Consent

Not applicable.

Competing interests

Authors have declared that no competing interests exist.

References

1. White GB(1973).In: Manson's tropical disease 20th Edition, ELBS low price edition. London.
2. Malebo, H.M., Imeda, C., Kitufe, N.A., Katani, S.J., Sunguruma, R., Magogo, F., Tungu, P.K., Nyigo, V.A., Wiketye, V., Mwaiko, G.L., Ogondiek, J.W., Mbogo, G.P., Mhame, P.P., Matata, D.Z., Malima, R., Magesa, S.M., Massaga, J.J., Malecela, M.N. and Kitua, A.Y. (2013). Repellence effectiveness of essential oils from some Tanzanian *Ocimum* and *Hyptis* plant species against afro-tropical vectors of malaria and lymphatic filariasis. *J. Med. Plants Res.* 7(11), 653-660. <http://www.academicjournals.org/JMPR>, DOI:10.5897/JMPR012.732.
3. **World Malaria Report, 2012. World Health Organization, Geneva, Switzerland.**
4. Sharma VP. 1996, Re-emergence of malaria in India. *Indian Journal of Medical Research.*;103:26–45.]
5. Karunamoorthi, K., Sabesan, S., 2013. Insecticide resistance in insect vectors of disease with special reference to mosquitoes: a potential threat to global public health. *Health Scope* 2 (1), 4–18. <http://dx.doi.org/10.5812/jhs.9840>.
6. **World Malaria Report, 2012. World Health Organization, Geneva, Switzerland.**
7. Karunamoorthi, K., 2012a. Medicinal and aromatic plants: a major source of green pesticides/risk-reduced pesticides. *J. Med. Aromat. Plants* 1, e137. <http://dx.doi.org/10.4172/2167-0412.1000e137>
8. WHO. *International Travel and Health Vaccination Requirement and Health Advice*. Geneva, Switzerland: World Health Organization; 1995.
9. Gupta RK, Rutledge LC. Role of repellents in vector control and disease prevention. *American Journal of Tropical Medicine and Hygiene*. 1994;50(6, supplement):82–86. **[PubMed]**

- 231 10. Malebo, H.M., Nguruwe, R., Lugimbana, L., Sambu, E., Malecela, E., Kamugisha, M.L., Senkoro,
232 K. & Magesa, S.M. (2005). Efficacy of *Ocimum suave* volatile oil formulation against man-biting
233 mosquitoes in Muheza, north-east Tanzania. *Tanzania Health Research Bulletin* 7, 25-30.
- 234 11. Hayes, Hand book of pesticides toxicology, volume one, third edition
- 235 12. Usavadee Thavara, Laboratory and Field Evaluations of the Insect Repellent 3535
- 236 13. Sukumar, K. Michael, J.P., Boobar, L.R. (1991) Botanical derivative in mosquito control: A
237 review. *J Am Mosq Control Assoc.* 7, 210-237.
- 238
- 239
- 240
- 241
- 242