

ANTI PLASMODIAL ACTIVITY OF BEE STING IN *PLASMODIUM BERGHEI* INFECTED MICE

(Please note correction in blue and word to be deleted in underline in red)

Original research article

ABSTRACT

Aim: Based on traditional claims and practice, the antiplasmodial activity of bee stings and its effect on haematological indices was investigated in *P. berghei* infected mice.

Methodology: Sixteen albino mice were intraperitoneally infected with chloroquine sensitive *P. berghei* strain and divided into four groups each consisted of four animals. Group I was set up as negative control of 0.2ml normal Saline/kg body weight, group II as 5mg chloroquine/kg body weight, group III had suppressive treatment and group IV was administered curative treatment. The thin blood smear was used to determine the parasite counts and the haematological parameters were estimated on day 7.

Results: The result of percentage chemosuppression shows that bee stings suppress the parasitaemia to 56.6%. Also, the suppressive and curative groups show longer mean survival period of 15.0 and 20.0 respectively. The haematological studies show that the level of packed cell volume (PCV) and haemoglobin concentration (HB) of infected untreated group was significantly ($p < 0.05$) lower when compared with all other experimental groups, whereas chloroquine treated group shows significant increase compared to the bee treated groups. The Red blood cell (R.B.C.) counts were significantly ($p < 0.05$) lowered in infected untreated group when compared with suppressive and chloroquine treated groups. However the white blood cell (WBC) counts were significantly ($p < 0.05$) higher in infected bee sting treated when compared to the infected untreated and infected chloroquine treated groups.

Conclusion: Based on the result obtained, this study confirms the antiplasmodial activity of bee stings and suggests its potential as drug agent or lead against malaria.

Keywords: Antiplasmodial, bee sting, *P. berghei*, Mice

INTRODUCTION

Malaria is an important cause of morbidity and mortality in tropical countries. The increase in resistance of the malaria parasite to the currently available antimalarial drugs result in the death of millions of people over the past 30 years (Gessler *et al.*, 1995). For many years, traditional medicines have been used for treatment of malaria and they are the source of artemisinin and quinine derivatives. They are of plant source and are effectively used in malaria treatment (Bodeker and Willcox, 2000). The high cost and difficulties encountered by the rural settlers in obtaining these drugs make them go for available traditional medicine such as herbal remedy which are sometimes perceived as being more effective than conventional antimalarial drugs including artemisinin combination therapy (ACT). In view of the problem associated with antimalarial drug resistance, new drug or drug combination are urgently required for the treatment of malaria. Preferably, the new drugs should have novel mechanism of action or be chemically different from drug in current use (Tijani *et al.*, 2008).

Bee (*Apis mellifera*) sting is obtained from the tip of the abdomen of honey bee of the class insect. It is believed that the venom released by bee during stings boost the immune system's in order to fight many disease including arthritis, malaria and so on. Bee sting are painful and are therefore avoided by many people. The bee dies soon after it has stung and released its venom (Luciano *et al.*, 2002). This is due to the destruction of its abdomen, digestive tract, muscle and nerve. Mellitin is the most important and abundant component of bee sting (about 52%) and it acts as a strong anti-inflammatory agent that induce production of cortisol in the body (Meier and White, 1995). *In vitro* intra erythrocytic development of malaria parasite is strongly inhibited by phospholipase A2 from bee sting. This inhibition depends on enzyme activities that require the presence of serum lipoprotein in the parasitic culture medium (Luciano *et al.*, 2002). Bee sting has been reported traditionally for antimalarial activities but scientific information is limited or absent on the therapeutic efficacy and safety for clinical application. Therefore, in this study the anti plasmodial activities of bee sting in *P. berghei* infected mice was investigated.

MATERIALS AND METHODS

58 **Experimental animals**

59 Swiss adult albino mice weighing 20-24g were obtained from animals breeding unit of the
60 Department of Biochemistry, University of Ibadan, Oyo State. They were housed in plastic cages
61 and maintained under standard laboratory conditions.

62 **Parasite**

63 Chloroquine– sensitive strain of *Plasmodium berghei* (NK-65) was obtained from the National
64 Institute for Pharmaceutical Research and Development (NIPRD), Idu, Abuja, Nigeria.

65 **Honey bee**

66 Life healthy honey bees were obtained from bee hive in Bosso Dam, Minna, Niger state. They
67 were house in wooden cage and were constantly given honey and fruits.

68 **Animals Grouping:**

69 The mice were group into four (4) groups of four (4) mice each. The groups were labeled as;
70 group 1 (infected not treated), group 2 (infected treated with standard), group 3 (infected
71 suppressive treated) and group4 (infected curativetreated).

72 **Innocation of parasite:**

73 A donor mouse which had been infected with *Plasmodium berghei* was anesthetized and the
74 blood was immediately diluted with normal saline and passage intraperitoneallyto healthy mice
75 with 0.2ml of the blood containing about 1×10^7 parasitized red blood cells.

76 **Parasitaemia count**

77 The method of Giemsa blood smear was usedto count the number of the parasite red blood cells.
78 The blood was taken from the tail of mice and thin blood smear was prepared using
79 Giemsa stain and the parasitemia was calculated by determining the average of red blood cells
80 infected divided by total red blood cells times 100.

81

82 **Treatment**

The experimental mice were treated as follows:

Group 1: infected and received 0.2ml of normal saline daily.

Group 2: infected and received 5mg per kg body weight chloroquine intraperitoneally (treatment commenced after 72 hours infection).

Group 3: infected and received intradermal bee sting daily (Suppressive test: this was done according to the methods of Knight and Peters (1980). The treatment commenced immediately after infecting the animals and the parasitaemia was determined from drop of blood collected from the tail of the mice for smeared on to the slide to make thin film. The film were then stained with Giemsa and view microscopically at magnification of x100 to determine the average percentage (%) parasitemia

Average % suppression =

$$\frac{\text{Average \% parasitemia of infected untreated} - \text{Average \% parasitemia of infected treated}}{\text{Average parasitemia of infected untreated}} \times 100$$

Average parasitemia of infected untreated

Group 4: infected and received intradermal bee sting daily for 5 days. (Curative test i.e. treatment commenced after 72 hours of *P. berghei* infection). On each day, drops of blood were collected from the tail each mouse smeared on the slide to make a thin film. The film were stained with Giemsa stain and view microscopically at magnification of x100

Mean Survival Time

The mean survival time of the mice in each group over a period of 28 days inoculations (Do – D28) was calculated. The deaths occurring within this period were recorded. The blood smear of the mice that survive beyond 28 days was also monitored. The Mean Survival Time (MST) for each group was calculated as

$$\text{MST} = \frac{\text{Sum of survival time of all mice in a group (days)}}{\text{Total number of mice in that group}}$$

Hematological Parameters

Automated haematologic analyzer SYSMEX KX21, a product of SYSMEX Corporation, Japan, was used to determine the haematological indices including the packed cell volume (PCV), red blood cell count (RBC), white blood cell count (WBC) and Haemoglobin concentration (Hb) as described by Dacie and Lewis (1991).

Ethical Clearance

Ethical Clearance was given by Federal University of Technology, Minna/Nigerian Ethical Review Board (CUERB) in accordance with International standard on care and use of experimental animals.

Consent

Not Applicable

Competing of Interests

Authors have declared no competing interest.

Statistical analysis

The data obtained were subjected to Analysis of Variance (ANOVA) using statistical package (SPSS) version 16. Means were separated using Duncan's Multiple Range Test (DMRT). Significance was accepted at $P < 0.05$. The data is given as mean $\pm$ SEM.

RESULTS

CURATIVE

The average parasitaemia of the infected untreated group, infected treated with 5mg of chloroquine and the bee sting, early treated group mice (prophylactic) are shown in figure 1. The negative control group (infected untreated) show a gradual increase in the parasitaemia count when compare to the bee sting treated group which show decrease in the level of parasitaemia up to day 7. The chloroquine treated group on the other hand show gradual increase in the parasitaemia till day 3 when thereafter continue to decline.




Figure 1: Changes in *in vivo* Antiplasmodial Activity of Honey bee sting Against *P. berghei* infected mice

SUPPRESSIVE

The percentage chemosuppression by bee sting and chloroquine treated group is shown in Table 1. The percentage chemosuppression by bee stung and chloroquine was 56.60 and 67.90 respectively.

Table 1: Percentage Chemosuppression by Bee sting and Chloroquine

Treatment	Parasitaemia Count	% Suppression
-----------	--------------------	---------------

151	Infected not treated	53.00 ± 0.00	-
152	Infected chloroquinetreated	17.00 ± 0.20	67.90
153	Infected Suppressive treated	23.10 ± 0.10	56.60

154 Values are expressed as Mean ±SEM.

155

156

157 Meansurvival period (M.S.P)

158 The mean survival period of all the experimental groups are shown in the Table 2. The infected
 159 untreated group shows the lowest M.S.P of 12 days while the infected treated with standard,
 160 suppressive and curative show survival days of 17,15and 20 days respectively.

161 **Table 2: Average Mean survival period of *Plasmodium berghei*infected mice treated with**
 162 **chloroquine and bee sting**

163	GROUP	SURVIVAL DAYS
164	Infected 0.2ml of normal saline	12.0
165	Infected 0.2ml of chloroquine	17.0
166	Infected Suppressive treated	15.0
167	Infected Curative treated	20.0

168

169 Haematological parameters

170 The results of haematological studies of *P. berghei* infected mice are presented in Table 3. The
 171 level of PCV and HB of infected untreated group was significantly (11.70 ±1.05and 3.90±0.55)
 172 lower when compare with all other experimental group andchloroquine treated group shows
 173 significant(p<0.05) increase when compared to the bee treated group. The R.B.C count

wassignificantly ($p<0.05$) lowered in infected untreated groupwhen compare with curative andchloroquine treated groups. However the WBC counts wassignificantly ($p<0.05$) higher in infectedbee sting treated groups when compare to the infected untreated and infected chloroquine treated mice.

Table 3: Effects of Bee Stings on Haematological parameter of *P.berghei* infected mice

	PCV(%)	Hb(g/dL)	RBC($\times 10^6$ cells/ μ L)	WBC($\times 10^3$ cells/ μ L)
INT	11.70 $\pm$ 1.05 ^a	3.90 $\pm$ 0.55 ^a	6.90 $\pm$ 0.61 ^a	9.02 $\pm$ 1.23 ^b
IQT	25.23 $\pm$ 2.10 ^c	8.41 $\pm$ 0.70 ^c	7.80 $\pm$ 0.34 ^b	7.95 $\pm$ 0.96 ^a
IST	22.61 $\pm$ 2.68 ^b	7.53 $\pm$ 0.88 ^b	6.50 $\pm$ 0.56 ^a	10.21 $\pm$ 1.04 ^c
ICT	23.11 $\pm$ 3.01 ^b	7.70 $\pm$ 0.14 ^b	12.01 $\pm$ 1.98 ^c	10.09 $\pm$ 1.21 ^c

Values are expressed as Mean $\pm$ SEM. Each mean is an average of four replicate (n=4). Values with different superscript (a, b, or c) are significantly different ($p<0.05$). INT: infected not treated, IQT: infected chloroquine treated, IST: infected suppressive treated, ICT: infected curative treated.

DISCUSSION

Many studies have been carried out in recent years on the pharmacological effects of bee products such as honey, propolis, bee pollen and bee venom (Gheldof and Engeseth, 2002; Taormina *et al.*, 2001). They have been reported to have analgesic, antimicrobial, anti-inflammatory and immunological benefits (Tonkset *et al.*, 2007). But little or no scientific data exist to validate the activities of traditional medicine of bee stings against malaria. Bhattacharya (2013) has reported that plasmodium parasites are susceptible to toxins and alcohol. The current anti malarial drugs derived from plants or synthesized (such as chloroquine and artemisinin) are beset with the problem of resistance and toxicity from the use of single compounds. . It has also been reported that the chloroquine resistance was as a result of a single point mutation in the transporter gene *Pfcr*t (Mehlota *et al.*, 2008) and that artemisinin resistance in *P. falciparum* has also been detected in Cambodia (Noedlet *et al.*, 2008). Therefore, it is important that their claimed

for antimalaria activities are investigated in order to establish their efficacy and determine the potential as a source of new antimalaria drug.

Some studies have reported that bee sting poses some therapeutic properties such as boosters of the immune system's ability to fight many diseases including arthritis and so on. The most destructive component of bee sting has been reported as phospholipase A2, the enzyme that inhibits the intraerythrocytic development of malaria parasite (Luciano *et al.*, 2002).

The significant ($P < 0.05$) decrease in parasitaemia of both the chloroquine and the bee sting treated groups compared to the negative control reported in this study could be attributed to the potency of bee sting against *Plasmodium berghei*. Although, the initial increase in the parasitaemia observed in the bee sting prophylactic group may be attributed to the accumulation of the venom before it inhibits the development of the parasite. The longer mean survival period observed in the bee sting treated group compared to the negative control group further showed the efficacy of the bee sting as an antimalaria agent.

Haematological parameters like anaemia, thrombocytopenia and leukocytosis have been documented in malaria infection (Fleming, 1981). Reactive oxygen species are produced by the body during the activation of the immune system thereby resulting in haemoglobin degradation (Loria *et al.*, 1999). Therefore, haematological parameters were always used to assess the effect of malarial parasite on blood components.

The significant decrease in the level of PCV of the infected untreated group is an indication of anaemic condition caused by the malarial infection. However, the infected suppressive and infected curative treated groups show an increase in the packed cell volume level. Thereby, indicating the ameliorative potentials of bee sting on the effect caused by malaria on haematological parameters. White blood cells (WBC) play an active role in the immune system. Thus, the significant increase in the WBC counts in the bee sting treated group when compared with other experimental groups suggest that bee sting stimulates immune response.

CONCLUSION

In conclusion, the result of this study indicates that bee sting possess some antimalaria property which is justify by significant plasmodial parasite clearance compared to chloroquine and ameliorative effect on symptoms cause by the infection.

REFERENCE

- Bhattacharya D. (2013). Transmission Blocking of Year Round Resistant Malaria in Koraput (India) by OMARIA – A New Antimalarial Phytotherapy, *British Journal of Pharmaceutical Research*, Vol. 3(1): pp.54-77.
- Bodeker, G. and Willcox, M.L.(2000). Conference report: the first international meeting of the Research Initiative on Traditional Antimalarial Methods (RITAM), *Journal of Alternative and Complementary Medicine*, 6, 195-207.
- Dacie, J. V. and Lewis, S. M. (1991). Practical Haematology, 7th edition. Churchill Livingstone. Edingburgh
- Fleming, A.F. (1981). Hematological Manifestation of Malaria and Other Parasitic Disease. *Clinical Hematology*.10, 983-1011.
- Gessler C, Nkunya H, Chollet J, Heinrich M, Tanner, M. (1995). Tanzanian medicinal plants used traditionally for the treatment of malaria: *in vivo* antimalarial and *in vitro* cytotoxic activities. *Phytotherapy Research*, 9:504–508.
- Gheldof, N., and Engeseth, N.J. (2002). Antioxidant capacity of honeys from various floral sources based on the determination of oxygen radical absorbance capacity and inhibition of *in vitro* lipoprotein oxidation in human serum samples. *Journal of Agricultural and Food Chemistry*, 50 (10), 3050-3055.
- Knight DJ and Peters W (1980). The antimalarial action of N-benzoyloxydihydrotriazines. The action of clociguanil (BRL50216) against rodent malaria and studies on its mode of action. *Annals of Tropical Medicine and Parasitology*, 74: 393-404.

Loria, P., Miller, S., Foley, M. and Tilley, L., (1999). Inhibition of Peroxidative Degradation of Haem as the Basis of Action of Chloroquine and Other Quinoline Antimalarials. *Biochemical Journal*; 339: 363-370.

Luciano, A., Junitsu I., Anil G. and Helge Z. (2002) Bee sting Phospholipase Inhibits Malaria Parasite Development in Transgenic Mosquitoes. *Journal of Biological Chemistry*, 43(1): 40839–40843

Mehlotra RK, Mattera G, Bockarie MJ, Maguire JD, Baird JK, Sharma YD et al. (2008). Discordant patterns of genetic variation at two chloroquine resistance loci in worldwide populations of the malaria parasite *Plasmodium falciparum*. *Antimicrobial Agents Chemotherapy* 6: 2212-2222.

Meier, J and White, J (1995). Clinical Toxicology of Animal Venoms and Poisons. CRC Press, Volume 236: 768

Noedl H, Se Y, Schaefer K, Smith BL, Socheat D, Fukuda MM. (2008). The artemisinin resistance in Cambodia 1 study, C.: evidence of artemisinin resistant malaria in Western Cambodia. *N Engl. J. Med.* 2008; 359: 2619-2620.

Tijani AY, Uguru MO, Salawu OA (2008). Anti-pyretic, anti-inflammatory and anti-diarrhoeal properties of *Faidherbia albida* in rats. *African Journal of Biotechnology*. 7(6): 696-700.

Taormina, P.J., Niemira, B.A., Beuchat, L.R. (2001). Inhibitory activity of honey against food-borne pathogens as influenced by the presence of hydrogen peroxide and level of antioxidant power. *International Journal of Food Microbiology*, 69, 217-225.

Tonks, A. J., Dudley, E., Porter, N.G., Parton, J., Brazier, J., Smith, E.L., & Tonks, A. (2007). A 5.8-kDa component of manuka honey stimulates immune cells via TLR4. *Journal of Leukocyte Biology*, 82(5), 1147-1155.