

2 **How different is the dietary intake of individuals with** 3 **metabolic syndrome?**

4

5

6 **ABSTRACT**

7 **Background:** Metabolic syndrome has turned out to be a chief public health concern,
8 but the role of diet in the etiology of metabolic syndrome is not well understood.

9 **Aims:** This study aimed at assessing whether individuals with metabolic syndrome
10 had a specific dietary intake and how it compared with the Recommended Nutrient
11 Intakes for Malaysia and the Medical Nutrition Therapy Guidelines by the Malaysian
12 Dietitians' Association.

13 **Study design:** Cross-sectional study

14 **Place of study:** Putrajaya, Malaysia

15 **Duration of study:** Eight weeks

16 **Methodology:** Waist circumference, systolic blood pressure, diastolic blood pressure,
17 triglycerides, HDL cholesterol and fasting plasma glucose were collected in 659
18 randomly selected subjects. A total of 275 out of 322 subjects with metabolic
19 syndrome completed the self-administered three-day food record. Data were analysed
20 with NutritionistPro, and were compared with the Recommended Nutrient Intakes for
21 Malaysia (2010), based on age and gender. Metabolic syndrome was defined
22 according to the 'Harmonized' criteria.

23 **Results:** Subjects' mean age was 34.51 ± 8.30 years, 202 Females: 73 Males.
24 Comparison of subjects' food records with the Recommended Nutrient Intakes for
25 Malaysia revealed that protein consumption was significantly higher among study
26 subjects compared to the Recommended Nutrient Intakes for Malaysia (62 g/day for

27 males; 55 g/day for females), $P < .001$. All subjects (100%) had low fibre (< 20 g/ day)
28 intake. Conversely, most subjects (72.0%) had high sodium intake (≥ 2400 mg/ day).

29 **Conclusion:** This study highlights the importance of assessing the dietary intake of
30 each individual with metabolic syndrome. There is a discrepancy between the dietary
31 intake among individuals with metabolic syndrome and the Recommended Nutrient
32 Intakes for Malaysia or Medical Nutrition Therapy Guidelines by the Malaysian
33 Dietitians' Association. Notwithstanding, our study cannot declare that diet induces the
34 disease, yet it definitely sheds more light on the need for appropriate and professional
35 dietary measures to be integrated in the management of individuals with metabolic
36 syndrome.

37 **Keywords:** metabolic syndrome, dietary intake, medical nutrition therapy, Recommended
38 Nutrient Intakes

39 1 INTRODUCTION

40 Metabolic syndrome is a cluster of risk factors, including obesity, dyslipidemia, hypertension, glucose
41 intolerance and insulin resistance, that increases the risk for cardiovascular disease and type 2
42 diabetes mellitus [1,2]. A remarkable increase in the number of people with metabolic syndrome has
43 occurred worldwide [3]. A national survey conducted from January to December 2008 revealed that
44 the overall prevalence of metabolic syndrome was 32.1%, 34.3%, 37.1% and 42.5%, based on
45 definitions from the World Health Organisation, the National Cholesterol Education Program ATP III,
46 the International Diabetes Federation and 'Harmonized', respectively [4]. Identification of particular
47 foods and nutrients associated with metabolic syndrome is imperative for the formulation of dietary
48 recommendations [5]. Further, the combined effect of food and nutrients, which are consumed
49 together, may be synergistic. In this context, dietary intake analysis, which concentrates on
50 macronutrients and their composition, may impart supplementary data to relate diet to disease risk.

51 Notwithstanding, previous studies have recognised food intake composition and examined their
52 association with metabolic syndrome, but these studies were predominantly from Europe [6,7], Korea
53 [8,9,10,11], the Middle East [12,13,14], North America [15], Samoa [16], and the United States

54 [17,18]. Dietary habits of Asians, including Malaysians, are considerably different from those countries
55 [19].

56 Despite the evident links to metabolic syndrome, little is known about the dietary intake of individuals
57 with metabolic syndrome. Management for individuals with metabolic syndrome generally includes
58 diet modifications, albeit there is a lack of supporting evidence. Hence, the aim of this study,
59 conducted among government employees with metabolic syndrome, was to analyse their dietary
60 intake and how they compared with the Recommended Nutrient Intakes for Malaysia [20] and the
61 Medical Nutrition Therapy Guidelines by the Malaysian Dietitians' Association [21].

62

63 **2 METHODOLOGY**

64 **2.1 Study procedure**

65 Subjects were requested to fill in a three-day food record, which was checked for completeness and,
66 where necessary, was clarified by asking the subjects questions. We also collected data that were
67 routinely obtained in a health examination, including anthropometric measures (waist circumference),
68 blood pressure measurement and biochemical data. This study conformed to the principles of the
69 Helsinki Declaration and was approved by the Ethics Committee of the Faculty of Medicine and
70 Health Sciences, Universiti Putra Malaysia (Reference No UPM/FPSK/100-9/2-MJKEtikaPen
71 (JPD_Jan (12) 04). Written informed consent was obtained from each subject. The flow chart of data
72 collection was demonstrated in Figure 1.

73

74

75

76

77

78




Figure 1 Flow chart of data collection

2.2 Study subjects

A total of 675 government employees were randomly selected from a list of 3173 government employees at the five government agencies, using a Table of Random Numbers.

Of the 322 government employees with metabolic syndrome, 275 subjects (73 males and 202 females) aged 18 to 59 years completed the self-administered three-day food record.

2.3 Dietary assessment

Dietary intake data was obtained through use of a food record for three days (two consecutive weekdays and one weekend day). Each subject was familiarised with portion sizes and the relevant procedures for successfully completing a food record. The records were checked for any mis-recorded or missing information when subjects returned the records.

To estimate compliance with respect to recording everything, a ratio between total daily energy intake (EI) and basal metabolic rate (BMR) was calculated [22]. The basal metabolic rate was obtained from the Lifecorder e-Step activity monitor (Suzuken Company Limited, Nagoya, Japan). A ratio of EI/BMR below 1.2 was used as the cut-off point to identify non-compliance in recording [23].

2.4 Anthropometric measurement

Waist circumference was measured at the end of normal expiration between the lowest rib and the iliac crest using a non-elastic measuring tape to the nearest 0.1 cm. Waist circumference was categorised as 'normal' (< 90 cm for men and < 80 cm for women) and 'high risk' ($\geq$ 90 cm for men and $\geq$ 80 cm for women) according to the cut-off points proposed by the 'Harmonized' definition [24].

2.5 Blood pressure measurement

Duplicate measurements of blood pressure (in mmHg) were measured two minutes apart with a digital automated blood pressure monitor, Omron HEM-907 (Omron, Japan), after the subjects had rested in a seated position for five minutes [25]. An average of the two readings was taken. The right arm was placed on a table and a cuff was placed on the right upper arm. The mean value of the two measurements was computed. Elevated blood pressure was defined as either elevated systolic blood pressure ($\geq$ 130 mmHg) or elevated diastolic blood pressure ($\geq$ 85 mmHg) alone, or a combination of both [24].

2.6 Biochemical measurements

Blood samples were collected in the morning after subjects had undergone an overnight fast. A total of 5 ml of venous blood was collected with a syringe and needle. All vials were delivered in a cool box packed with dry ice to an accredited pathology laboratory (Gribbles Pathology (M) Sdn Bhd, Bangi, Malaysia) after collection of all blood samples.

2.7 Definition of metabolic syndrome

Metabolic syndrome and metabolic risk factors were defined according to the 'Harmonized' criteria [24]. There are five metabolic risk factors according to the 'Harmonized' definition, including waist circumference, HDL cholesterol, triglycerides, fasting glucose, systolic blood pressure and diastolic blood pressure (Table 1).

127

Table 1: Five metabolic risk factors according to 'Harmonized' definition

Clinical measure	Insulin resistance	Body weight	Lipid	Blood pressure	Glucose
Harmonized (Alberti <i>et al.</i> , 2009) [24]	None, but any three of the five risk factors	Increased waist circumference (Gender- and ethnicity- specific)	TG ≥ 1.7 mmol/L or on TG treatment HDL cholesterol < 1.0 mmol/L in men or < 1.3 mmol/L in women or on HDL cholesterol treatment	$\geq 130/85$ mmHg or on hypertensive treatment	FPG ≥ 5.6 mmol/L or diabetes

128

129 2.8 Statistical analysis

130 NutritionistPro (First Data Bank) was used to analyse dietary data. Data was analysed using SPSS
 131 version 20. Nominal or ordinal variable values were depicted as frequency and percentage, while
 132 interval or ratio variable values were presented as mean $\pm$ standard deviation. One sample *t*-test was
 133 used to test whether there was a significant difference between the sample means and the
 134 Recommended Nutrient Intakes for Malaysia [20] and the Medical Nutrition Therapy Guidelines by the
 135 Malaysian Dietitians' Association [21].

136 3 RESULTS

137 Of the 322 government employees with metabolic syndrome, 275 subjects (73 males and 202
 138 females) aged 34.51 ± 8.30 years completed the self-administered three-day food record (response
 139 rate 85.4%). Table 2 shows the distribution of subjects based on individual metabolic risks.

140

141

Table 2 Distribution of subjects based on individual metabolic risks

Metabolic Risk Factors	Number of subjects (Percentages)	
	Yes	No
Abdominal obesity	265 (96.4)	10 (3.6)
Elevated systolic blood pressure	198 (72.0)	77 (28.0)
Elevated diastolic blood pressure	102 (37.1)	173 (62.9)
Elevated blood pressure	221 (80.4)	54 (19.6)
Low HDL cholesterol	219 (79.6)	56 (20.4)
Elevated triglycerides	135 (49.1)	140 (50.9)
Elevated plasma glucose	49 (17.8)	226 (82.2)

142

143 Based on Table 3, a comparison of the subjects' food records with the Recommended Nutrient
 144 Intakes for Malaysia revealed that protein consumption was significantly higher among study subjects
 145 compared to the Recommended Nutrient Intakes for Malaysia (62 g/day for males; 55 g/day for
 146 females), ($P < .001$). All subjects (100%) had low fibre (< 20 g/day) intake. Conversely, most subjects
 147 (72.0%) had high sodium intake (≥ 2400 mg/day), as portrayed in Table 4.

148

149 **4 DISCUSSION**

150 Dietary intake data was obtained using a food record for three days, which were composed of two
 151 consecutive weekdays and one weekend day [26,27,28,29,30,31,32,33]. Keeping a food record for
 152 three days has proven to be the best approach for assessing actual nutrient intake [34,35,36]. The
 153 choice of the number of days is typically dependent upon pragmatic considerations such as the
 154 willingness of the subject to be compliant [37]. Recording all food consumed for more than one or two
 155 consecutive days is a notorious behaviour modification method for reducing food intake. In spite of
 156 this, a food record is often employed as the gold standard for validating other dietary assessment
 157 methods when observation is not possible [38].

158

Table 3 Comparison between the sample means with Recommended Nutrient Intakes for Malaysia (Ministry of Health, 2005) and Medical Nutrition Therapy
 159 (MNT) Guidelines by Malaysian Dietitians' Association (2005) using One sample *t*-test

Nutrients	Age	Male (n=73)					Female (n=202)				
		Mean±SD	RNI	P-value	MNT	P-value	Mean±SD	RNI	P-value	MNT	P-value
Energy (kcal)											
	19-<30	2110.19±194.15	2440	< .001			2249.46±254.44	2000	< .001		
	30-<51	2350.13±372.55	2460	.049			2270.57±72.39	2180	< .001		
	51-<60	2296.67±515.35	2460	.518			2289.81±258.91	2180	.090		
Carbohydrate (g)		327.77±55.34	200	< .001			327.40±3.01	180	< .001		
			330	.732				230	< .001		
Carbohydrate (%)		57.70±5.97	55	< .001	50	< .001	57.92±5.01	55	< .001	50	< .001
			70	< .001	60	.002		70	< .001	60	< .001
Sugars (%)		6.88±3.25	<15				6.64±3.25	<15			
Protein (g)		82.70±18.76	62	< .001			80.64±14.42	55	< .001		
Protein (%)		14.54±2.53			15	.124	14.23±1.93			15	< .001
					20	< .001				20	< .001
Fat (g)		73.67±17.54	54	< .001			71.91±16.08	46	< .001		
			82	< .001				70	.092		
Fat (%)		29.01±4.56	20	< .001	25	< .001	28.49±5.10	20	< .001	25	< .001
			30	.067	30	.067		30	< .001	30	< .001

PUFA (%)	5.10±2.41	6	.002		5.16±2.13	6	< .001
		10	< .001			10	< .001
MUFA (%)	5.58±2.34	12	< .001		5.53±2.01	12	< .001
		15	< .001			15	< .001
SFA (%)	6.38±2.55	<10			5.72±2.29	<10	
Fiber (g)	6.60±2.97	20	< .001		6.83±3.20	20	< .001
		30	< .001			30	< .001
Sodium (mg)	3099.65±1117.57			<2400	2962.03±925.62		<2400

160

161

162

163

164

165

166

167

168

169

170

171

Table 4 Distribution of subjects based on Recommended Nutrient Intakes for Malaysia (Ministry of Health, 2005) and Medical Nutrition Therapy (MNT)

172

Guidelines by Malaysian Dietitians' Association (2005)

	Recommended Nutrient Intakes for Malaysia				Medical Nutrition Therapy (MNT) Guidelines			
		Male (n=73)	Female (n=202)	Total (n=275)		Male (n=73)	Female (n=202)	Total (n=275)
Carbohydrate (%)	<55	21 (28.8)	53 (26.2)	74 (26.9)	<50	5 (6.8)	13 (6.4)	18 (6.5)
	55-70	51 (69.9)	149 (73.8)	200 (72.7)	50-60	44 (60.3)	117 (57.9)	161 (58.5)
	>70	1 (1.4)	0 (0.0)	1 (0.4)	>60	24 (32.9)	72 (35.6)	96 (34.9)
Sugars (%)	<15	70 (95.9)	200 (99.0)	270 (98.2)				
	≥15	3 (4.1)	2 (1.0)	5 (1.8)				
Protein (%)					<15	46 (63.0)	137 (67.8)	183 (66.5)
					15-20	26 (35.6)	65 (32.2)	91 (33.1)
					>20	1 (1.4)	0 (0.0)	1 (0.4)
Fat (%)	<20	2 (2.7)	8 (4.0)	10 (3.6)	<25	15 (20.5)	50 (24.8)	65 (23.6)
	20-30	46 (63.0)	118 (58.4)	164 (59.6)	25-30	33 (45.2)	76 (37.6)	109 (39.6)
	>30	25 (34.2)	76 (37.6)	101 (36.7)	>30	25 (34.2)	76 (37.6)	101 (36.7)
PUFA (%)	<6	50 (68.5)	132 (65.3)	182 (66.2)				
	6-10	21 (28.8)	67 (33.2)	88 (32.0)				
	>10	2 (2.7)	3 (1.5)	5 (1.8)				
MUFA (%)	<12	72 (98.6)	201 (99.5)	273 (99.3)				

	12-15	1 (1.4)	1 (0.5)	2 (0.7)				
	>15	-	-	-				
	SFA (%)	<10	68 (93.2)	193 (95.5)	261 (94.9)			
		≥10	5 (6.8)	9 (4.5)	14 (5.1)			
	Fiber (g)	<20	73 (100.0)	202 (100.0)	275 (100.0)			
		20-30	-	-	-			
		>30	-	-	-			
Sodium (mg)					<2400	18 (24.7)	59 (29.2)	77 (28.0)
					≥2400	55 (75.3)	143 (70.8)	198 (72.0)

173 However, all methods for assessing dietary intake are subject to a large error sample, especially in
174 overweight or obese populations, in whom a tendency to underestimate food consumption has been
175 demonstrated [39]. The rate of under-reporting of total calorie intake was 3.3% (n = 9) of the total
176 subjects: 7 of the under-reporters were male, while another 2 were female.

177 According to the Medical Nutrition Therapy Guidelines (MNT) by the Malaysian Dietitians' Association
178 [21], 50 to 60% of total energy intake from carbohydrate [40], is recommended. Lower intake of 50%
179 should be considered for individuals with metabolic syndrome who have elevated triglycerides or low
180 HDL cholesterol [41]. Nevertheless, fat consumption should not be less than 25% of total energy
181 intake in order to avoid replacement with excess carbohydrate, which might result in higher
182 triglycerides and lower HDL cholesterol.

183 In this study, it was demonstrated that individuals with metabolic syndrome record a diet richer in
184 protein (%) and poorer in fibre (g) when compared to the recommendations. Fibre consumption was
185 lower than recommended (20 to 30 g/day) in all the subjects, which may have contributed to fat
186 accumulation and insulin resistance.

187 A study conducted by Aljada and colleagues demonstrated that high dietary fat consumption is related
188 to the activation of the proinflammatory transcription factor (nuclear factor kappa-beta) and oxidative
189 stress [42]. On the contrary, high fruit and fibre consumption has no proinflammatory capacity
190 compared with a diet rich in fat, even if the diet has a similar calorie content [43].

191 The healthy and effective approach for long-term weight loss is a reduced-calorie diet, comprising of a
192 modest deficit of 500 to 1000 kcal/day. Sustained dietary modifications may require a registered
193 dietitian's intervention to safeguard a sufficient micronutrient consumption, such as iron, calcium and
194 folate, while decreasing calorie intake. A prospective cohort study [44] has demonstrated that a
195 Mediterranean-style diet was inversely related to the cumulative metabolic syndrome incidence. A
196 Mediterranean-style diet enhances the mental and physical domains of health-related quality of life,
197 such as vitality, physical health, physical function, emotional role and self-perception of health [45],
198 and it reduces the odds of glucose, LDL-cholesterol levels [46], triglycerides and low HDL-cholesterol
199 values [47]. In addition, the Dietary Approaches to Stop Hypertension (DASH) diet combined with
200 lifestyle interventions in the PREMIER study [48] improved blood pressure. Furthermore, a protein

intake of 10–35% of total calorie consumption is suggested by the Institute of Medicine for the general population. A diet rich in complex carbohydrates with 14g of fibre per 1000 calories daily and low in added sugars, $\leq 25\%$ of calorie consumption, is suggested for individuals at risk of or with metabolic syndrome [49].

Afshin and colleagues [50] carried out a comparative risk assessment analysis to approximate the mortality of cardiometabolic disease attributable to 4 metabolic and 11 dietary risk factors in 20 countries of the adult population in the Middle East. The findings demonstrated that a suboptimal diet was the principal risk factor for cardiometabolic disease among 11 countries in the Middle East, equivalent to 48–72% of cardiometabolic disease mortalities. Suboptimal systolic blood pressure was the principal risk factor for cardiometabolic disease mortalities among 8 countries in the Middle East, accounting for 45–68% of cardiometabolic disease mortalities. Suboptimal fasting plasma glucose and body mass index were the third and fourth principal risk factors for cardiometabolic disease death in most Middle East countries. Low consumption of fruits was responsible for 8–21% of cardiometabolic disease mortalities, and low consumption of whole accounted for 7–22% of cardiometabolic disease mortalities among individual dietary factors. These findings emphasise significant differences and similarities in the effect of the metabolic and dietary risk factors on cardiometabolic disease death in Middle Eastern countries, and they inform policy decisions in preventing cardiometabolic disease.

The distribution of calorie from macronutrients in the diets of subjects as observed in this study (Carbohydrate: 57% for males and 58% for females; Protein: 14% for males and 14% for females; Fat: 29% for males and 28% for females) is comparable to that of the diets of Malaysians (Carbohydrate: 59%; Protein: 14%; Fat: 27%). Mean carbohydrate intake of Malaysian adults was about 232 g (59% of the total calorie intake). Malaysian adults reported a mean protein intake of 59 g, which contributed to 14% of total calorie intake. On the other hand, mean fat intake was approximately 50 g for the whole population [51]. In addition, The findings from the dietary survey among type 2 diabetes patients attending an outpatient clinic in Malaysia revealed a similarity in macronutrient contribution to the total calorie intake (Carbohydrate: 57%; Protein: 15%; Fat: 28%) [52]. The study subjects' sodium intake (about 3100 mg for males and 2962 mg for females) was

higher than that of Malaysian adults' mean sodium intake, which was approximately 2575 mg. The mean sodium intake was higher in males than females by about 500 mg [51].

This study had some limitations that warrant indication. First, findings originating from a cross-sectional study do not necessarily signify causality. Second, this study was carried out among government employees. Therefore, the findings obtained may not be applied to the general population.

5 CONCLUSION

This cross-sectional study among government employees further highlights the importance of assessing the dietary intake of each individual with metabolic syndrome. There is a discrepancy between the dietary intake composition among individuals with metabolic syndrome and the Recommended Nutrient Intakes for Malaysia or the Medical Nutrition Therapy Guidelines from the Malaysian Dietitians' Association. Notwithstanding, our study cannot declare that diet induces the disease, yet it definitely sheds more light on the need for appropriate and professional dietary measures to be integrated in the management of individuals with metabolic syndrome.

ACKNOWLEDGEMENTS

The authors have declared that no competing interest exists. The Ministry of Higher Education funded this study. The funders had no role in the study design, data collection and analysis, decision to publish or preparation of the manuscript. We thank the subjects involved in this study for their cooperation.

AUTHOR CONTRIBUTIONS

Conceived and designed the experiments: Chee Huei Phing Hazizi Abu Saad Barakatun Nisak Mohd Yusof Mohd Nasir Mohd Taib. Performed the experiments: Chee Huei Phing Hazizi Abu Saad. Analyzed the data: Chee Huei Phing Hazizi Abu Saad. Wrote the paper: Chee Huei Phing Hazizi Abu Saad

256 **REFERENCES**

- 257 [1] Mortillo, S., Filion, K.B., Genest, J., Joseph, L., Pilote, L., Poirier, P. et al. The metabolic syndrome
258 and cardiovascular risk a systematic review and meta-analysis. **J Am Coll Cardiol**. 2010; 56(14):
259 1113-1132.
- 260 [2] Ford, E.S., Li, C., Sattar, N. Metabolic syndrome and incident diabetes: Current state of evidence.
261 **Diabetes Care**. 2008; 31(9): 1898-1904.
- 262 [3] Cameron, A.J., Shaw, J.E., Zimmet, P.Z. The metabolic syndrome: Prevalence in worldwide
263 populations. **Endocrinol Metab Clin North Am** Endocrinology Metabolism Clinics of North
264 America. 2004; 33(2): 351-375.
- 265 [4] Wan Nazaimoon, W.M., Aziz, A.I., Amir, S., Ikram, S.I., Kamarul, I.M., Khalid, A.K. et al.
266 Prevalence of metabolic syndrome and its risk factors in adult Malaysians: Results of a
267 nationwide survey. **Diabetes Res Clin Pract**. 2011; 91(2): 239-245.
- 268 [5] Kimokoti, R.W., Brown, L.S. Dietary management of the metabolic syndrome. **Clin Pharmacol**
269 **Ther**. 2011; 90(1): 184-187.
- 270 [6] Heidemann, C., Scheidt-Nave, C., Richter, A., Mensink, G.B. Dietary patterns are associated with
271 cardiometabolic risk factors in a representative study population of German adults. **Br J Nutr**.
272 2011; 106(8): 1253-1262.
- 273 [7] Paganiotakos, D.B., Pitsavos, C., Skoumas, Y., Stefanadis, C. The association between food
274 patterns and the metabolic syndrome using principal components analysis: The ATTICA study. **J**
275 **Am Diet Assoc**. 2007; 107(6): 979-987.
- 276 [8] Cho, Y.A., Kim, J., Cho, E.R., Shin, A. Dietary patterns and the prevalence of metabolic syndrome
277 in Korean women. **Nutr Metab Cardiovasc Dis**. 2011; 21(11): 893-900.
- 278 [9] Hong, S., Song, Y.J., Lee, K.H., Lee, H.S., Lee, M., Jee, S.H. et al. A fruit and dairy dietary
279 patterns and glucose tolerance abnormalities in Japanese men. **J Nutr**. 2012; 136(5): 1352-
280 1358.
- 281 [10] Kim, J., Jo, I. Grains, vegetables and fish dietary pattern is inversely associated with the risk of
282 metabolic syndrome in South Korean adults. **J Am Diet Assoc**. 2011; 111(8): 1141-1149.
- 283 [11] Song, Y., Joung, H. A traditional Korean dietary pattern and metabolic syndrome abnormalities.
284 **Nutr Metab Cardiovasc Dis**. 2012; 22(5): 456-462.

285 [12] Amini, M., Esmailzadeh, A., Shafaezadeh, S., Behrooz, J., Zare, M. Relationship between major
 286 dietary patterns and metabolic syndrome among individuals with impaired glucose tolerance.
 287 Nutrition. 2010; 26(10): 986-992.

288 [13] Esmailzadeh, A., Kimiagar, M., Mehrabi, Y., Azadbakht, L., Hu, F.B., Willett, W.C. Dietary
 289 patterns, insulin resistance and prevalence of the metabolic syndrome in women. **Am J Clin**
 290 **Nutr.** 2007; 85(3): 910-918, 2007.

291 [14] Naja, F., Nasreddine, L., Itani, L., Adra, N., Sibai, A., Hwalla, N. Association between dietary
 292 patterns and the risk of metabolic syndrome among Lebanese adults. **Eur J Clin Nutr.** 2013;
 293 52(1): 97-105.

294 [15] Denova-Gutierrez, E., Castanon, S., Talavera, J.O., Gallegos-Carrillo, K., Flores, M.,
 295 Dosamantes-Carrasco, D. et al. Dietary patterns are associated with metabolic syndrome in an
 296 urban Mexican population. **Nutrition.** 2010; 140(10): 1855-1863.

297 [16] DiBello, J.R., McGrahey, S.T., Kraft, P., Goldberg, R., Campos, H., Quesada, C. et al. Dietary
 298 patterns are associated with metabolic syndrome in adult Samoans. **Nutrition.** 2009; 139(10):
 299 1933-1943.

300 [17] Deshmukh-Taskar, P.R., O'Neil, C.E., Nicklas, T.A., Yang, S.J., Liu, Y., Gustat, J. et al. Dietary
 301 patterns associated with metabolic syndrome, sociodemographic and lifestyle factors in young
 302 adults: the Bogalusa Heart study. **Public Health Nutr.** 2009; 12(12): 2493-2503.

303 [18] Lutsey, P.L., Steffen, L.M., Stevens, J. Dietary intake and the development of the metabolic
 304 syndrome: the Atherosclerosis Risk in Communities study. *Circulation.* 2008; 117(6): 754-761.

305 [19] FAO Statistics Division. Food and Agriculture Organization of the United Nations. Accessed 1
 306 May 2011. Available: <http://faostat.fao.org>.

307 [20] Ministry of Health, National Coordinating Committee on Food and Nutrition. Recommended
 308 Nutrient Intakes for Malaysia. A report of the Technical Working Group on Nutritional Guidelines.
 309 Malaysia: Ministry of Health; 2005.

310 [21] Malaysian Dietitians' Association. Guidelines for Dietary Management of Hyperlipidemia.
 311 Malaysia: Malaysian Dietitians' Association; 2005.

312 [22] Macdiarmid, J., Blundell, J. Assessing dietary intake: Who, what and why of under-reporting, **Nutr**
 313 **Res Rev.** 1998; 11(2): 231-254.

314 [23] Goldberg, G., Black, A., Jebb, S., Cole, T., Murgatroyd, P., Coward, W. et al. Critical evaluation of
 315 energy intake data using fundamental principles of energy physiology: 1. Derivation of cut-off
 316 limits to identify under-recording, *Eur J Clin Nutr* . 1991; 45(12): 569-581.

317 [24] Alberti, K., Eckel, R.H., Grundy, S.M., Zimmet, P.Z., Cleeman, J.I., Donato, K.A. et al.
 318 Harmonizing the metabolic syndrome. *Circulation*. 2009; 120(16): 1640-1645.

319 [25] Murphy, M.H., Murtagh, E.M., Boreham, C.A.G., Hare, L.G., Nevill, A.M. The effect of a worksite
 320 based walking programme on cardiovascular risk in previously sedentary civil servants. *BMC*
 321 *Public Health*. 2006; 6(1): 136-143.

322 [26] Kimm, S.Y., Glynn, N.W., Kriska, A.M., Fitzgerald, S.L., Aaron, D.J., Similo, S.L. et al.
 323 Longitudinal changes in physical activity in a biracial cohort during adolescence. *Med Sci Sports*
 324 *Exerc*. 2000, 32(8): 1445-1454.

325 [27] Manios, Y. Design and descriptive results of the "Growth, exercise and nutrition epidemiological
 326 study in preSchoolers": The Genesis study. *BMC Public Health* 2006; 6: 32-40.

327 [28] Manios, Y., Kourlaba, G., Kondaki, K., Grammatikaki, E., Birbilis, M., Oikonomou, E., et al. Diet
 328 quality of preschoolers in Greece based on the Healthy Eating Index. The Genesis study. *J Am*
 329 *Diet Assoc*. 2009; 109(4): 616-623.

330 [29] Myszkowska-Ryciak J, Szymanska J, Harton A, Gajewska D. Evidence of high fat and sugar
 331 intake, and low fiber and selected minerals intake in adult female coeliac patients. *JPCCCR*. 2010;
 332 4(2): 112-115.

333 [30] Roy, M.S., Stables, G., Collier, B., Roy, A., Bou, E. Nutritional factors in diabetics with and
 334 without retinopathy. *Am J Clin Nutr*. 1989; 50(4): 728-730.

335 [31] Samuel-Hodge, C.D., Fernandez, L.M., Hernandez-Roldan, C.F., Johnston, L.F., Keyserling, T.C.
 336 A comparison of self-reported energy intake with total energy expenditure estimated by
 337 accelerometer and basal metabolic rate in African-American women with type 2 diabetes.
 338 *Diabetes Care*. 2004; 27(3): 663-669.

339 [32] Soo, K.L., Wan Abdul Manan, W.M., Abdul Manaf, H., Lee, Y.Y. Dietary practices among
 340 overweight and obese Chinese children in Kota Bahru, Kelantan. *Mal J Nutr*. 2011; 17(1): 87-95.

341 [33] Wada, K., Nakamura, K., Tamai, Y., Tsuji, M., Sahashi, Y., Watanabe, K. et al. Seaweed intake
 342 and blood pressure levels in healthy pre-school Japanese children. *Nutr J*. 2011; 10: 83-89.

343 [34] Bazzarre, T., Yuhas, J. Comparative evaluation of methods of collecting food intake data for
 344 cancer epidemiology studies. *Nutr Cancer*. 1983; 5(3-4): 201-214.

345 [35] Gersovitz, M., Madden, J., Smiciklas-Wright, H. Validity of the 24 hour dietary recall and seven
 346 day record for group comparisons. *J Am Diet Assoc*. 1978; 73(1): 48-55.

347 [36] White, J., Jago, R., Thompson, J.L. Dietary risk factors for the development of insulin resistance
 348 in adolescent girls: A 3-year prospective study. *Public Health Nutr*. 2012; [Epub ahead of print].

349 [37] Rennie, K.L., Wareham, N.J. The validation of physical activity instruments for measuring energy
 350 expenditure: Problems and pitfalls. *Public Health Nutr*. 1998; 1(4): 265-271.

351 [38] Barrett-Connor, E. Nutrition epidemiology: How do we know what they ate? *Am J Clin Nutr*. 1991;
 352 54(1): 182S-187S.

353 [39] Willett, W. *Nutritional Epidemiology*. USA: Oxford University Press, 1998.

354 [40] American Diabetes Association. Report of the Expert Committee on the Diagnosis and
 355 Classification of Diabetes Mellitus. *Diabetes Care*. 2003; 26: S3.

356 [41] NCEP ATP III. Final Report of the Third Report of the National Cholesterol Education Program
 357 (NCEP) Expert Panel on detection, evaluation and treatment of High Blood Cholesterol in *Adults*
 358 (*Adult Treatment Panel III, 2001*). NIH; 2001.

359 [42] Aljada, A., Mohanty, P., Ghanim, H., Abdo, T., Tripathy, d., Chaudhuri, A. et al. (2004). Increase
 360 in intranuclear nuclear factor κ B and decrease in inhibitor κ B in mononuclear cell after a mixed
 361 meal: Evidence for a proinflammatory effect. *Am J Clin Nutr*. 2004; 79(4): 682-690.

362 [43] Dandona, P., Aljada, A., Chadhuri, A., Mohanty, P., Garg, R. Metabolic syndrome: A
 363 comprehensive perspective based on interactions between obesity, diabetes and inflammation.
 364 *Circulation*. 2005; 112 (II): 1448-1454.

365 [44] Tortosa, A., Bes-Rastrollo, M., Sanchez-Villegas, A., Basterra-Gortari, F.J., Nunez-Cordoba, J.M.,
 366 Martiniz-Gonzalez, M.A. Mediterranean diet inversely associated with the incidence of metabolic
 367 syndrome. The SUN prospective cohort. *Diabetes Care*. 2007; 30(11): 2957-2959.

368 [45] Landaeta-Diaz, L., Fernandez, J.M., Da Silva-Grigoletto, M., Rosado-Alvarez, D., Gomez-
 369 Garduno, A., Gomez-Delgado, F. et al. Mediterranean diet, moderate-to-high intensity training
 370 and health-related quality of life in adults with metabolic syndrome. *Eur J. Prev Cardiol*. 2013;
 371 20(4): 555-564.

372 [46] Paletas, K., Athanasiadou, E., Sarigianni, M., Paschos, P., Kalogirou, A., Hassapidou, M. et al.
373 The protective role of the Mediterranean diet on the prevalence of metabolic syndrome in a
374 population of greek obese subjects. J Am Coll Nutr. 2010; 29(1): 41-45.

375 [47] Babio, N., Bullo, M., Basora, J., Martiniz-Gonzalez, M.A., Fernande-Ballart, J., Marquez-
376 Sandoval, F. et al. Adherence to the Mediterranean diet and risk of metabolic syndrome and its
377 components. Nutr Metab Cardiovas. 2009; 19(8): 563-570.

378 [48] Lien, L.F., Brown, A.J., Ard, D., Loria, C., Erlinger, T.P. et al. Effects of PREMIER lifestyle
379 modifications on participants with and without the metabolic syndrome. Hypertension. 2007;
380 50(4): 609-616.

381 [49] Mitch, W.E. Beneficial responses to modified diet in treating patients with chronic kidney disease.
382 Kidney Int Suppl. 2005; 67(94): S133-S135.

383 [50] Afshin, A., Micha, R., Khatibzadeh, S., Fahimi, S., Shi, P., Powles, J. et al. The impact of dietary
384 habits and metabolic risk factors on cardiovascular and diabetes mortality in countries of the
385 Middle East and North Africa in 2010: A comparative risk assessment analysis. BMJ Open. 2015;
386 5: e006385.

387 [51] Mirnalini, K., Zalilah, M.S., Safiah, M.Y., Tahir, A., Siti Haslinda, M.D., et al. Energy and nutrient
388 intakes: Findings from the Malaysian Adult Nutrition Survey (MANS). Mal J Nutr. 2008; 14(1): 1-
389 24.

390 [52] Moy, F.M., Suriah, A.R. Anthropometric and dietary intake of type 2 diabetes patients attending
391 an outpatient clinic. Mal J Nutr. 2002; 8(1): 63-73.

392