

Original Research Articles

The Interaction between Malaria parasites and Blood Groups in Port Harcourt, Nigeria.

ABSTRACT

The link between blood group Rhesus antigens and different malaria parasites as well as the prevalence of other Plasmodium species were studied among adults in Port Harcourt, Nigeria. Thick blood smears and filter paper blood spots were made from finger prick for microscopy and molecular genotyping of parasite strains. Two hundred and forty six participants aged 16-60 years attending the Braithwaite Memorial Hospital and blood donors presenting at the University of Port Harcourt Teaching Hospital Blood Bank were enrolled into the study. Blood samples were analysed for ABO antigen grouping using commercially prepared potent antisera. Isolated DNA was extracted from dried filter papers and analysed for Plasmodium specie identification using nested PCR method. Results obtained reveal that blood group O was most prevalent while the least was blood group AB. Parasitaemia results by microscopy shows that blood group O Rh positive was the highest with 163 (66.2%) followed by blood group A Rh positive 43 (17.5%), B Rh positive 26 (10.6%), O Rh negative 7 (2.85%), AB Rh positive 5 (2.03%), B Rh negative 1 (0.41%) and A Rh negative 1(0.41%). Nested PCR revealed the presence of low prevalence of *P.ovale* and *P.malariae* species (4.3% each) and a very high prevalence of *P. falciparum* (82.1%) as well as mixed infections of the species. These results show the association of blood groups, rhesus positivity and Plasmodium parasites as well as the presence of other species in the study area.

Key Words: malaria parasites, blood group, interaction

1. Introduction

The global burden of malaria infection remains more severe in Africa especially among children and pregnant mother. Malaria is a protozoan infection in humans caused by the parasite Plasmodium which until recently was thought to be classified into five species *P. falciparum*, *P. ovale*, *P. malariae* , *P. vivax* and *P. knwolesi*. However current study ¹ (Sutherland et al., 2010) has shown that there are two non-recombining species of *ovale*

(*ovale curtisi* and *ovale wallikeri*) which are non-sympatric in nature ²(Oguike et al., 2011). Of all these, *Plasmodium falciparum* is the agent of the most malignant form of malaria, usually presenting with severity mostly in children in sub-Saharan Africa.

The pathogenesis of Plasmodium infection entails merozoite invasion of erythrocyte, which implies an implicit interaction between the red cell membrane proteins and the invading plasmodium antigens. Blood group antigens are constituents of these proteins that are embedded in erythrocyte cellular membranes and are encoded by alleles at a single gene locus or at closely linked gene loci of which more than 300 specificities have been reported (Johnson, 1992³; Contreras, 2001)⁴. Additionally, blood group antigens have been reported to serve as genetic markers of several clinical conditions (Nwauche et al 2003, Vestibo et al, 1993, Slater, 1993, Brooks et al, 1988)⁵⁻⁸ while several workers have shown a definite relationship between the ABO and Rh blood group antigens with malaria (Ademowo et al, 1993; Cserti et al, 2007)^{9,10}. The clearest example of this is the interaction between Plasmodium vivax and the Duffy antigen (Langhi & Bordin, 2006; Carlos et al, 2007)^{11,12}.

1.1. Study Area

The study was carried out in two major hospitals: the University of Port Harcourt Teaching Hospital (UPTH) and the Braithwaite Memorial Specialist Hospital (BMH) both in Port Harcourt, Nigeria. The study was conducted from September 2010 to September 2011. Malaria is holoendemic in Nigeria. Nigeria has two main seasons, the dry seasons which occur from October to March and the wet or rainy season occurring from April to September with peak of the rains between May and July when malaria transmission is very intense. Port Harcourt exhibits lengthy and heavy rain seasons with annual rainfall of more than 3,550 millimeters and average temperatures between 25°C-28°C.

The main vector of malaria in Nigeria is *Anopheles gambiae* but *A. funestus* and *A. arabiensis* are also commonly encountered. *A. melas* is found in the coastal areas. (NPC & NMCP, 2012)¹³. The commonest species responsible for to 98% of the confirmed malaria cases in Nigeria is *Plasmodium falciparum*. The other specie that has been reported in Nigeria is the *P. malariae*, but it usually occurs as a mixed infection with *P. falciparum*. (NPC & NMCP, 2012)¹³.

2. Methods

2.1. Study Participants

Two hundred and forty six participants: 142 males (57.72%) and 104 females (42.28%) aged 16-60 years attending the Braithwaite Memorial Hospital and blood donors presenting at the University of Port Harcourt Teaching Hospital Blood Bank were enrolled into the study. Consent was obtained from the participants through the signed informed consent forms. Ethics approval was obtained from the Ethics Committee of both hospitals and the London School of Hygiene and Tropical Medicine.

2.2. Screening for ABO and Rhesus D antigen

Blood donors were sampled and screened for ABO and Rhesus D antigen. The ABO and Rhesus D antigen testing was carried according to previously described method (Dacie and Lewis, 2001¹⁴) using commercially prepared potent antisera (anti-A, anti-B, anti-D). Briefly described, 3mls of blood were transferred to sterilin containers and red cells were separated and suspended in saline. One drop of 20%cell suspension was added to two drops of the antisera (cell to serum) in the following tubes-Saline, albumin and anti-human globulin (AHG) and were incubated at 37°C. Positive samples were identified by agglutination.

Thick blood smears and dry blood spots on Whatman filter papers were made from finger prick for microscopy and molecular genotyping of parasite strains. Participants were recruited if the blood samples collected were found *Plasmodium*-positive by microscopy. Genotyping of the filter paper blood spots specie identification was carried out in the London School of Hygiene and Tropical Medicine, London for identification of *Plasmodium* species.

2.3. Isolation of DNA

Parasite DNA was extracted from dried filter papers using Chelex method as previously reported (Dlamini et al, 2010)¹⁵. Briefly described, blood spots punched out from dried filter papers were soaked in a deep well plate in 0.5% saponin 1 x PBS solution and incubated overnight at 37°C. Debris and saponin were removed after centrifugation and the filter papers were washed in PBS solution. They were subsequently suspended in 6% Chelex®100 and incubated in a water bath for 25 minutes at 100°C in 6% Chelex®100 and centrifuged thereafter to remove resin. Approximately 100µl of supernatant containing the DNA was removed and stored in -20°C freezer.

2.4. *Plasmodium* species identification by nested PCR

Amplification of the small sub-unit of ribosomal gene (SSU rRNA) of the *Plasmodium* species was carried out on extracted DNA using nested PCR to identify the different species present. (Snounou et al., 1993)¹⁶. PCR reactions was carried out under the following conditions: 95°C for 5min/ [58°C for 2min-72°C for 2min-94°C for 1min] x25 cycles/58°C for 2min-72°C for 5min using primers (rPLU5+rPLU6) as listed below.

rPLU6	5'	TTAAAATTGTTGCAGTTAAAACG	3'genus F2
rPLU5 new	5'	CYTGTTGTTGCCTTAAACTTC	3'genus R

The amplified products were nested with primer mix consisting of 5 μ M Primer mix (Pf/Pm/Pv/Po) under the cycling conditions– 95°C for 5mins, (58°C for 2mins; 72°C for 2mins; 94°C for 1min) x30cycles; 58°C for 2mins, 72°C for 5mins, 4°C forever for species differentiation using Snounou protocol¹⁸. The amplified products were run on agarose gel and size against Hyperladder IV to identify the different species. PCR conditions and primer sequences used are listed below.

rFAL1 5'-TTAAACTGGTTTGGGAAAACCAAATATATT-3'

rFAL2 5'-ACACAATAGACTCAATCATGACTACCCGTC-3'

rMAL1 5'-ATAACATAGTTGTACGTTAAGAATAACCGC-3'

rMAL2 5'-AAAATTCCCATGCATAAAAAATTATACAAA-3'

rVIV1 5'-CGCTTCTAGCTTAATCCACATAACTGATAC-3'

rVIV2 5'-ACTTCCAAGCCGAAGCAAAGAAAGTCCTTA-3'

Pad Po 5' CTGTTCTTTGCATTCTTATGC 3'

Cycling conditions – 95°C for 5mins, (58°C for 2mins; 72°C for 2mins; 94°C for 1min) x30cycles; 58°C for 2mins, 72°C for 5mins, 4°C forever. The band size obtained for the various species differ and each was identified according to the band size. The band sizes obtained are here below listed.

Table 1: Plasmodium species with the expected band sizes

P. falciparum primers(rfal1+rfal2)	205 bp product
P. malariae (rmal1 + rmal2)	144bp
P. vivax (rviv1+rviv2)	120bp
P. ovale (pad Po)	800bp

3. Results

Observation from the baseline distribution of blood groups shows that the largest blood group in the set of study samples was blood group O (Fig.1). Results of the blood group screening against microscopy for *Plasmodium falciparum* shows that blood group O Rh positive was the highest with 170 (69.1%) followed by blood group A 44 (17.9%), B 27 (11%), AB 5 (2%) [Fig. 2]. Close observation of the results also show that the samples from Rhesus positive were more parasitaemic than the Rhesus negative samples Table 2.

From the analysis of the PCR data obtained from the specie identification, we found a similar pattern of distribution of the parasitemia among the blood groups with Blood group O having the highest number of parasitaemia positive. However, we observed that a very high percentage (71.5%) of the samples were aparasitaemic suggesting a very poor microscopic diagnosis. Further observation from the obtained PCR positive results, showed that the samples with Rhesus positivity had higher parasitaemia (Table 2). Analysis of the specie identification from the PCR results, revealed that the most prevalent specie is *P. falciparum* (82.9%), followed by *P. malariae* (5.7%), *P. ovale* (5.7%) and then mixed infections of *falciparum* and *ovale* (1.4%), *falciparum* and *malariae* (4.3%). Closer observation showed that the few *ovale* and *malariae* parasites seen were predominantly in the blood group O Rhesus positive samples while 25% (1/4) of each of these parasites were from the blood group B Rhesus positive samples. The mixed infections were all from the blood group O Rhesus positive samples. There was no *P. vivax* seen in all the samples analysed (Table 2).

Table 2. Distribution of Plasmodium species and ABO-Rh antigens using PCR

Bld Gp	A+	A-	B+	B-	AB+	AB-	O+	O-	Total
Pf	11		3				40	4	58
Pm			1				3		4
Po			1				3		4
Pf+Pm							3		3

Pf+Po							1		1
Total	11		5				50	4	70

Fig. 1 distribution of blood group among the study subjects.





Fig. 2 Pie chart showing *P. falciparum* parasitaemia by microscopy and blood group distribution



4. Discussion

A total of 246 adults presenting blood donors were recruited into the study. Results obtained showed that the most prevalent blood group was Blood group O while the least was blood group A. This is in consonance with earlier studies (Jeremiah, 2006, Nwauche et al, 2012, Otajevwo and Igoniwari, 2014)¹⁷⁻¹⁹ and the much earlier study on nationwide distribution of ABO blood group as reported by Worlledge et al, 1974²⁰.

Analysing the results of the Rhesus positivity, the observation was that Plasmodium parasite selects for the Rhesus D antigen hence the samples which were Rhesus positive had higher prevalence of parasitaemia regardless of the ABO blood grouping. The combined analysis of the blood grouping and Rhesus antigen, against microscopy revealed that blood group O Rh positive had the highest parasitaemia positive while the least was seen in the blood group A Rhesus negative. This observation is in consonance with previous reports of Singh et al, 2015²¹ of higher prevalence of malaria among Rhesus positive persons.

Further findings of the study reveal that the most prevalent specie in the study sample is *Plasmodium falciparum*. This is consistent with various studies (^{22,23}) that have reported the high prevalence of *P. falciparum* in Nigeria. National Malaria control (NMC, 2012)¹³ reports that *P. falciparum* is the most virulent of the Plasmodium species and accounts for about 98% of the malaria cases in the country. In malaria endemic areas of tropical Africa, almost all malaria attacks are attributed to *P. falciparum*, possibly due either to underdiagnosis of *P. ovale* and *P. malariae* clinical attacks, to partial cross-immunity between malaria species, or to rapidly acquired species-specific protective immunity against *P. ovale* and *P. malariae* (Roucher et al, 2014)²⁴. The presence of these species *seen* though with a very low prevalence is in consonance with the report from Malaria Indicator Survey (2010)¹³ and it buttresses the importance of further studies on the prevalence of other species of Plasmodium in Nigeria.

In a 20 year study in Senegal, Roucher et al, 2014²⁴ established that *Plasmodium ovale* and *malariae* were a common cause of morbidity in the villagers. This underlines the importance of surveillance to these parasites which have been overshadowed by the abundant presence of falciparum malaria and understanding these interactions could have an important influence on the deployment of interventions such as malaria vaccines (Mueller et al, 2007)²⁵.

Plasmodium ovale was the last of the malaria parasites of humans to be described and it is primarily distributed throughout sub-Saharan Africa (Collins and Jeffrey, 2005)²⁶. The presence of *P. ovale* was first reported in Nigeria in 1930 (James et l, 1930)²⁷. It is important that the diagnosis of ovale malaria be made appropriately to enhance proper and accurate management since *Plasmodium ovale* is a relapsing malaria parasite with a latent liver stage that often persists for many months. There have been reports of relapses occurring as long as 45 months after infection (Collins and Jeffrey 20015, Marty et al, 1987)^{26,28}. On the other

hand, *Plasmodium malariae* has been reported to carry a high rate of mortality both in children and adults (Mueller et al, 2007²⁵) and have been attributed to be responsible for chronic nephrotic syndrome in a US traveler 14 years after leaving Nigeria.(Hedelius et al, 2011)²⁹.

A major limitation to the study is the inexperience of the microscopists such that a greater percentage of the study participants were actually aparsitaemic. This resulted in lower number that would be used for statistical analysis. Despite this limitation, we were able to present a picture of the interaction between ABO Rh blood group and Plasmodium species in the study environment. Further studies with a larger sample size is therefore very imperative

In conclusion, this study is first to document the interaction between different Plasmodium species and ABO antigen blood groups in Nigeria. The study showed that there is higher prevalence of malaria parasitaemia among blood group O Rhesus positive persons regardless of the Plasmodium specie thereby underscoring the relationship between blood group O antigen, Rhesus positivity and parasitaemia. It also highlighted the presence of mixed infections in Port Harcourt and its relationship with blood group O Rhesus positive. There is need for further investigations of the interaction between blood groups and various malaria parasites since this may be beneficial in malaria vaccine development.

References

1. Sutherland CJ, Tanomsing N, Nolder D, Oguike M, Jennison C, Pukrittayakamee S et al. Two nonrecombining sympatric forms of the human malaria parasite *Plasmodium ovale* occur globally. *J Infect Dis*, 2010, 1544-50.
2. Oguike MC, Betson M, Burke M, Nolder D, Stothard JR, Kleinschmidt I et al. *Plasmodium ovale curtisi* and *Plasmodium ovale wallikeri* circulate simultaneously in African communities. *Int J Parasitol*, 2011; 41, 677-83.
3. Johnson CL. Blood group antigens and antibodies. In: Tilton R., Balows A, Hohnadel DC, Reiss RF. (Eds). *Clinical Laboratory Medicine*. St. Louis. Mosby Yearbook Inc. 1992. Pp. 1064-1074.

4. Contreras, Lubenko A. Antigens in human blood. In: Hoffbrand AV, Lewis SM, and Tuddenham EGD (Eds) Postgraduate Haematology 2001. Fourth Edition, Arnold, pp182-214.
5. Nwauche CA and Ejele OA. Red cell antigens and the practice of transfusion medicine in Nigeria. Nigeria Journal of Orthopaedics & Trauma, 2003; vol 2(2), 68-7.
6. Vestbo JHO, Suadican P, Sorensen H, Gyntelberg F. The genetic markers for Chronic Bronchitis and Peak expiratory flow in the Copenhagen male study. Danish Med. Bull. 1993, 40 (3): 378-80.
7. Slater G, Itzkowitz S, Azer A.H. (Jr). Clinicopathologic correlations of ABO and Rhesus blood type in Colorectal cancer. Dis Col Rect, 1993, 36(1): 5-7.
8. Brooks A.M. Gilles WE. Blood groups as genetic markers in Glaucoma. J Ophthal, 1988, 72(4):2.
9. Ademowo OG, Falusi AG and Mewoyeka OO. Prevalence of asymptomatic parasitemia in an urban and rural community in South-Western Nigeria. Cent Afr J Med 1993; 41 (1) : 18-21.
10. Cserti CM and Dzik WH. The ABO blood group system and *Plasmodium falciparum* malaria. BLOOD, 2007; 10(7): 2250-8.
11. Langhi DM Jr, Bordin JO. Duffy blood group and malaria. Hematology. 2006 Oct; 11(5):389-98.
12. Carlos E Cavasini Luiz C de Mattos, Álvaro AR D'Almeida Couto, Vanja SC D'Almeida Couto, Yuri Gollino, Laurence J Moretti, Cláudia R Bonini-Domingos, Andréa RB Rossit, Lilian Castilho and Ricardo LD Machado. Duffy blood group gene polymorphisms among malaria vivax patients in four areas of the Brazilian Amazon region Malaria Journal 2007, 6:167.
13. National Population Commission & National Malaria Control Programme (NPC & NMC). 2012. Nigeria Malaria Indicator Survey, 2010 Final Report, Federal Republic of Nigeria
14. Dacie JV, Lewis SM. Practical Haematology, 9TH Ed. London: Churchill Livingstone, 2001: 19-56.
15. Dlamini SV, Beshir K. & Sutherland CJ. Markers of anti-malarial drug resistance in *Plasmodium falciparum* isolates from Swaziland: identification of pfmdr1-86F in natural parasite isolates. Malar J, 2010, 9, 68.

16. Snounou G, Viriyakosol S, Zhu XP, Jarra W, Pinheiro L, Do-Rosario VE, Thaithong S, & Brown KN. High sensitivity of detection of human malaria parasites by the use of nested polymerase chain reaction. *Mol Biochem Parasitol*, 1993 61, 315-20.
17. Jeremiah ZA. Abnormal haemoglobin variants, ABO and Rh groups among students of African Descent in Port Harcourt. *Africa. Health Sci.* 2006; 6:177-81
18. Nwauche CA, Eneh AU, Pughikumo CO, Chijioke-Nwauche I. ABO and Rhesus blood groups in mothers and their newborns in Port Harcourt, Nigeria. *Port Harcourt Medical Journal* 2012; 6 (4), 455-459.
19. Otajevwo FD and Igoniwari SF. Malaria Parasitaemia Association with ABO. Blood Types among Students of a Private University in Western Delta, Nigeria. *International Journal of TROPICAL DISEASE & Health* 4(5): 540-554, 2014
20. Worlledge S, Ogiemudia SE, Thomas CO, Ikoku BN, Luzzatto L. Blood group antigens and antibodies in Nigeria. *Ann. Trop Med Parasitol*; 1974; 68: 249-264
21. Singh G, Urhekar AD, Singh R. A study on correlation on malaria infection with A, B, O and RH blood group system. *Journal of Parasitology and Vector Biology*. 2015. Vol.7 (4), 67-73
22. Nigeria Malaria Indicator Survey (2010)
23. Okolie NJ. Occurrence of different Plasmodium species in malaria patients in Owerri, Nigeria. *International Journal of Natural and Applied Sciences* Vol. 2 (4) 2006: pp. 409-411.
24. Roucher C, Rogier C, Sokhna, Tall A, Trape J. A 20-Year Longitudinal Study of Plasmodium ovale and Plasmodium malariae Prevalence and Morbidity in a West African Population. *PLOS ONE*. 2014, 9(2): 1-13.
25. Mueller I, Zimmerman PA, Reeder JC. Plasmodium malariae and Plasmodium ovale - the “bashful” malaria parasites. *Trends Parasitol*; 2007 23: 278–283.
26. Collins WE and Jeffrey GM. *Plasmodium ovale*: parasite and disease. *Clin Microbiol.* 2005 18(3): 570–81.
27. James SP, Nicol WD, Shute PG. The specific status of Plasmodium ovale Stephens. *Am. J. Trop. Med.* 1935; 15:187-88.
28. Marty PB, Chapdelaine Y. Le Fichoux, and J. M. Chabert JM. . Paludisme anémiant à *Plasmodium ovale* après 45 mois d'incubation. *Presse Med.* 2017, 16:357.
29. Hedelius R, Fletcher JJ, Glass WF, Susanti A, Maguire JD. Nephrotic syndrome and unrecognized Plasmodium malariae infection in a US Navy sailor 14 years after departing Nigeria. *J Travel Med*; 2011; 18:288–291.