

**Original Research Articles****The Synergy of Diet, Exercises and Lifestyle Modification for the Prevention of Osteoporosis****Abstract:**

This paper did a review of literature on osteoporosis, the risk factors, the consequences, and the prevention through the synergy of diet, exercises and appropriate lifestyle modification. Osteoporosis is a condition in which bones become dangerously thin and fragile over time. Fractures are the most serious consequence of osteoporosis, and up to 20% of all people who suffers a hip fracture die within a year. Other problems associated with osteoporosis are loss of height and a stooped posture due to vertebral fractures, severe back and hip pain, and breathing problems caused by changes in the shapes of the skeleton. Women are at greater risk than men for osteoporosis because they have 10 – 25% less bone in their skeletons. Women's bones become dangerously thin sooner than men's bones after menopause when estrogen production is diminished. Estrogens improves calcium absorption and reduces the amount of calcium the body excretes. Preventing osteoporosis requires building as much bone as possible during ones young years and then maintaining it as one age. Diet and exercises also play key roles. Other lifestyle modifications, such as abstaining from cigarette smoking, and alcohol consumption one imperative in preventing osteoporosis.

26 **Key Words:**

27 Osteoporosis, Diet, Exercise, Fractures, Estrogen, Menopause,  
28 Calcium, Copper.

29 **Introduction:**

30 Osteoporosis (thinning and weakening of the bones related to  
31 loss of calcium stores) is a debilitating disease that is primarily age  
32 – and gender – related. It is also described as a condition in which  
33 bones become dangerously thin and fragile overtime [1].  
34 Osteoporosis is a disorder characterised by a reduction in bone  
35 density due to calcium loss and resulting in brittle bones. A  
36 National Institute of Health [2]. Consensus panel noted that  
37 although prevalent in white menopausal women, osteoporosis  
38 occurs in all populations and at all ages and has significant  
39 physical, psychological, and financial consequences.

40 Fractures are the most serious consequence of osteoporosis up  
41 to 20% of all people who suffer hip fractures die within a year.  
42 Other problems associated with osteoporosis are loss of height and  
43 a stooped posture due to vertebral fractures, severe back and hip  
44 pain, and breathing problem caused by changes in the shapes of  
45 the skeleton [1].

46 Osteoporosis occurs more frequently in women than in men  
47 [3]. Women are at greater risk than men for osteoporosis because  
48 they have 10 – 25% less bone in their skeleton. As they lose bone  
49 mass with age, women's bones become dangerously thin sooner

50 than men's bones [1]. Bone loss accelerates in women during the  
 51 first 5 – 10years after the onset of menopause because of the drop  
 52 in estrogen production. Estrogen improves calcium absorption and  
 53 reduces the amount of calcium the body excretes.

54 Although osteoporosis is more common in women, it is also  
 55 common among men. More than 2million American men suffer from  
 56 osteoporosis and millions more at risk [3]. Each year 80,000 men  
 57 suffer a hip fracture due to this disorder [3]. Osteoporosis does not  
 58 begin in old age. The bone weakness that increases the risk of  
 59 osteoporosis may actually begin before birth. Infants  
 60 undernourished in the womb are small when born, remain smaller  
 61 than their age peers, and have low bone mass at age 25, and have  
 62 an increased risk of fracture in late adulthood [4]. Other risk factors  
 63 for osteoporosis are diet, lack of exercise, and lifestyle [4], [1], [3].  
 64 According to them, a positive family history, or heredity, and low  
 65 levels of estrogen and the two primary risk factors in women.  
 66 Following menopause, estrogen production is diminished. Estrogen  
 67 is a hormone essential for optimal calcium balance in women.  
 68 Reduced levels of estrogen lead to negative calcium balance and a  
 69 rapid onset of bone demineralization. This softening of the bones  
 70 predisposes to fractures, particularly in the spine, the end of the  
 71 radius in the forearm, and the neck of the femur at the hip joint.

72 ➤ An estimated 10million Americans over age 50 have  
 73 osteoporosis, and another 34millioin are at risk [4] and many

more are in the developing nations of the world where the health system is weak, and where poor vital statistics abound.

➤ [3] And [1] affirmed that most bone mass is built by age 18, and after, bone density peaks between ages 25 and 35, and that bone mass is slowly lost over time. So to prevent osteoporosis, the best strategy is to build as much bone as possible during one's youth and then to do everything one can to maintain it as you age. Since up to 50% of bone loss is determined by controllable lifestyle factors, especially diet and exercise habits, this paper reviewed the synergy of diet, physical exercises, and lifestyle modification for the prevention of osteoporosis.

### **Causative Factors of Osteoporosis**

Osteoporosis does not begin in old age. The bone weakness that increases the risk of osteoporosis may actually begin before birth. Infants undernourished in the womb are small when born, remain smaller than their age peers at age 1, have low bone mass at 25, and have an increased risk of fracture in late adulthood [3]. Other risk factors of osteoporosis according to [3] include:

- ❖ Age; Risk of osteoporosis increases as one grows older.
- ❖ Being Female: women have less bone tissue than do men and tend to experience a rapid loss of bone in the first few years after menopause.

❖ Body Size-: Small, thin – boned women are at greater risk of osteoporosis.

❖ Ethnicity: white and Asian women are at highest risk.

❖ Family History: Having parents with a history of osteoporosis as well as fractures in adulthood can place individuals at increased risk.

❖ Sex Hormones: Abnormal absence of menstrual periods (amenorrhea) or menopause can increase risk of osteoporosis.

❖ Anorexia

❖ Lifetime diet low in calcium and vitamin D.

Other risk factors according to [1] include: early menopause (before age 45), abnormal or irregular menstruation, a history of anorexia, and a thin small frame. Other risk factors are thyroid medication, corticosteroid drugs for arthritis or asthma, and long-term use of the contraceptive Depo-Provera.

Similarly, Booth and Lees, according to [4] indicated that inactivity (lack of exercise) increase the risk of osteoporosis by 59%. Also several dietary substances may have a negative effect on bone health. Alcohol consumption reduces the body's ability to absorb calcium and may interfere with the bone-protecting effects of the hormone estrogen. High sodium intake increases calcium loss in the urine, and this may lead to loss of calcium from the skeleton. Caffeine may also cause small losses of urinary calcium.

Furthermore, chronic excess intake of retinal, one form of vitamin A, is associated with decreased bone density. Drinking lots

123 of Soda, which often replaces milk in the diet and which is high in  
124 phosphorous (a mineral that may interfere with calcium absorption)  
125 has been shown to increase the risk of bone fracture in teenage  
126 girls [4].

## 127 **Deleterious Consequences of Osteoporosis**

128 Osteoporosis (thining and weakening of the bones related to  
129 loss of calcium stores), is a debilitating disease that is primarily  
130 age- and gender-related. A [2] of Health (NIH) consensus panel  
131 noted that although prevalent in white post menopausal women,  
132 osteoporosis occurs in all populations and at all ages and has  
133 significant physical, psychological and financial consequences.[4]  
134 observed that each year, osteoporosis causes 1.5 million fractures,  
135 including 300,000 hip and 700,000 vertebral fractures. Although  
136 more common in women, more than 2 million American men suffer  
137 from osteoporosis, and millions more at risk. Each year, 80,000  
138 men suffer a hip fracture. Osteoporosis is known as a silent killer.  
139 The disease itself causes no pain but following a serious bone  
140 fracture, nearly a third or more of women and men die due to  
141 accompanying illness within a year.

142 Also, osteoporosis may lead to a condition in which losses in  
143 bone density become so severe that a bone will break after even  
144 slight trauma or injury. A chronic disease, osteoporosis is silent for  
145 years or decades before a fracture occurs. Women who have smaller  
146 skeletons, are more vulnerable than men; in extreme cases, their  
147 spines may become so fragile that just bending causes severe pain.

148 But although commonly seen as an illness of women, osteoporosis  
149 occurs frequently in men. One in every two women and one in four  
150 men over 50 will have an osteoporosis-related fracture in their life-  
151 times [3].

152 Furthermore, experts agreed that fractures are the most  
153 serious consequences of osteoporosis; and that up to 20% of all  
154 people who suffer a hip fracture die within a year[4], [1], [3]. Also,  
155 other problems associated with osteoporosis are loss of height and  
156 stooped posture due to vertebral fractures, severe back pain, hip  
157 pain, and breathing problems caused by changes in the shape of  
158 the skeleton [1]. Severe breathing problems may also occur. Severe  
159 breathing problems may lead to short of breathe, lack of oxygen to  
160 the systems and organs of the body that are very vital, and  
161 complications in the cardiovascular and nervous systems may  
162 ensue.

163 In addition to the morbidity and mortality, osteoporosis may  
164 lead to many man-hour losses due to absence from duty. Huge sum  
165 of money are expended in the treatment and management of this  
166 disease at individual level, besides cost of drugs purchase by the  
167 government, in addition to reduced productivity.

## 168 **Prevention of Osteoporosis**

169 Preventing osteoporosis requires building as much bone as  
170 possible during the young years and then maintain it as you age.  
171 Diet and exercise play key roles. Diets containing calcium, vitamin

172 D, and vitamin K are very useful in preventing osteoporosis. Also  
 173 weight bearing aerobic activities must be performed regularly  
 174 throughout life to have lasting effects. Strength training improves  
 175 bone density, muscle mass, strength, and balance, protecting  
 176 against both bone loss and falls, a major cause of fractures.

177 Both men and women can benefit from high impact exercises,  
 178 such as running and jumping. Studies by [5] and [6] indicated that  
 179 young, premenopausal women had significant higher bone mineral  
 180 density; and that moderate exercise reduces hip fracture in both  
 181 men and women by about 20%. Two other lifelong strategies for  
 182 reducing the effects of osteoporosis are avoiding tobacco and  
 183 managing depression and stress. Smoking reduces the body's  
 184 oestrogen levels and is linked to earlier menopause and more rapid  
 185 post menopausal bone loss. In addition, alcohol reduces the body's  
 186 ability to absorb Calcium and may interfere with the bone-  
 187 protecting effects of the hormone oestrogen. Therefore, smoking and  
 188 alcohol consumption must be avoided in order to prevent  
 189 osteoporosis.

## 190 **Osteoporosis and Diet**

191 Up to 50% of bone loss is determined by controllable lifestyle  
 192 factors, especially diet and exercise habits [1]. Key nutrients  
 193 essential for building and maintaining bone mass, and which must  
 194 be factored into individual's diet include the following;

1. Calcium-: [1] affirmed that consuming an adequate amount of Calcium is important throughout life to build and maintain bone mass. Milk, yogurt, and calcium fortified orange juice, bread and cereals are all good sources of calcium [7] [8]. Nutritionist suggests that one can obtain calcium from food first and then take supplements only if needed to make up the difference [9]. Another source of calcium in diet is through insects consumption, particularly vonocorous variegates, and *Migratoria locusta* [1]. Insects are known to supply many minerals, most of which are essential for building and maintain bone mass.
2. Vitamin D-: Vitamin D is necessary for bones to absorb calcium. It has been suggested by [1] that a daily intake of 5ug is recommended for adult age 19-50. Vitamin D can be obtained from foods and is manufactured by the skin when it is exposed to sunlight. Candidates for vitamin D supplements include people who do not eat many foods rich in Vitamin D; those who have dark skin or who do not expose their face, arms, and hands to the sun for 5-15 minutes a few times each week; and people who live in Northern latitudes of Boston, Oregon-California border where the sun is weaker [4].
3. Vitamin K-: Vitamin K promotes the synthesis of proteins that help keep bones strong. Broccoli and leafy vegetables are rich in Vitamin K. Good plant sources of vitamin K include vegetable oils (Soya beans, olive), green and leafy vegetable such as Kale, peas, and spinach; while meats, and milk

contain much lower amounts. Since vitamin K appears to enhance the function of osteocalcin, a protein that plays an important role in strengthening bones [4], consumption of vitamin K is recommended for the prevention of osteoporosis.

4. Vitamin C-: Vitamin C plays an important role in bone health. Vitamin C with calcium and other minerals to build bone; it also helps to reduce the connective tissue collagen, which forms the scaffolding in bones. Vitamin C is obtained from food sources such as fruit and vegetables, primarily the Citrus fruits and leafy parts of green vegetables. Excellent sources include oranges, grape fruit, broccoli, and salad greens. Other food sources are green peppers, potatoes, strawberries and tomatoes. Some products are fortified with vitamin C and they could also be obtained from supplements.

5. Magnesium-: Magnesium aids in bone formation. The body stores about 50-60% of its magnesium in the skeletal system, which may serve as a reserve during short periods of dietary deficiency. [10], indicates that magnesium influences bone metabolism and helps prevent bone fragility. Sources of magnesium include nuts, seafood, green leafy vegetables, fruit vegetables, black beans, and whole-grain products.

6. Potassium-: Potassium helps bones retain calcium [1]. Potassium is found in most foods and is especially abundant in bananas, citrus fruits, fresh vegetables, milk, meat and fish.

7. Manganese-: Manganese may help lessen calcium loses. Apart from the function of Manganese in many enzymes involved in

energy metabolism, manganese is also involved in bone formation. Sources of manganese include whole grain products, dried peas and beans, leafy vegetables and bananas.

8. Fluoride: Fluoride helps form bones and teeth. Sources of fluoride include milk, egg yolks, drinking water and seafoods .

9. Zinc: Zinc, according to [11] helps in bone formation, and wound healing. Good sources of Zinc are found in animal protein, such as meat, milk and seafood such as Oysters.

10. Copper: Copper is essential in helping maintain collagen. [12], also indicate that copper is involved in bone formation, being an essential cofactor in enzyme involved in the synthesis of various bone matrix constituents. Copper is widely distributed in foods and is high in seafoods, meat, nuts, beans, and whole grain products.

11. Boron: Boron may increase Calcium absorption. Boron is known to be involved in the metabolism of steroid hormones, such as estrogen. [13] Indicated that the effects of boron on calcium and estrogen metabolism could favourably affect bone development. Boron is found naturally in many plant foods, particularly dried fruits, nuts, peanut butter, legumes, fresh vegetables, apple source, milk and dairy products, grape juice and wine.

## **Exercise and osteoporosis**

The beneficial effects of exercise on health has been known for centuries. [14] and others discussed the cost of physical inactivity overtime. The shorter-term cost of physical inactivity is metabolic deterioration and weight gain; the intermediate-term cost is an increase in disease, such as type 2 diabetes, and the long-term cost is an increased premature mortality. [4] affirmed that physical inactivity increases osteoporosis by 59%. Clinical, epidemiological and basic research evidence clearly supports the inclusion of regular physical activity as a tool for the prevention of chronic disease, including osteoporosis and the enhancement of overall health. Specifically, [4] asserted that regular moderate physical activity and exercise, among others, increases bone density and strength, thus preventing osteoporosis. In the same vein, [3] opined that exercise, both aerobic exercise and weight training can help preserve bone density.

In addition, [15] concluded that physical activity offers strong protection against osteoporosis in postmenopausal women but is less effective in preventing this disorder in premenopausal women. Furthermore, evidence has accumulated suggesting that physical activity can protect both men and women against loss of bone mineral density, especially if they have a history of athletic performance. To this end, weight-bearing aerobic activities must be performed regularly throughout life to have lasting effects on osteoporosis prevention. Strength training improves bone density, muscle mass, strength and balance,

protecting against both bone loss and falls, a regular cause of fractures. Even for people in their seventies, low-intensity strength training has been shown to improve bone density.

### **Osteoporosis and Lifestyle Modification:**

Lifestyle modification in the case, means doing certain things you have not been doing but which promotes your health and abandoning things which are detrimental to your health. Lifestyle modification is very pertinent in preventing osteoporosis. One type of healthy lifestyle modification is the participation in physical activities. Clinical, epidemiological and basic research evidence clearly supports the inclusion of regular physical activity as a tool for the prevention of chronic disease, including osteoporosis and the enhancement of overall health. Specially, [4] affirmed that regular moderate physical activity and exercise, among others increases bone density and strength, thus preventing osteoporosis. In view of this, [3] suggests that that one should exercise regularly and that both aerobic exercise and weight training can help preserve bone density. The values of physical activity may have informed the [16] and the American Medical Association [AMA] to launch a programme entitled “Exercise is Medicine” designed to encourage physicians and other health care professionals to include exercise as part of the treatment for every patient.

Another lifestyle modification that may help in preventing osteoporosis is ceasation of cigarette smoking. Cigarette smoking has many deleterious health consequences, including a risk factor for osteoporosis. Smoking cigarette reduces the body's estrogen levels and is linked to earlier menopause, and more rapid postmenopausal bone loss [1]. Also, [3] affirmed that smokers tend to be thin and enter menopause earlier, thus extending the period of jeopardy from estrogen loss. But since estrogen improves calcium absorption and reduces the amount of calcium the body excretes, smoking cigarette should be avoided so as to preserve the body's estreogen levels.

### **Conclusion**

- This paper did a review of literature on osteoporosis, the risk factors, the deleterious consequences, and preventive measures through the synergy of diet, physical exercises and appropriate lifestyle modification, and therefore concludes as follows;
1. Osteoporosis affects both males and females, but the occurrence is greater among women than men.
  2. Osteoporosis may begin among undernourished infants in the womb.
  3. Occurrence of osteoporosis is faster among women shortly after menopause when estrogen secretion is diminished, and calcium is depleted.

4. Osteoporosis may be prevented through physical exercises, diet rich in calcium, Copper and vitamin D; and through ceasation of cigarette smoking and alcohol consumption.

### **Recommendation**

In corollary to the conclusion, the following recommendations were made;

1. Everybody should be encouraged to participate in physical exercises at youth to build up bone density as a reservoir to draw from in later years.
2. Since osteoporosis is much more common among women, and the risk factors include post menopause when estrogen secretion is diminished, diets rich in Calcium, Copper and Vitamin D should be taken by women, particularly the postmenopausal women.
3. To prevent osteoporosis, ingestion of alcohol, and cigarette smoking should be avoided.
4. Pregnant women should be placed on balanced diet to prevent infant undernourishment in the womb.
5. Individuals should regularly go for medical check-up to examine their bone tissue levels.

### **References**

- [1] Insel, P.M., and Roth,W.T (2006) Core Concept in Health, 10<sup>th</sup> ed. New York, Mc Graw Hill

- [2]National Institutes of Health. (2006). National Institutes of Health State-of-the-Science Conference Statement: Multivitamin/mineral supplements and chronic disease prevention. *Annals of Internal Medicine*. 145:364-71
- [3]Hales, D.(2009). *An Invitation to Health*, 2009-2010 ed, United States Wadsworth.
- [4]Williams, M.H; Anderson, D.E and Rawson, E.S (2013). *Nutrition for Health, Fitness & Sport*. 10<sup>th</sup> ed. New York, MC Graw Hill
- [5] Vainionpaa, A; Korpelainen, R; Leppaluoto, J; And Jamsa, T (2005) Effects of high-impact exercise on bone mineral density: A randomised controlled trial in premenopausal women. *Osteoporosis international*. 16, 191-197
- [6] Holdrup, S; Sorensen, T.I.A; Stronger, U; Lauritzen, J.B; Schroll,M; and Gronbaak, M (2001). Leisure-time physical activity levels and changes in relation to risk of hip fracture in men and women. *American Journal of Epidemiology*. 154,60-68
- [7]Ajala, J.A. (2006). *Understanding Food and Nutrition*. Ibadan, May Best Publishers.
- [8]Alade, Ibukun-Olu (2001). *Public Health Nutrition* (2<sup>nd</sup> ed). Ilorin, S.O.A Fasco Ventures Press
- [9]Talwar, G.P; Srivastava, L.M; and Mudgil, K.D. (1989). *Textbook of Biochemistry and Human Biology*. Prentice Hill of India Private Limited, India

- [10]Sojka, J; and Weaver, C(1995). Magnesium supplementation and osteoporosis. Nutrition Reviews. 53:71-80
- [11]Micheletti, A; et al (2001) Zinc status in athletes: Relation to diet and exercise. Sports medicine 31:577-82
- [12] Lowe, N; et al (2002). Is there a Potential therapeutic value of copper and zinc for osteoporosis? Proceedings of the Nutrition Society.61:181-85
- [13] Nielsen, F. (2008) Is boron nutritionally relevant? Nutrition reviews.66:183-91
- [14]slentz, C; et al (2007). Modest exercise prevents the progressive disease associated with physical inactivity. Exercise and Sports Sciences Reviews 35:18-23
- [15]Harris, S.S; Casperson, C.J DeFries, G.H; and Estes, H. (1989). Physical activity counselling for healthy adults as a primary preventive intervention in the clinical setting:Report for the U.S preventive Services Task Force. Journal of the American Medical Association, 261,3590-3598
- [16]American College of Sports Medicine (2004). Position stand: Physical activity and bone health. Medicine and Science in Sports and Exercise. 36(11). 1985-1996.