

Original Research Article

Human Papilloma virus awareness and uptake of HPV vaccination among human immunodeficiency virus positive patients in Nigeria.

Abstract

Background. The contribution of Human Papillomavirus (HPV) to the aetiology of cervical cancer is well established. Vaccination of at risk population is recommended best practice.

Aim: To determine the awareness of HPV and uptake of HPV vaccination among Human immunodeficiency virus (HIV) positive women attending the antiretroviral clinic.

Study design: A cross sectional questionnaire based survey

Place and duration of study: Antiretroviral clinics in the University of Port Harcourt Teaching Hospital between 21st October -21st of December 2014.

Method: Questionnaire based survey of 162 HIV positive patients was conducted. Data on sociodemographic characteristics, awareness of HPV infection and previous uptake of HPV vaccination was collated and analyzed using SPSS Version 20. Data was presented as frequency, percentages and $P < 0.05$ set as significant.

Results. The mean age of respondents was 35.17 ± 7.14 years. Fourteen percent (23) were aware of HPV causing cervical cancer with 10.4%(17) aware of the existence of HPV vaccination. None of the patients were ever vaccinated against HPV. The main reason for lack of uptake was non-availability of the HPV vaccine.

Conclusion. Awareness of HPV as a precursor of cervical cancer and HPV vaccine uptake among this high risk population is abysmally Poor. There is an urgent need for more sensitization of this group of patients to embrace preventive strategies to reduce the scourge of cervical cancer while concerted efforts should be made to increase availability of vaccine.

Introduction.

Eighty percent of cervical cancer related burden rest in developing countries of the world and it contributes to a significant proportion of all gynaecological oncology deaths [1, 2]. This had been associated with poor health seeking behaviour, late presentation and the lack of use of preventative strategies related to cervical cancer and precursors [3-5].

The link between oncogenic strains of human papillomavirus as the precursors of cervical cancer had been well established by several series [6,7]. A woman of reproductive age has a 50-90% risk of acquisition of the HPV virus during her reproductive lifespan with different oncogenic potentials [8,9]. These are largely categorized into low risk strains associated with development of warts and high risk strains which have been associated with the development of cervical cancer [1,6,7]. There are over 30 variants of HPV which target the genital mucosa with 15 strains considered as oncogenic [6].

31 Globally the commonest oncogenic serotypes are 16 and 18 while others like 31,33,35,45,52,57,58,66,67,69,70,73 and 82
32 had also been identified [1,7,10].

33 Spontaneous regression of HPV infection however, is the norm in a patient with normal cellular immune response without
34 integration of the HPV virus into the cellular genome of the cervix[11].

35 About 50% of women do not develop any measurable antibody response following natural infection with the virus however
36 in women who develop detectable antibody, levels are too low to confer any suitable immunity against reinfection [6,12].

37 HPV vaccine introduction marks a landmark achievement in the primary prevention of cervical cancer owing to the public
38 health challenge cervical cancer had caused in our environment. Its introduction and usage had contributed significantly in
39 reducing the incidence of cervical cancer related morbidities among the immunized group of females in the developed
40 countries [7,13]. but reports of poor uptake had been documented, influenced by sociocultural, political and economic
41 reasons in developing countries[14,15].

42 Globally two vaccines against the HPV are available: Cervarix, a bivalent vaccine produced by GlaxoSmithKline
43 (Brentford UK), against serotypes 16,18 and Gardasil which is a quadrivalent vaccine (Merck and Co USA) protective
44 against serotypes 6,11,16, and 18 which are administered over six months period[13]. The exact duration of antibody
45 protection from immunization is unknown but immunological studies had observed a 11-13 fold increase above natural
46 immunity. However reports of up to 7.3years antibody detection after vaccination with cervirex had been documented with
47 statistical models projecting an up to 10 years protection[16,17].

48 The persistence rate of the HPV virus in patients with immunological compromise is higher with eventual integration of
49 the HPV genome into the cervical epithelial matrix resulting in causation of premalignant transformation [18,19]. Patients
50 in this category, such as the human immunodeficiency virus patients have been associated with a high prevalence of
51 cervical cancer and also the acquisition of the HPV virus. These observations thus put cervical cancer presence as an AIDS
52 defining disease by the Centre for Disease Control, USA (CDC)[20]. With the advent of antiretroviral therapy, life
53 expectancy of these patients is on the increase bringing to light other co-morbidities such as cervical cancer. It thus
54 becomes imperative that these at risk group should embrace all available primary preventive measures to improve outcome
55 as is evident in the developed countries.

56 It is in the light of these, that this study seeks to determine the awareness of this high risk group to HPV and to ascertain if
57 these group of patients attending the antiretroviral clinic had utilized the recommended best practice of HPV vaccination.

58 Method: A cross sectional prospective study was carried out among 162 women attending the antiretroviral clinic between
59 October 21st and December 21st 2014. A pretested interviewer administered questionnaire was used to obtain information
60 from the patients. The antiretroviral clinic operated from Monday- Friday with an average of forty attendees presenting at

61 the clinic per day. Patients were counseled on the study and consent obtained. Those who refused to give consent to
62 participate in the study were excluded from the study.

63 A total number of 4 participants were recruited per day, selecting one out of 10 patients until the desired sample size was
64 obtained using a simple random technique over the two months period.

65 Ethical approval was obtained from the Ethics committee of the hospital for the study

66 The desired sample size was determined using the Lesh-Fish formula for simple proportions [21] which is $N = \frac{Z^2 Pq}{d^2}$
67 while the $P = 6.9\%(0.069)$ which is the HPV vaccination uptake among adolescent daughters in Lagos Nigeria [22].

68 Allowing an error margin of 0.05 at 95% confidence interval, a minimal sample size of 105 respondents was calculated,
69 however a total of 162 respondents were recruited for this Study.

70 Information on the age, parity, marital status, level of education, occupation of spouse, knowledge of route of transmission,
71 awareness of human papillomavirus and its relationship to cervical cancer was obtained from the respondents. Information
72 on previous vaccination with any available brand of HPV vaccine was also obtained from the respondents. The social
73 classes of the respondents were determined using the Olusanyan et al formulae [23].

74 The information was collated and analyzed using SPSS version 20 software. Results of analysis were presented as
75 frequency tables, percentages and test of significant was done using student t test with $P < 0.05$ considered as significant.

76 Results: The mean age of respondents was 35.17 ± 7.14 years with age range of 21-61 years. 48.8%(80) were in the age
77 group 25-34 years, 42.1%(69) were in the group 35- 44years. Other age distributions are as shown in table 1.

78 The review of parity of respondents showed that 56.7%(93) were para0-1, with the least with parity greater than five
79 4.9%(8).

80 Fifty seven point nine percent (90) of respondents were in the middle social class while 25% were in the high social class
81 with 17.1% (28) in the low socioeconomic class. Concerning marital status 75.6% were married, 17.7%(17) were single
82 while 6.7%(11) were divorced. These are shown in table 1.

83 Table 3 highlights the awareness of HPV, HPV vaccination and uptake of vaccination. Among respondents, 14%(23) were
84 aware of HPV causing cervical cancer, 3.7%(6) had correct knowledge of the route of transmission of HPV while
85 10.4%(17) were aware of the existence of HPV vaccination.

86 Comparing uptake of uptake of HPV vaccination and awareness, 9.8%(16) were aware of HPV vaccination and had not
87 taken previously while 90.2%(148) were not aware of the vaccination and hence did not take it.

88 Fig 1 depicts the reasons for failure of uptake of vaccination as expressed by the respondents: 53.7%(88) mentioned the
89 absence of the vaccine as the reason for lack of use, 24.4%(40) sited fear of side effects, cost was expressed by 19.5%(32)
90 and it been unnecessary in 15.2%(25)

91 Discussion

92 The relationship between human papillomavirus and cervical cancer had been established by several studies and had
93 formed the platform for the development of vaccines as primary level of prevention among the populace. This study
94 revealed an abysmal level of awareness of the link between HPV and cervical cancer among the group of patients surveyed
95 who had been observed to have a higher risk of development of cervical cancer. This is even more worrisome because the
96 stage of engagement of these patients was a tertiary hospital which makes it more glaring that these facilities have not done
97 enough to sensitize the patients about the risk of cervical cancer they are exposed to.

98 Both human immunodeficiency virus and the human papillomavirus have the same route of transmission and the risk
99 factors associated with these diseases are largely similar. Various series have associated HIV as medium for enhancing the
100 acquisition of HPV [24, 25]. In addition to the fact that most of the patients are not aware of HPV, larger proportions were
101 also not knowledgeable of the fact that the HPV virus is sexually transmitted. The lack of understanding of this fact implies
102 that the patients are likely to still engage in risk factors associated with its transmission since most of the patients studied
103 are still in their reproductive age. The prevalence of sexually related disease conditions are largely associated with females
104 of reproductive age [26-28] which makes it imperative the need to reinforce the message to reduce the risk of HPV
105 infection among this group of high risk patients.

106 In a bid, to reducing the scourge of human papillomavirus, the development of variants of vaccines have been used with
107 good outcome in developed countries especially in patients with a higher propensity for development of cervical cancer.
108 The use of this preventive strategy has been poor in females in the developing countries and worst still it was observed that
109 the all patients surveyed had not previously been immunized against the HPV inspite of the fact that about 50% of these
110 women are still in the recommended age group for the vaccination. Despite the fact that about 10.4% of women are aware
111 of the existence of the HPV vaccination, this did not influence the uptake of the vaccine among this group. This implies that
112 there must be other factors that are responsible for the lack of uptake of HPV vaccination.

113 In a review of the possible reasons for the lack of uptake of HPV vaccination, a large number of the respondents blamed
114 lack of availability of the vaccine to be the reason for lack of use. This is a common scenario in most developing countries
115 as a result of the inability of governments to procure the vaccine because of other competing health needs and lack of
116 understanding of the burden of the disease. This is in contrast to what is obtained in developed countries where
117 governments have legislative backing and had incorporated vaccination into their national immunization programmes hence
118 increasing uptake with good outcome [29-32]. The assistance of non-governmental agencies like GAVIS in providing
119 vaccines for low and medium income countries had gone a long way in making available vaccines in such environments
120 [33] even though more need to be done in this direction.

The fear of side effects was sighted as the reason for non use of the vaccine as it is obtainable with most vaccines [34]. This erroneous perception can only be removed by proper public health education which had positively affected uptake in some countries.

Cost was another limiting factor that was identified by less than a fifth of the women studied. The effect of cost on the uptake of vaccinations was highlighted by other series [35,36]. However, this study did not identify cost as a major factor since a look at the sociodemographic parameters of the respondents showed that about 82.9% were of middle and higher socioeconomic class.

Furthermore, most of the women also had socioeconomic support from their spouse since a large proportion of the women were married. There is therefore need to explore the marriage link and extend sensitization of the benefits of vaccinations to their husbands who had been observed to pay an important role in the decision making process of women in this environment [37].

In conclusion, the observation of poor uptake of HPV vaccination among this high risk group may infer a possible increase cervical cancer burden on the health system if stringent measures are not taken to avoid its occurrence. A built in primary/community health care enlightenment programme is urgently needed to improve uptake among other programmes among HIV positive women in Nigeria.

Reference:

1. Anorlu RI. Cervical cancer: the Sub Saharan African perspective. *Reprod Health Matters*.2008. 16(32): 41-49.
2. Okonofua F. Human Papillomavirus vaccination and prevention of cervical cancer in Africa. *Afr J Reprod Health*.2007.11: 7-12.
3. Canadian Task force on preventive health care. Recommendations on screening for cervical cancer.*CMAJ*.2013.185: 35-45
4. Sasieni P, Adams J, Cuzick J. Benefits of cervical cancer screening at different ages: Evidence from the UK audit of screening histories. *Br J cancer*.2003.89: 88-93
5. Awodele O, Ademoye A, Awodele D, Kwashi V, Awodele I, Dolapo D. A study on cervical cancer screening amongst nurses in Lagos University Teaching Hospital, Lagos Nigeria. *J Cancer Education*.2011. 26(3): 497-504.
6. Human Papillomavirus: Genital HPV infection. Facts Sheet. Centre for Disease Control and prevention.2014. Accessed 21st January, 1pm 2015.
7. Bosch FX, Lorinez A, Munoz N, Meijer CJ, Shan KV. The causal relationship between HPV and cervical cancer. *J Clin Pathol*. 2002.55(4): 244-265.
8. Megan C, Mark S, Sholom W, Ana CR, Allan H, Wim Q et al. A prospective study of absolute risk determinants

- 151 of human papillomavirus incidence among young women in Costa Rica. BMC Infect Dis. 2013.13:308-17
- 152 9. Winer RL, Feng Q, Hughes JP, O'Reily S, Kivat NB, Koutsky LA. Risk Factors of female human papillomavirus
- 153 acquisition associated with first male sex partners. J Infect Dis .2008.197:279-282.
- 154 10. Faridi R, Amreen , Khalida K, Muhammed I. Oncogenic potentials of human papillomavirus and its relationship
- 155 with cervical cancer. Virol J. 2011.8:269- 271
- 156 11. Magnet S. HPV immune response to infection and vaccination. Stanley Infectious Agent and Cancer.2010.5: 19-
- 157 25
- 158 12. Safaeian M, Porras C, Schiffman M et al. Epidemiological study of anti- HPV16/18 seropositivity and subsequent
- 159 risk of HPV 16 and 18. J Natl cancer Inst. 2010.102(21): 1653-1662.
- 160 13. Andre FE, Booy R, Bock HL, Clemens J, Datta SK, John TJ et al. Bulletin of world Health Organization.
- 161 Vaccination greatly reduces disease, disability and death and inequality worldwide. 2008.86:81-160
- 162 14. Jannah W, Ernestina C, Deborah W. Human Papillomavirus vaccine implementation in low and middle income
- 163 countries: healthy system experiences and prospect. Vaccine. 2013.31(37): 3811- 3817.
- 164 15. Jannie GE, Amrita M. Global challenges of implementing Human Papillomavirus vaccine. International Journal of
- 165 equity in Health.2011.10: 27-32.
- 166 16. Luciano M, Aldo V. HPV Vaccine: an overview of immune response, clinical protection and new approaches for
- 167 the future. Journal of translational medicine.2010.8: 105-13
- 168 17. Diane MH, Stephen LV. Next generation cancer protection: The Bivalent HPV vaccines for females. ISRN Obstet
- 169 Gynaecol.2011.10.4572204.doi.10.5402/2011/45204
- 170 18. Dimie O, Bolande OM, Onyemelukwe GC. Human Papillomavirus (HPV) infection is associated with HIV – 1
- 171 infection and AIDS in HIV infected adult patients from Zaria, Northern Nigeria. Pan Afr Med J.2013.15:
- 172 38.doi.10.11604/Pamj.2013.15.38.2349.
- 173 19. Veldhuijzen NJ, Snigder PJ, Reiss P, Meijer CJ, Van De Wijgert JA. Factors affecting transmission of mucosal
- 174 human Papillomavirus. Lancet Infect Dis.2010.10: 862-874.
- 175 20. Centre for Disease control and prevention. Revised classification system for HIV infection and
- 176 expanded Case Surveillance definition for AIDS among Adolescence and Adults.1993.MWWR. 14:1-
- 177 19
- 178 21. Jekel JF, Katz DL, Elmore JF. Sample size, randomization and probability theory. Epidemiology,
- 179 biostatistics and preventive medicine. 2nd edition Philadelphia. WB Saunders 2001: 196-204.
- 180 22. Beatrice NZ, Mobalanle RB, Ifeoma PO. Mother's Human Papillomavirus Knowledge and willingness to

- 181 vaccinate their adolescent daughters in Lagos, Nigeria. *Int J Women Health*.2013.5.371-377.
- 182 23. Olusanya BO, Okpere E, Ezimokhai M. The importance of social class in voluntary fertility control in a
183 developing country. *West Afr J Med*. 1985.4(1): 205-12
- 184 24. Massad LS, Ahdieh L, Benning et al. Evaluation of cervical cytology among women with HIV 1; evidence from
185 surveillance cytology in the women interagency HIV study. *J Acquired immunodeficiency syndromes*.200. 27(5):
186 432-442.
- 187 25. Sun X-W, Kuhn L, Ellerbrock TV, Chiasson MA, Bush TJ, Wright TC. Human Papillomavirus infection in
188 women infected with the human immunodeficiency virus. *New England J Med*.1997. 337(10).1343-49.
- 189 26. Cotton SC, Sharp L, Masson LT, Little J, Seth R, Neal K et al. lifestyle and sociodemographic factors associated
190 with high risk HPV infection in UK women. *Br J Cancer*. 2007.97(1): 133-139.
- 191 27. Sichanh C, Quet F, Chanthavilay P, Diendere J, Latthaphasavang V, Longquet C, Buisson Y. Knowledge
192 awareness and attitudes about cervical cancer among women attending in an HIV treatment centre in Laos PDR.
193 *BMC Cancer*. 2014.14:161-70.
- 194 28. Sally A, Anthony S, Maryam M, Ayotunde F, Olaniyan O, Offiong R,Wheeler CM. Adebamow C. HIV associated
195 high risk HPV infection among Nigerian women. *BMC infect Dis*.2013.13: 521-26.
- 196 29. Gelrian A, Nikolajski C, Schwarz E, Borrero S. Racial Disparity in awareness of human Papillomavirus. *J*
197 *Women's Health*. 2011.20(8): 1165-1173.
- 198 30. Podolsky R, Gremer M, Atrio J, Hochman T, Arslan AA. HPV vaccine acceptability by Latino Parents: A
199 comparison of Salvadoran Populations. *J Pediatr Adolesc Gynaecol and US*. 2009. 22: 205-215.
- 200 31. Perlman S, Wamai R, Bain P, Welty T, Welty E, Ogembo J. Knowledge and Awareness of HPV vaccine and
201 acceptability to vaccinate in Sub-Saharan African: A systematic Review. Hozbor D (ed). *PLOS One* .2014.9(3):
202 90912.
- 203 32. Kobetz E, Manard J, Hazan G, Sengul T, Joseph T, Nissan J et al. Perception of HPV and Cervical Cancer among
204 Haitian immigrants women. Implication for vaccine acceptability. *Education for Health*.2011. 24(3): 479
- 205 33. Susan AC.A long and winding road: Getting the HPV vaccine in the developing world. *Guttmacher Policy*
206 *Review*. Summer.2007.10.3
- 207 34. Deborah WJ, Keith T, Pieter R, Kathy B, Riziki P, Selephina S et al. Reasons for receiving for not receiving HPV
208 vaccination in primary school girls in Tanzania. A case control study. *PLOS ONE*.2012. 7(10) e45231: 1-10
- 209 35. Ugwu EO, Obi SN, Ezechukwu PC, Okafor II, Ugwu AO. Acceptability of human Papillomavirus vaccine and
210 cervical cancer screening among female health care workers in Enugu, South East Nigeria. *Niger J Clin*

211 Pract.2013.16: 249-52.

212 36. Odetola TD, Ekpo K. Nigerian Women's Perceptions about human papillomavirus immunization. J Community

213 Med Health Edu. 2012.2(11): 1-5.

214 37. Mustapha CD, Ismaila ZM. Male knowledge, Attitudes and Family Planning Practices in Northern Nigeria. Afri J

215 Reprod Health. 2006.10(3): 53-65.

216 Table 1: Sociodemographic characteristics

217

Variables	Frequency	Percentage
Age		
15-24	10	6.1
25-34	80	48.8
35-44	69	42.1
45-54	3	1.8
55-64	2	1.2
Parity		
0-1	93	56.7
2-3	47	28.7
4-5	16	9.7
>5	8	4.9
Social status		
Low	28	17.1
Middle	95	57.9
High	41	25.0
Marital Status		
Single	29	17.7
Married	124	75.6
Divorced	11	6.7

Table 2: Awareness of HPV, knowledge of HPV vaccination and uptake of HPV vaccination

Variables	Frequency	
	Yes n(%)	No n(%)
Awareness of HPV causing cancer	23(14.0)	141(86.0)
Transmission route of HPV	6(3.7)	158(96.3)
Awareness of HPV vaccination	17(10.4)	147(89.6)

Comparing awareness of HPV causing cancer against awareness of HPV vaccination (OD1.41 P=0.1999;CL 0.72-2.75)

Fig 1: Reasons for failure of uptake of vaccination




Table 3 Awareness of HPV and previous uptake of HPV vaccination

	Uptake of HPV vaccination		Total n(%)
	No n(%)	Yes n(%)	
Awareness of HPV vaccination			
Yes	16(9.8)	0(0.0)	16(9.8)
NO	148(90.2)	0(0.0)	148

Uptake of HPV was not affected by awareness of the HPV vaccination. (P=0.90)