

Original Research Article

Soil-Transmitted Helminth Infections and Associated Risk Factors in a Neglected Region in the Upper Nkongho-mbo Area, South-west Region, Cameroon.

ABSTRACT

Introduction

Soil-transmitted helminth (STH) infections are among the most prevalent chronic human infections worldwide and affect the poorest and most deprived communities. Few studies have been carried out to evaluate the state of STH infections in the Upper Nkongho-Mbo region of Cameroon; hence levels of infection remain unknown.

Methodology

A cross-sectional study on the prevalence and intensity of soil-transmitted helminth infections as well as the influence of risk factors was carried out in the Upper Nkongho-Mbo area, South-West Region, Cameroon, between November 2012 and July 2013. Faecal and soil samples were collected and analysed using the formol-ether concentration technique and the sucrose floatation centrifugation technique respectively. Information on hygienic standards was also obtained.

Results

Results obtained showed that out of 327 people sampled, 145 were infected with one or more STHs giving an overall prevalence of 44.34%. The prevalence of round worm infection (39.14%, 128) was significantly higher ($\chi^2 = 26.95$; $p \leq 0.001$) than that of whipworm (8.26%, 27) and hookworm (13.46%, 44). The prevalence of infection varied with age group ($p \leq 0.001$). The 21-50 years age group had the highest prevalence (55.44%) while the 6-12 years age group had the lowest prevalence (30.38%). Intensity of infection also varied significantly with age groups ($p < 0.001$). The highest intensity was observed in the 21-50 years age group (113.5 ± 24.8), and the lowest in the 6-12 years age group (20.95 ± 9.33).

26 Soil samples examined had a prevalence of 26.79% (30) compared with 44.34% observed in
27 the faecal samples. The proportion of soils contaminated was statistically different from the
28 proportion of faecal samples contaminated ($\chi^2 = 4.307$; $p = 0.038$). Age and toilet type were
29 found to be important predictors of infection ($p = 0.041$ and $p = 0.01$).

30 **Conclusion**

31 These findings provide evidence for the high risk of acquiring STH infections from the
32 middle age group in the Upper Nkongho-Mbo region of Cameroon. Effective measures are
33 therefore necessary to reduce contamination of the region.

34 **KEY WORDS**

35 Soil transmitted helminthes, Co-infection, Prevalence, Cameroon.

36 **INTRODUCTION**

37 Thousands of rural and impoverished villagers throughout the tropics and subtropics are often
38 chronically infected with several different species of parasitic worms [1] About 2 billion
39 people harbour these infections worldwide, of which 300 million suffer associated severe
40 morbidity and even death. The World Health Organization estimates indicate that
41 geohelminthiasis account for more than 40% of the disease burden due to all tropical diseases
42 [2,3,4]. In developing countries, where control measures are often difficult to implement,
43 STHs remain a significant health problem. Approximately one third of the world's population
44 is infected with at least one species of STH, with *Ascaris lumbricoides* infecting 800 million
45 people, *Trichuris trichiura* 600 million, hookworm 600 million and resulting in up to 135,000
46 deaths annually [5]. STHs have also been classified among the most prevalent neglected
47 tropical diseases as they persist exclusively in the poorest populations and thus are often
48 forgotten [6]. Important contextual determinants for human infection are poverty, lack of
49 sanitation and inadequate hygiene [7,8] Efforts at controlling STHs are principally directed
50 at school children because they are at the greatest risk of morbidity [9]. This may not be the
51 best control strategy, and may not achieve the goal of eradication by 2020. Adults have been

known to have a higher infection rate than school children and to harbour a greater variety of intestinal helminths [10]. Engagement in agricultural practices puts adults at a greater risk of infection with hookworm [11]. Thus, in many communities adults constitute an important reservoir of infection which is largely neglected by most current control programs. Due to a high rate of re-infection, despite long periods of intervention, it has been found that many communities that are subjected to routine targeted deworming still maintain a low level of infection [12]. In Cameroon, STH infections are endemic with rural areas being more infected than urban areas [13,14,15]. Despite more than a decade of serious intervention following the MDAI (combination therapy with albendazole and ivermectin) targeted chemotherapy protocol [16]. STH infections persist in many rural areas. As a result, it can be concluded that STH infections contribute significantly to the vulnerability of our rural populations. This study was conducted to determine the prevalence and intensity of STHs in the Upper Nkongho-Mbo area and also identify, in real time, the role of risk factors in transmission in the study area.

2. MATERIALS AND METHODS

2.1 Study population and design

The study was carried out between November 2012 and July 2013 in the UpperNkongho-Mbo area. This area is found in the Kupe-Manenguba division of the South-West region of Cameroon. Faecal samples were collected from participants in three villages using clean dry wide-mouth specimen bottles with tight covers. This was a cross sectional study where sample collection was selectively purposive, since the study aimed at targeting a wide age range within the defined study area [17]. Equally, 112 soil samples were collected from toilet areas, community playing grounds and school yards (200gms of top soil each).

76 **Ethical considerations**

77 This study was carried out with the approval of the Ethical Review Committee on Health
78 Research, regional delegation of health for the South West Region Buea, Cameroon.
79 Informed written consent was obtained from each study participant. Each participant was
80 free to withdraw consent at any time. All personal information of the participants was treated
81 strictly confidential.

83 **Sample size and sampling techniques**

84 Fresh faecal samples were collected from 327 participants during the study period using
85 sterile dry, clean, air tight, wide mouth screw cap labelled containers and analyzed within 24
86 hours of collection, following standard procedures. Stool samples were analyzed using the
87 formal-ether concentration method while soil samples were analysed using the sucrose
88 floatation-centrifugation technique [18, 19]. Equally a semi-structured questionnaire was
89 administered to all the 327 participants to elicited demographic information, toilet type,
90 safety of water source, social determinants, health seeking behaviours and health education.

91 **Data Analysis**

92 Data was entered into Microsoft Excel 2007 and analysed using Minitab version 16.
93 Descriptive analysis was performed on the data and the results presented in tables and graphs.
94 Statistical tests were carried out at 95% confidence interval ($\alpha= 0.05$).The data was classified
95 by age, sex and village for statistical comparison. The data was tested for normality using the
96 Kolmogorov-Smirnov test for normality and counts were found to be not normally
97 distributed. Mean parasite intensities between sites and between groups were compared using
98 non-parametric Kruskal-Wallis test (H) or the Mann-Whitney U test (W). Chi square
99 Goodness-of-Fit test was used to compare proportions. Binary logistic regression was used to
100 test the determinants of infection.

101

3.0 RESULTS

3.1 Characteristics of the study population

A total of 327 participants provided samples for the study. This included 164 males (50.15%) and 163 females (49.85%). Out of the three villages included in the study, 91 respondents were from Lebok, 49 from Mbefe, and 187 from Njungo (Table 1). Participants were distributed into five principal age groups broken down as follows: 0-5 years, 6-12 years, 13-20 years, 21-50 years and >50 years. The proportion of each age group is presented in Table 1.

3.2 Prevalence and intensity of infection with STHs in the study area

Out of the 327 stool samples examined, 145 (44.34%) of them were infected with one or more species of helminth ova shown in Plate 1. Overall prevalence of infection was similar ($P > 0.05$) in Mbefe (46.94%), Njungo (45.46%) and Lebok (40.66%). However, the prevalence of whip worm varied significantly from village to village ($\chi^2 = 8.95$, $P < 0.05$), being highest in Mbefe (14.29%) and lowest in Lebok (2.2%) as shown in Table 2. Prevalence of roundworms was significantly higher than whipworm and hookworm ($\chi^2 = 26.95$, $P < 0.01$). This overall pattern was consistent in all three villages in the study area (Table 2).

Overall prevalence of co-infections with two or more STHs was 15.0% (Table 3). Co-infection was most common in Njungo (16.6%) than in Lebok and Mbefe. Co-infection with roundworm and hookworm showed the highest prevalence (7.7%) while the lowest was with whipworm and hookworm (0.9%) as shown in Table 3.




Plate 1: Eggs of soil-transmitted helminths identified during the parasitological survey. A and B, *A. lumbricoides*, one cell stage; C, *A. lumbricoides*, blastula; D and E, *T. trichiura*; F, hookworm.

Table 1: Demographic characteristics of the study population

Variable		Villages			Total
		Lebok	Mbefe	Njungo	N (%)
		N (%)	N (%)	N (%)	
Sex	Male	52 (15.9)	21 (6.42)	91 (27.83)	164 (50.15)
	Female	39 (11.93)	28 (8.56)	96 (29.36)	163 (49.85)
Age groups in Years	0-5	4 (4.4)	3 (6.12)	27 (14.44)	34 (10.4)
	6-12	53 (58.24)	12 (24.49)	39 (20.89)	101 (31.8)
	13-20	4 (4.4)	10 (20.41)	35 (18.72)	49 (14.99)
	21-50	18 (19.78)	13(26.53)	61 (32.62)	102 (31.19)
	>50	12 (15.9)	11 (22.45)	25 (13.37)	58 (17.75)

Total 91 (27.83) 49 (14.99) 187 (57.19) 327 (100)

Table 2: Prevalence of soil-transmitted helminth infections among subjects in three principal communities in the study area

Prevalence (%)							
Village	N	Roundworm % (N)	whipworm % (N)	Hookworm% (N)	Overall prevalence % (N)	χ^2	P-value
Lebok	91	37.36 (34)	2.2 (2)	10.99 (10)	40.66 (37)	39.74	< 0.001
Mbefe	49	44.9 (22)	14.29 (7)	8.16 (4)	46.94 (23)	34.51	< 0.001
Njungo	187	39.57 (74)	8.02 (15)	15.51 (29)	45.46 (85)	25.84	< 0.001
Total	327	39.14 (128)	8.26 (27)	13.46 (44)	44.34 (145)	26.95	< 0.001
χ^2		0.74	8.95	2.38	0.49		
P-value		0.69	0.011	0.30	0.784		

(Chi-square Goodness-of-Fit Test; χ^2 = Chi-square statistic)

138 Table 3: Prevalence (%) of co-infections in the study area

Village	Prevalence (%); (Number infected)					χ^2	P-value
	round worm and whipworm	Roundworm and hookworm	whipworm and hookworm	All three STHs	Total		
Lebok	1.1 (1)	6.6 (6)	0 (0)	1.1 (1)	8.8 (8)	12.1	0.007
Mbefe	8.2 (4)	4.1 (2)	0 (0)	4.1 (2)	16.3 (8)	8.2	0.04
Njungo	4.8 (9)	9.1 (17)	1.6 (3)	1.07 (2)	16.6 (31)	9.9	0.02
Total	4.3 (14)	7.7 (25)	0.9 (3)	1.5 (5)	15.0 (47)	8.1	0.05
χ^2	5.366	1.894	3.2	2.8998	2.8101		
P-value	0.07	0.39	0.202	0.24	0.25		

139

140 3.3. Intensity of infection with soil-transmitted helminths in the study area

141 Overall, roundworm had the highest intensity of infection in the study area with mean worm
 142 burden being 59.9 ± 10.0 epg while whipworm had the lowest intensity (3.07 ± 0.91 epg).
 143 While intensity of infection with roundworm and hookworm did not differ between the
 144 communities, intensity of infection with whipworm was statistically different with intensity
 145 in Njungo being higher than in Lebok (Table 4).

146 Table 4: Intensity of infection with STH in the different villages

Village	Intensity of infection (epg $\pm$ SE)		
	Roundworm	Whipworm	Hookworm
Lebok	34.6 $\pm$ 15.3	1.29 $\pm$ 1.16	4.35 $\pm$ 2.47
Mbefe	56.1 $\pm$ 20.2	2.12 $\pm$ 0.91	3.10 $\pm$ 1.64
Njungo	73.2 $\pm$ 14.9	4.19 $\pm$ 1.46	3.87 $\pm$ 0.89

Total	59.9 ±10.0	3.07 ±0.91	3.87 ±0.89
H	1.67	6.85	2.48
p	0.433	0.032	0.289

(Kruskal-Wallis test; H = Kruskal-Wallis statistic)

Roundworms remained the most implicated worm type. However Mann-Whitney test showed that the intensity of infection with STHs was roughly comparable among males and females (Table 5). In contrast to trends observed in Lebok and Njungo, women tended to harbour higher worm burdens than men in Mbefe.

Table 5 Sex-related intensity of infection with soil-transmitted helminths in the study area

	Mean intensity of infection (epg) ±SE					
	Roundworm		Whipworm		Hookworm	
	Males	Females	Males	Females	Males	Females
Lebok	51.3 ±	14.91 ±	2.25 ±	0.000 ±	5.31 ±	2.88 ±
	26.5	4.44	2.03	0.000	4.09	1.73
Mbefe	41.6 ±	67.0 ±	0.33 ±	3.46 ± 1.54	2.95 ±	3.21 ±
	14.5	33.8	0.33		2.17	2.40
Njungo	73.1 ±	73 ± 19.7	5.81 ±	2.66 ± 1.13	3.58 ±	4.15 ±
	22.5		2.76		1.09	1.41
Significance	W = 26849.5, p =			W = 26758.0, p =		
	0.88			0.98		

(Mann-Whitney test; W = Mann-Whitney statistic).

Overall, the 21-50 years age group appeared to have the highest prevalence of infection (56.5%) followed by the 13-20 years age group (51.02%). These differences however were

not statistically significant (Table 6). In all three villages, prevalence of STH infection differed significantly with respect to age. Apart from the 21-50years age group, the prevalence of infection with STHs differed significantly for each age group across the three villages. In addition, apart from the 21-50 years age group where the prevalence was similar in all the villages, each age group differed significantly from village to village.

163

Table 6: Age-related Prevalence of STH infections in the study area

Number infected (%) in each age group								
Village	N	0-5	6-12	13-20	21-50	>50	χ^2	P-value
Lebok	91	50 (2)	26.4 (13)	75 (3)	55.6 (10)	66.7 (8)	25.2077	<0.001
Mbefe	49	100 (3)	58.3 (7)	60 (6)	38.5 (5)	18.18 (2)	67.0749	<0.001
Njungo	187	40.7 (11)	28.2 (10)	45.7 (16)	60.7 (37)	40 (10)	12.8633	0.01
Total	327	31.2 (14)	28.9 (30)	53.1 (26)	59.8 (55)	41.7 (20)	8.515	0.07
χ^2		32.003	17.067	7.1277	5.2407	28.373		
P-value		< 0.001	< 0.001	0.03	0.07	<0.001		

165

Out of 112 soil samples, 30 (26.79%) were infected with one or more species of STHs. The most common STH in the soil was roundworm (14.29%) while the least was whipworm (5.36%). Although not statistically significant, soil samples from Mbefe were most infected (33.33%) when compared with the infection rate of soils in Lebok (31.11%) and Njungo (18.61%) (Table 7).

170

Table 7: Rate of soil contamination with STHs in the Upper Nkongho-Mbo area

Village	Prevalence (%)			Total	χ^2	P-value
	Roundworm	Whipworm	Hookworm			
Lebok	15.56 (7)	6.67 (3)	13.33 (6)	31.11 (14)	3.610	0.164
Mbefe	16.67 (4)	8.33 (2)	16.67 (4)	33.33 (8)	3.338	0.188
Njungo	11.63 (5)	2.33 (1)	9.30 (4)	18.61 (8)	6.040	0.049
Total	14.29 (16)	5.36 (6)	12.5 (14)	26.79(30)	4.166	
χ^2	0.959	3.323	2.079	4.550		
P-value	0.619	0.190	0.354	0.103		

3.4 Predictors of soil-transmitted helminth infections.

Binary logistic regression between STH infection and a number of determinants revealed that age and toilet type were significant predictors of STH infections ($p = 0.041$ and $p = 0.01$ respectively). Soil-transmitted helminth infections tended to increase as age increased; while the use of the jetty toilet increased the risk of infection with STHs. Possession of a pit toilet was negatively associated with STH infection (table 8).

Table 8: Predictors of STH infection

Variable		Odds ratio	95% CI		P-value
STH	Coefficient		Lower	Upper	
Sex	- 0.108999	0.9	0.57	1.4	0.634
Age	0.0106227	1.01	1.00	1.06	0.041
Toilet	- 0.682883	0.55	0.3	0.85	0.01

4. DISCUSSION

The prevalence and intensity of intestinal helminth infections among different populations are functions of many different factors, most importantly environmental factors, parasitic factors, and host factors. The study revealed an overall prevalence of 44.04% for STHs. Prevalence rates were higher than those reported in other parts of Cameroon [20, 21, 22]. However, other researchers found much higher prevalence rates among the Orang Asli in rural Malasia [23], although this study was not restricted to children. The prevalence of STH infections was similar in Lebok, Mbefe and Njungo, probably because the villages are ecologically similar.

Ascaris infections occurred at higher prevalence than whipworm and hookworm infections, consistent with findings of Ntonifor, et al. [8]. This is probably because ecological determinants such as humidity and altitude favour the development of Ascaris spp while hampering the development of whipworm. Hookworm infections were more prevalent than infections with whipworms. Findings in Kotto_Barombi and Marumba II villages similarly indicated higher prevalence of whipworm than hookworm [13]. The higher prevalence of hookworm noted in this study is probably because hookworm infections usually increase during the rainy season and also it is a typical rural farming population where most of them work bare footed in their farmlands. In addition, whipworm is less common at high altitudes [24].

Intensity of infection was generally low, consistent with the findings of Allen et al. [12] who explained that in regions where STHs are targeted for elimination with periodic deworming, high worm burdens are not very common. The data showed extreme variability supporting the fact that STHs are typically over-dispersed in the population with a few individuals harbouring infections of high intensity. Roundworm occurred at the highest intensities in the region while whipworm occurred at the lowest intensities. These trends are consistent with the findings of other researchers [10, 15].

210 Despite initiation and completion of the annual deworming campaign shortly after the start of
 211 these studies, the prevalence of infection in the targeted age group remained high. This is
 212 roughly comparable to the findings of Ndamukong et al [25]. .Prevalence of infection was
 213 significantly higher in Mbefe than in Lebok and Njungo in these age groups. This is probably
 214 because the samples of Mbefe were collected before the anthelmintic campaign began and so
 215 these children had not received any anthelminthics. Such a high prevalence of infection is
 216 consistent with the results of other reseachers [26]. The most affected age group was the 21-
 217 50 years age group consistent with the findings of Pinheiro, et al. [27] and Ntonifor et al., [8].
 218 These findings are probable indicators that adults in the community are more important
 219 epidemiologically than school-aged children. This can be because this is the most active age
 220 group, daily involved in agriculture and outdoor activities that put them at greater risk of
 221 infection with STHs. Unfortunately, this group is largely ignored by the deworming exercise
 222 which is targeted at children through schools. Thus infection with STHs would tend to
 223 amplify in this age group. Such tendencies would account at least in part for the differences
 224 in infection prevalence observed between age groups. Such heavily infected age groups
 225 would equally impact environmental contamination with ova considerably. In addition, the
 226 deworming programmes target only children therefore leaving the adults as the main sources
 227 of contamination.

228 The overall rate of soil contamination (26.79%) was similar to the findings of Cassenote, et
 229 al. [28] but much less than that reported by Horiuchi, et al. [29]. Soil analysis revealed that
 230 the soils were seeded with the various ova of these worms. This moderate rate of Soil
 231 contamination is probably due to the use of the jetty toilet. Roundworms were more prevalent
 232 in soil samples than hookworm and whipworm, even though soil contamination with
 233 hookworm had the highest mean intensities. This is probably because hookworm
 234 development peaks during the rainy season [30]. This seasonal surge might not necessarily

affect the distribution of soil-transmitted helminth infections in the long run. Although prevalence of STHs was higher in faecal samples than in the soil, the pattern of distribution of soil-transmitted helminth infections in the soil was identical to the pattern of infection in study subjects. This suggests that the soil plays a major role in the epidemiology of soil-transmitted helminth infections in the Upper Nkongho-Mbo area. However, the generally low intensities might point to the cultural use of some plants in foods and drinks in the study area. It is possible that the consumption of menag (a bitter-sweet liquor) garnished with *Anona senegalensis* (locally called ‘bitter’), and the use of *Piper guineense* (bush pepper) as a spice in most of their foods helps to reduce worm burdens. These plants have been shown to have anthelmintic activities [31, 32]. This might affect worm fecundity and survival in the human host and thus reduce faecal egg counts. However, the possible anthelmintic activity of menag garnished with “bitter” needs to be investigated further. Age and toilet type were identified as important predictors of infection. Older people tended to be more at risk of becoming infected with STHs. The use of the jetty increased the risk of becoming infected probably because this helped to seed the surroundings with infective helminth eggs. Contamination of the environment with infective stages of STHs is considerable in the absence of adequate sanitation [5, 33].

1. WHO. Global School-based Student Health Survey: Honduras. 2014 Fact Sheet. 2 p.
2. Hotez PJ, Molyneu, DH, Fenwick A., Kumaresan J., Sonia PH., et al. Review; Current concepts: control of neglected tropical diseases. *New England Journal of Medicine* 2007 **357**: 1018-1027.
3. WHO. Monitoring anthelmintic efficacy for soil-transmitted helminths; Report of WHO expert committee; Geneva, 2008a

- 259 4. WHO. Public health significance of intestinal parasitic infections. Bulletin of WHO;
260 2008b **65** (5): 575 - 588.
- 261 5. Hotez, P. J. One world health: Neglected Tropical Diseases in a Flat World. PLoS
262 Negl Trop Dis. 2009, 3, 405
- 263 6. Liese B., Rosenberg M. and Schratz A. Programmes, partnerships and governance for
264 elimination and control of neglected tropical diseases. Lancet. 2010, 375, 67–76
- 265 7. Vandemark LM., Jia TW. and Zhou XN. Social science implications for control of
266 helminth infections in Southeast Asia. Adv Parasitol. 2010, 73, 137–170
- 267 8. Ntonifor HN, Green AE, Bobda MOS, and Ebot T.J. Epidemiology of Urinary
268 Schistosomiasis and Soil transmitted Helinthiasis in a recently established focus
269 behind Mount Cameroon. Int. J. Current microbial. app. Scs. 2015, 4(3) :1056-1066.
- 270 9. De Silva NR., Brooker S., Hotez PJ., Montessor A., Engels D., and Savioli L. Soil-
271 transmitted helminth infections: updating the global picture. Trends in Parasitol; 2003,
272 62: 221-261
- 273 10. Anantaphruti M.T., Waikagul J., Maipanich W., Nuamtanong S. and Pubampen S.
274 Soil-transmitted helminthiasis and health behaviours among school children and
275 community members in a west-central border area of Thailand. Southeast Asian J of
276 Trop Med Pub Hlth; 2004, **35** (2): 260-266.
- 277 11. Brooker S., Whawell S, Kabatereine N.B., Fenwick A. and Anderson R.M.
278 Evaluating the epidemiological impact of national control programs for helminths.
279 Trends in Parasitol y;2004, **11**: 537-545.
- 280 12. Allen H, Sithey G., Padmasiri E.A. and Montresor A. Epidemiology of soil-
281 transmitted helminths in the western region of Bhutan, . Southeast Asian J of Tropil
282 Med Pub Hlth;; 2004, **35** (4): 777-779.
- 283 13. Nkengazong, L., Njiokou, F; Wanji, S., Teukeng, F., Enyong, P., et al. Prevalence of
284 soil-transmitted helminths and impact of albendazole on parasitic indices in
285 KottoBarombi and Marumba II villages (South-West Cameroon. .Af J of Envttal Sc
286 and Tech.2010, 4 (3): 115-121.

- 287 14. Kimbi H.K., Lum E., Wanji S., Mbuh J.V., Ndamukong-Nyanga J.L., Eyong EEJ. and
288 Lello J. Co-infections of asymptomatic malaria and soil-transmitted helminths in
289 school children in localities with different levels of urbanization in the mount
290 Cameroon region. *Journal of Bacteriology and Parasitology*; 2012, 3: 134
- 291 15. Mbuh JV., Ntonifor HN. and Ojong J. The epidemiology of soil-transmitted helminth
292 and protozoan infections in South-West Cameroon. *Journal of Helminthology*; 2012
293 86 (01): 30 - 37.
- 294 16. WHO. First WHO report on NTDs 2010: working to overcome the global impact of
295 NTDs. 2010. Geneva.
- 296 17. Teddlie C., Yu F. .Mixed methods of sampling: a typology with examples. *Journal of*
297 *mixed methods research*; 2007, **1**(1): 77 - 100.
- 298 18. Uga S., Matsumira T., Aoki N. and Kataoka N. Prevalence of Toxocara eggs in the
299 sandpits of public parks in Hyogo Prefecture, Japan. *Japanese J. Parasitol.* 1989, 38:
300 280 - 284.
- 301 19. Rai S.K., Uga S., Ono K., Rai G. and Matsumura T. Contamination of the soil with
302 helminth parasite eggs in Nepal; 2000, **31** (2):
- 303 20. Nkuo-Akenji T.K., Chi P.C., Cho J.F., Ndamukong K.J. and Sumbele I. Malaria and
304 helminth co-infection in children living in a malaria endemic setting of mount
305 Cameroon and predictors of anaemia. *J. Parasitol.*; 2006, 92 (6):1191-1195.
- 306 21. Makoge VD., Mbah G.A., Nkengazong L., Sahfe N.E., and Moyou R.S.. *Falciparum*
307 malaria, helminth infection and anaemia in asymptomatic pupils in four villages in
308 Cameroon .*European J. Zool. Res* ;2012, **1** (2): 54-59.
- 309 22. Tchuenté L.AT., Noumedem C.D., Ngassam P., Kenfack C.M., Gipwe N.F et al.
310 .Mapping of schistosomiasis and soil-transmitted helminthiasis in the regions of

- 311 Littoral, North-West, South and South-West, Cameroon and recommended treatment.
312 BMC Infectious Diseases;2013, 13 602.
- 313 23. Nasr N.A., Al-Mekhlafi H.M., Roslan M.A., and Bulgiba A. Towards an effective
314 control program of soil-transmitted helminth infections among Orang Asli in rural
315 Malasia. Part 1: prevalence and associated risk factors. Parasites and Vectors; 2013, 6:
316 27-33
- 317 24. Chammartin F., Scholte R.G.C, Malone J.B., Bavia M.E., Nieto P., Utzinger J. and
318 Vounatso P. Modelling the geographical distribution of soil-transmitted helminth
319 infections in Bolivia.. Parasites and Vectors;2013, 6: 152.
- 320 25. Ndamukong K.J.N. Epidemiology of intestinal nematodes in school age children of
321 the kumba urban area, South West, Cameroon. East Af Medl J.2005, 82 (11): 559-
322 564.
- 323 26. Egwunyenga O. and Ataikiru D.P. Soil-transmitted helminthiasis among school age
324 children in Ethiope East Local Government Area, Delta State, Nigeria. Af. J. Biotech;
325 2005, 4 (9): 938-941.
- 326 27. Pinheiro I.O., Castro M.F., Mitterothe A., Pires F.A.C., Abramo C., et al. Prevalence
327 and risk factors for giardiasis and soil-transmitted helminthiasis in three
328 municipalities of Southeastern Minas Gerais state, Brazil. Parasitology Research;
329 2011, 108: 1123 - 1130.
- 330 28. Cassenote A. J., Pinto-Neto J.M., Lima-Catelam A.R. and Ferriera A.W. .Soil
331 contamination by eggs of soil-transmitted helminths with zoonotic potential in the
332 town of Fernandópolis, state of Sao Paulo, Brazil, between 2007 and 2008, .Revista
333 da SociedadBrasileira de Medicina Tropical; 2011, 44 (3): 371-374
- 334 29. Horiuchi S., Paller V.G. and Uga S. Soil contamination by parasite eggs in a rural
335 village in the Philippines.Tropical Biomed; 2013, 30 (3): 495-503.

- 336 30. Schad G.A. Review of recent advances in knowledge of biology of human hookworm
337 infection. Scientific group on intestinal protozoan and helminthic infections.WHO;
338 Geneva, Switzerland. 1980.
- 339 31. Alawa C.B., Adamu A.M., Gefu J.O., Abdu P.A., Chiezey N.P., Alawa J.N. and
340 Bowmann D.D. Invitro screening of two Nigerian medicinal plants
341 (*Vernoniaamygdalina* and *Annonasenegalensis*) for anthelmintic activity. *Veterinary*
342 *Parasitology*; 2013, 113 (1): 73-81.
- 343 32. Ekanem A.P., Wang M., Simon,J.E., Obiekezie A.I. and Morah,F.).Invivo and invitro
344 activities of the seed extracts of *Piper guineense*Schuum. andThonn. against skin and
345 gill monogenean parasites of goldfish (*Carassiusauratusauratus*).
346 *Phytotherapyresearch*; 2014, 18 (10): 793-797.
- 347 33. Ziegelbauer K., Speich B., Mausezahl D., Bos R., Keiser J., and Utzinger J. Effect of
348 sanitation on soil-transmitted helminth infection: systematic review and meta-
349 analysis. *PLOS Medicine*; 2009, 9: e1001162.